



2014 Analytic Report

**Sponsoring Washington State Agencies:
Department of Social and Health Services'
Division of Behavioral Health and Recovery
Office of Superintendent of Public Instruction
Department of Health
Liquor and Cannabis Board
Department of Commerce**

Prepared by:
Looking Glass Analytics, Inc.

April 2016

Washington State Healthy Youth Survey 2014

Analytic Report

Department of Social and Health Services

1115 Washington St. S.E.
P.O. Box 45000
Olympia, WA 98504-5000

Department of Health

Town Center East
111 Israel Road S.E.
Tumwater, WA 98501-7835

Office of Superintendent of Public Instruction

Old Capitol Building
600 S. Washington
P.O. Box 47200
Olympia, WA 98504-7200

Liquor and Cannabis Board

3000 Pacific Ave. S.E.
P.O. Box 43075
Olympia, WA 98504-3075

Prepared by:

Looking Glass Analytics, Inc.

215 Legion Way SW
Olympia, WA 98501

April 2016

In collaboration with Anar Shah, Lillian Bensley, Martha Perla, Grace Hong, Jennifer Sabel,
and members of the Healthy Youth Survey Planning Committee

This report is available online at:
<http://www.AskHYS.net>

Suggested Citation:

Healthy Youth Survey 2014 Analytic Report. Washington State Department of Social and Health Services,
Department of Health, Office of the Superintendent of Public Instruction, and Liquor and Cannabis Board, April
2016

Data from the 2014 Washington State Healthy Youth Survey and previous administration of youth surveys in
Washington were used in this publication.

The Healthy Youth Survey was administered by the Washington State Department of Health, the Office of the
Superintendent of Public Instruction, the Department of Social and Health Services, and the Liquor and Cannabis
Board. The Healthy Youth Survey Planning Committee included members of these state agencies to oversee the
implementation of the 2014 survey.

Federal funding for this report was provided by the Substance Abuse Prevention and Treatment Block
Grant and the Community Mental Health Services Block Grant

Federal funding for the survey was provided by the U.S. Center for Substance Abuse Prevention, the Maternal and
Child Health Bureau, and the Centers for Disease Control and Prevention. Washington State support was provided
by the Department of Health - Division of Prevention and Community Health; the Office of Superintendent of
Public Instruction, the Department of Social and Health Services - Division of Behavioral Health and Recovery, the
Department of Commerce, the Family Policy Council and the Liquor and Cannabis Board. This report is supported
in part by the Maternal and Child Health Block Grant (B04MC08890) through the Department of Health and Human
Services.

Prepared for:

Department of Social and Health Services

Kevin Quigley

Secretary, Department of Social and Health Services

Chris Imhoff

Director, Division of Behavior Health and Recovery

Michael Langer

Chief, Office of Behavioral Health and Prevention

Katie Weaver Randall

Chief, Office of Decision Support and Evaluation

Sarah Mariani

Behavioral Health Administrator

Felix I. Rodriguez

Researcher, Office of Decision Support and Evaluation

Grace Hong

Epi Prevention Research Manager, Office of Decision Support and Evaluation

Martha Perla

SEOW Research Manager, Office of Decision Support and Evaluation

Stephen Smothers

Prevention Services Lead

Linda Becker

Prevention Research Manager, Office of Decision Support and Evaluation

Department of Health

John Weisman

Secretary

Kathy Lofy

State Health Officer

Cathy Wasserman

State Epidemiologist for Non-Infectious Conditions

Lillian Bensley

Epidemiologist

Jennifer Sabel

Epidemiologist

Kevin Beck

Survey & Operations Manager

Anar Shah

Epidemiologist

Office of Superintendent of Public Instruction

Randy I. Dorn

State Superintendent of Public Instruction

Ken Kanikeberg

Chief of Staff

Alan Burke

Deputy Superintendent, K-12 Education

Greg Williamson

Director, Learning and Teaching Support

Dixie D. Grunenfelder

Program Supervisor, Learning and Teaching Support

Krissy Johnson

Program Supervisor, Learning and Teaching Support

Liquor and Cannabis Board

Jane Rushford

Board Chair

Rick Garza

Director

Mary Segawa

Public Health Education Liaison

Contents

Acknowledgements.....	i
Executive Summary.....	ii
1. Introduction	1
Organization and Purpose of the Report	3
Participation	3
Cautions.....	4
Representativeness.....	4
Trends	4
Rounding Differences.....	5
School Dropouts.....	5
Developmental Changes	5
Self-Report Data.....	5
Correlational Data.....	6
2. Methods.....	7
Sampling	7
Survey Administration.....	7
Questionnaires	8
Translations	9
Reliability and Validity.....	9
Data Preparation and Analysis	9
Differences by Grade Level and Gender	10
Differences over Time.....	10
Chart Formatting.....	11
Calculating Confidence Intervals.....	11
Response Rates.....	13
Non-completion Rates by Form	14
3. Physical Activity and Dietary Behavior.....	18
Obesity and Overweight.....	18
Exercise and Physical Activity	18
60 Minutes of Exercise Daily.....	19
Physical Education Classes.....	20
Time Spent in Physical Education Classes.....	21
Television Watching and Video Game Playing.....	22
Nutrition	23
Fruit and Vegetable Consumption	23
Eating Dinner with Family.....	23
Drinking Sweetened Beverages	25
Drinking Sweetened Drinks at School.....	26
Buying Sweetened Drinks at School.....	27
Food Insecurity	28
4. Health Status and Health Care	29
Asthma.....	29

Lifetime Asthma	29
Current Asthma.....	30
Diabetes	31
Access to Care	32
Access to a Doctor	32
Access to a Dentist.....	33
5. Mental Health	34
Depressive Feelings	34
Suicide	35
Suicide Attempts.....	36
6. Sexual Behavior.....	37
Sexual Behavior	37
Lifetime Sexual Intercourse	37
Sexual Behaviors among Those who have had Sex	38
Sexual Orientation	39
Sex of Sexual Contacts	40
7. School Climate.....	41
School Safety, Bullying, and Harassment	41
Feeling Safe at School	41
Bullying.....	42
Harassment by Computer or Cell Phone	43
Harassment Due to Perceived Sexual Orientation	44
Weapon Carrying at School	45
Substance Use at School.....	46
Alcohol or Other Drug Use on School Property	46
Alcohol Drinking on School Property	47
Marijuana Use on School Property	48
Perceived Availability of School Staff to Discuss Substance-Related Problems	49
School Attendance	50
Skipping or Cutting School	50
Enjoying School	51
8. Unintentional Injury Behaviors	52
Motor Vehicle Safety	52
Riding with a Drinking Driver	52
Drinking and Driving.....	53
Marijuana Use – Riding and Driving.....	54
Texting or Emailing – Riding and Driving	55
Swimming Safety	56
Boat Safety	57
9. Intentional Injury Behaviors.....	58
Physical Fighting	58
Gangs	59
10. Alcohol, Tobacco, and Other Drug Use	60
Current Substance Use	60
Lifetime Substance Use	65

Alcohol Use.....	70
Lifetime Alcohol Use	70
30-Day Alcohol Use	71
Binge Drinking.....	72
Average Age of First Alcohol Use	73
Levels of Problem Drinking: Composite Scale.....	74
Perception of Access to Alcohol	75
Usual Sources of Alcohol.....	76
Perception of Risk from Daily Alcohol Consumption.....	76
Tobacco Use	78
Lifetime Cigarette Smoking.....	78
30-Day Cigarette Smoking	79
Average Age of First Cigarette Smoking	80
30-Day Chewing Tobacco Use.....	81
30-Day Cigar, Cigarillo or Little Cigar Smoking.....	82
30-Day Electronic Cigarettes, E-cigs or Vape Pen Use	83
Secondhand Smoke Exposure.....	84
Perception of Access to Cigarettes	85
Perception of Risk from Heavy Cigarette Smoking (Pack or More Daily)	86
Marijuana Use	87
Lifetime Marijuana Use.....	87
30-Day Marijuana Use	88
Levels of Marijuana Use.....	89
Average Age of First Marijuana Use	90
Perception of Access to Marijuana	91
Usual Type of Marijuana	92
Perception of Risk from Regular Marijuana Use.....	93
Other Drugs Not Including Alcohol, Tobacco, or Marijuana	94
30-Day Other Drug Use (Not Including Alcohol, Tobacco, or Marijuana).....	94
Prescription Opiate (Painkiller) Use.....	95
Prescription Drug Misuse.....	96
Lifetime Methamphetamine Use.....	97
Lifetime Inhalant Use.....	98
Lifetime Heroin Use	99
Lifetime Cocaine Use	100
Lifetime Steroid Use.....	101
11. Risk and Protective Factors.....	102
Community Domain: Risk Factors.....	107
Community Domain: Protective Factors.....	108
School Domain: Risk Factors	109
School Domain: Protective Factors.....	110
Peer-Individual Domain: Risk Factors	111
Peer-Individual Domain: Protective Factors	112
Family Domain: Risk Factors	113
Family Domain: Protective Factors	114
References	115
Appendix	121

Tables

Table 1 State Sample School Response Rates in 2014.....	13
Table 2 Student Response Rates in 2014 (Valid Surveys)	13
Table 3: Respondent Characteristics in 2014, Percent of Students (and 95% CI).....	17
Table 4: Current (30-Day) Substance Use by Year, Grade 6	61
Table 5: Current (30-Day) Substance Use by Year, Grade 8	62
Table 6: Current (30-Day) Substance Use by Year, Grade 10	63
Table 7: Current (30-Day) Substance Use by Year, Grade 12	64
Table 8: Lifetime Substance Use by Year, Grade 6	66
Table 9: Lifetime Substance Use by Year, Grade 8	67
Table 10: Lifetime Substance Use by Year, Grade 10	68
Table 11: Lifetime Substance Use by Year, Grade 12	69
Table 12 Average Age of First Use and Regular Use of Alcohol in 2014	73
Table 13 Average Age of First Cigarette Use in 2014.....	80
Table 14: Average Age of First Marijuana Use in 2014.....	90
Table 15 Risk Factors Included in 2014.....	103
Table 16 Protective Factors Included in 2014.....	104
Table 17 Profile of Community Risk Factors,	107
Table 18 Profile of Community Protective Factors,	108
Table 19 Profile of School Risk Factors,	109
Table 20 Profile of School Protective Factors,	110
Table 21 Profile of Peer-Individual Risk Factors,.....	111
Table 22 Profile of Peer-Individual Protective Factors,	112
Table 23 Profile of Family Risk Factors,	113
Table 24 Profile of Family Protective Factors,	114

Acknowledgements

The planning and implementation of the 2014 administration of the Washington State Healthy Youth Survey were the products of an important collaborative effort among members of the Healthy Youth Survey Planning Committee, local educators, health professionals, and community members throughout the state of Washington. The members of the Healthy Youth Survey Planning Committee, and the authors of and contributors to this report, thank the students, school administrators, parents, and local prevention and health professionals who encouraged and supported school participation in the survey. The survey would be of little use or consequence if these individuals had not demonstrated their commitment to addressing the health behaviors and related risk and protective factors identified in the survey.

Members of the Healthy Youth Survey Planning Committee include: Cathy Wasserman, Riley Peters, Kevin Beck, Lillian Bensley, and Anar Shah at the Department of Health; Dixie Grunenfelder and Krissy Johnson at the Office of Superintendent of Public Instruction; Steve Smothers, Linda Becker, Martha Perla, Grace Hong, and Felix Rodriguez at the Department of Social and Health Services; and Mary Segawa at the Liquor and Cannabis Board.

Other contributors include Eric Ossiander, Jennifer Sabel, and Karen Guillies at the Department of Health; and Jennine Knight at the Department of Social and Health Services.

Special thanks are also due to Joe Kabel, Susan Richardson, Pete Lund, Greg Stogner, Bill Luchansky, and Curtis Mack of Looking Glass Analytics, Inc. for their contributions to the 2014 survey effort.

Executive Summary

Background

The Washington State Healthy Youth Survey (HYS) measures health risk behaviors that contribute to the health and safety of youth in Washington State. The survey results serve two important functions: first as needs assessment data for program planning; and second as a global look at the effectiveness of statewide prevention and health promotion initiatives based on a range of education and health-related goals at the federal and state levels.

The 2014 administration of the Healthy Youth Survey represents a collaborative effort among the Department of Health; the Office of Superintendent of Public Instruction; the Department of Social and Health Services' Division of Behavioral Health and Recovery; the Liquor and Cannabis Board; and the contractor, Looking Glass Analytics, Inc. Representatives of these agencies served as members of the Healthy Youth Survey Planning Committee, which guided every aspect of the survey development and implementation.

The HYS 2014 was the 14th administration of a statewide survey among Washington's students. This report provides results of HYS 2014, including comparisons by grade and by gender. It also includes past survey results, and looks at changes from the HYS 2012 results and trends from 2002-2014.

Participation

Washington State schools were randomly selected for the HYS 2014 statewide sample. Of the sampled schools asked to participate, about 88 percent of Grade 6 schools, 90 percent of Grade 8 schools, 87 percent of Grade 10 schools, and 82 percent of Grade 12 schools took part in the survey.

All Grade 6, 8, 10 and 12 students in the sampled schools were eligible to participate in the survey. An estimated 79 percent of the Grade 6 students, 79 percent of the Grade 8 students, 67 percent of the Grade 10 students, and 50 percent of the Grade 12 students in the original random sample of schools took part in the survey (estimates were based on fall 2014 enrollment data from the Office of Superintendent of Public Instruction). Non-response is both a function of schools choosing not to participate AND students not participating. Student non-participation could come from a number of reasons, including being absent on the day of the survey, they or their parents opted them out, or they did participate but their survey was not valid.

A total of 192 schools and 35,262 students contributed data to the statewide sample. In addition, 188,962 students in 890 schools participated in the survey as non-sampled schools. These additional schools received reports of their own results, but those results are not included in this statewide report because the schools were not part of the representative statewide sample.

Results

Some behaviors increase with age and others decrease as part of normal development. Most of the results in this summary are presented as a range, reporting from the lowest to the highest rate. Significant increases or decreases from 2012 to 2014 are also included. Results are provided for the following topics:

- Physical activity and dietary behavior
- Health status and health care
- Mental health
- Sexual behavior
- School climate
- Unintentional injury behaviors
- Intentional injury behaviors
- Alcohol, tobacco and other drug use

Physical Activity and Dietary Behavior

- Self-reported data on height and weight indicate that obesity was 9 percent of Grade 8 students and 11 percent of Grade 10 and 12 students. In addition, overweight was 14 percent of Grade 8 and 10 students and 13 percent of Grade 12 students.
- Sixty minutes of physical activity on at least seven days a week are recommended for youth. Twenty-nine percent of Grade 6 students, 31 percent of Grade 8 students, 24 percent of Grade 10 students and 21 percent of Grade 12 students met the physical activity recommendation. From 2012, there was a significant decrease in meeting the physical activity recommendation among Grade 6 and 8 students.
- Participation in physical education classes on five days a week was 41 percent of Grade 8 students, 26 percent of Grade 10 students, and 22 percent of Grade 12 students. In addition, among those that took physical education, spending more than 20 minutes exercising or playing sports in an average class was 90 percent of Grade 8 students, 91 percent of Grade 10 and Grade 12 students.
- Watching television or playing video games for two hours or less on an average school day was 53 percent of Grade 6 students, 42 percent of Grade 8 and Grade 10 students, and 45 percent of Grade 12 students.
- Eating fruit and vegetables five or more times a day was 24 percent of Grade 8 students, and 22 percent of Grade 10 and 12 students. From 2012, there was a significant decrease in eating fruit and vegetables five or more times a day among Grade 10 students.
- Eating dinner with their family most of the time or always was 75 percent of Grade 6 students, 68 percent of Grade 8 students, 61 percent of Grade 10 students, and 53 percent of Grade 12 students.
- Drinking two or more sweetened beverages was 7 percent of Grade 6 students, 9 percent of Grade 8 students, 11 percent of Grade 10 students, and 12 percent of Grade 12 students.
- Drinking sweetened beverages at school (including after school or weekend activities) was 40 percent of Grade 8 students, 49 percent of Grade 10 students, and 44 percent of Grade 12 students. From 2012, there were significant decreases in drinking sweetened beverages at school among Grade 10 and 12 students.
- Among those who drank soft drinks at school, purchasing soft drinks at school was 17 percent of Grade 8 and Grade 12 students, and 25 percent of Grade 10 students. From 2012, there were significant decreases in buying sodas at school (for those who drank soda at school) among Grade 8, 10 and 12 students.
- Students who reported that their family had to cut meal size or skip meals because of lack of money for food in the past year was 11 percent of Grade 8, 13 percent of Grade 10, and 14 percent of Grade 12 students. From 2012, there were significant decreases in having to cut meal size or skip meals among Grade 8, 10 and 12 students.

Health Status and Health Care

- Doctor-diagnosed or lifetime asthma was 14 percent of Grade 6 students, 18 percent of Grade 8 students, 22 percent of Grade 10 students, and 21 percent of Grade 12 students. From 2012, there were significant decreases in lifetime asthma among Grade 6 and 12 students.
- Current asthma was 7 percent of Grade 6 students, 8 percent of Grade 8 and Grade 12 students, and 10 percent of Grade 10 students. From 2012, there were significant decreases in current asthma among Grade 6, 8 and 12 students.

- Visiting a doctor in the past year for a checkup was 65 percent of Grade 8 students, 66 percent of Grade 10 students, and 61 percent of Grade 12 students. From 2010, there were significant increases in seeing a doctor in the past year among Grade 8, 10 and 12 students.
- Visiting a dentist in the past year for a checkup, exam, teeth cleaning, or other dental work was 77 percent of Grade 8 students, 79 percent of Grade 10 students, and 76 percent of Grade 12 students.
- Experiencing depressive feelings (i.e., had ever felt so sad or hopeless almost every day for two weeks in a row that they stopped doing some usual activities) during the past year was 27 percent of Grade 8 students, 35 percent of Grade 10 students, and 34 percent of Grade 12 students. From 2012, there were significant decreases in experiencing depressive feelings among Grade 10 and 12 students.
- Ever had sexual intercourse was 9 percent of Grade 8 students, 27 percent of Grade 10 students, and 52 percent in Grade 12 students. From 2012, there was a significant decrease in having sexual intercourse among Grade 8 students.
- Sexual orientation was added to HYS in 2014. About 1 percent of Grade 8 students, and 2 percent of Grade 10 and 12 students identified as gay or lesbian. About 5 percent of Grade 8 students, and 6 percent of Grade 10 and 12 students identified as bisexual. Also about 1 percent of Grade 8 students, 2 percent of Grade 10 students, and 4 percent of Grade 12 students reported having same-sex contact and about 2 percent of Grade 8 students, 6 percent of Grade 10 students, and 7 percent of Grade 12 students reported having contact with both females and males.

School Climate

- Feeling safe at school was 89 percent of Grade 6 students, 86 percent of Grade 8 students, 85 percent of Grade 10 students, and 87 percent of Grade 12 students.
- Being bullied at school in the past month was 31 percent of Grade 6 students, 28 percent of Grade 8 students, 23 percent of Grade 10 students, and 16 percent of Grade 12 students. From 2012, there were significant decreases in being bullied among Grade 8, 10 and 12 students.
- Being harassed with a computer or cell phone was 13 percent of Grade 8 students, 12 percent of Grade 10 students, and 11 percent of Grade 12 students. From 2012, there was a significant increase in being harassed with a computer or cell phone among Grade 8 students.
- Being harassed because of their perceived sexual orientation was 11 percent of Grade 8 students, 9 percent of Grade 10 students, and 7 percent of Grade 12 students.
- Carrying weapons at school in the past month was 3 percent of Grade 6 students, 4 percent of Grade 8 students, and 6 percent of Grade 10 students, and Grade 12 students.
- Being drunk or high at school in the past year was 6 percent of Grade 8 students, 14 percent of Grade 10 students, and 18 percent of Grade 12 students. From 2012, there was a significant decrease in being drunk or high at school among Grade 8 students.
- Drinking alcohol at school in the past month was 3 percent of Grade 8 students, 6 percent of Grade 10 and Grade 12 students. In addition, using marijuana at school in the past month was 3 percent of Grade 8 students, 7 percent of Grade 10 students, and 8 percent of Grade 12 students.
- Having someone at school for students to discuss substance-related problems (such as a counselor, intervention specialist, or some other school staff member) was 65 percent of Grade 8 students, and 56 percent of Grade 10 and Grade 12 students.

- Skipping school in the past month was 17 percent of Grade 6 students, 15 percent of Grade 8 students, 18 percent of Grade 10 students, and 28 percent of Grade 12 students. From 2012, there was a significant increase in skipping school among Grade 12 students.
- Enjoying school almost always was 31 percent of Grade 6 students, 18 percent of Grade 8 students, 11 percent of Grade 10 students, and 10 percent of Grade 12 students. From 2012, there were significant decreases in almost always enjoying school among Grade 8, 10 and 12 students.

Unintentional Injury Behaviors

- Riding in a vehicle in the past month that was driven by someone who had been drinking alcohol was 6 percent of Grade 6 students, 17 percent of Grade 8 students, 18 percent of Grade 10, and 17 percent of Grade 12 students. From 2012, there was a significant decrease in riding with a driver who had been drinking among Grade 10 students.
- Five percent of Grade 10 students and 9 percent of Grade 12 students drove a vehicle in the past month after they had been drinking alcohol. From 2012, there was a significant decrease in driving after drinking alcohol among Grade 12 students.
- Riding in a vehicle in the past month that was driven by someone who had been using marijuana was 10 percent of Grade 8 students, 19 percent of Grade 10 students, and 26 percent of Grade 12 students.
- Nine percent of Grade 10 students and 17 percent of Grade 12 students drove a vehicle in the past month after they had been using marijuana.
- Riding in a vehicle in the past month that was driven by someone who had been texting or emailing was 27 percent of Grade 6 students, 47 percent of Grade 8 students, 57 percent of Grade 10 students, and 59 percent of Grade 12 students.
- Eight percent of Grade 10 students and 43 percent of Grade 12 students drove a vehicle in the past month while texting or emailing.
- Taking formal swimming lessons was 56 percent of Grade 6 students, 59 percent of Grade 8 and 10 students, and 64 percent of Grade 12 students. In addition, 85 percent of Grade 6 students, 87 percent of Grade 8 students, and 86 percent of Grade 10 and 12 students reported they felt comfortable playing or swimming in water over their head.
- Among those who had been in a small boat such as a canoe, raft, or motorboat, always wearing a life vest when boating was 42 percent of Grade 8 students, 32 percent of Grade 10 students, and 27 percent of Grade 12 students. From 2012, there were significant decreases in life vest while boating among Grade 8, 10 and 12 students.

Intentional Injury Behaviors

- Attempting suicide in the past year was 9 percent of Grade 8 students, 10 percent of Grade 10 students, and 8 percent of Grade 12 students.
- Any physical fighting (not just fighting at school) in the past year was 23 percent of Grade 6 students, 27 percent of Grade 8 students, 22 percent of Grade 10 students, and 16 percent of Grade 12 students. From 2012, there were significant decreases in physical fighting among Grade 8, 10 and 12 students.
- Gang membership in the past year was 6 percent of Grade 8 and 10 students, and 5 percent of Grade 12 students.

Alcohol, Tobacco, and Other Drug Use

- Alcohol, marijuana and tobacco continue to be the substances most widely used by youth in Washington.
- 30-day alcohol use was 2 percent of Grade 6 students, 8 percent of Grade 8 students, 21 percent of Grade 10 students, and 33 percent of Grade 12 students. From 2012, there were significant decreases in 30-day alcohol use among Grade 8 and 10 students.
- Binge drinking (i.e., five or more drinks on at least one occasion during the previous two weeks) was 2 percent of Grade 6 students, 5 percent of Grade 8 students, 11 percent of Grade 10 students, and 19 percent of Grade 12 students. From 2012, there were significant decreases in binge drinking among Grade 8 and 10 students.
- 30-day cigarette smoking was 1 percent of Grade 6 students, 4 percent of Grade 8 students, 8 percent of Grade 10 students, and 13 percent of Grade 12 students. From 2012, there were significant decreases in 30-day cigarette smoking among Grade 8 and 12 students.
- 30-day e-cig or vape pen use was 8 percent of Grade 8 students, 18 percent of Grade 10 students, and 23 percent of Grade 12 students.
- 30-day chewing tobacco was 1 percent of Grade 6 and 8 students, 4 percent of Grade 10 students, and 5 percent of Grade 12 students. From 2012, there were significant decreases in 30-day chewing tobacco use among Grade 8 and 12 students.
- 30-day marijuana use was 1 percent of Grade 6 students, 7 percent of Grade 8 students, 18 percent of Grade 10 students, and 27 percent of Grade 12 students. From 2012, there was a significant decrease in 30-day marijuana use among Grade 8 students.
- 30-day use of other drugs (not including alcohol tobacco or marijuana) was 1 percent of Grade 6 students, 2 percent of Grade 8 students, 4 percent of Grade 10 students, and 7 percent of Grade 12 students. From 2012, there was a significant decrease in 30-day use of other drugs among Grade 8 students.
- 30-day use of prescription pain medication to “get high” was 2 percent of Grade 8 students, 5 percent of Grade 10 students, and 6 percent of Grade 12 students. From 2012, there were significant decreases in 30-day use of pain medication to get high among Grade 8, 10 and 12 students.

As in previous survey administrations, there was a clear relationship between the number of risk and protective factors present and the use of alcohol, cigarettes and marijuana for students in Grade 8 (the only grade examined in terms of risk and protective factors for this report). As the number of risk factors for individual students increased, the more likely they were to use alcohol, cigarettes, and marijuana. Similarly, as the number of protective factors for individual students increased, the less likely they were to use alcohol, cigarettes, and marijuana.

This report details the findings from the 2014 administration of the Healthy Youth Survey. HYS 2014 continues Washington State’s ongoing effort to assess the health of youth throughout the state. The results of the survey will be used by stakeholders at the state, county, district, school, and community levels who are interested in developing and improving prevention and intervention programs to better the lives of youth.

1. Introduction

The Washington State Healthy Youth Survey (HYS) is an effort to measure health risk behaviors that contribute to morbidity, mortality, and social problems among youth in Washington State. These behaviors include alcohol, tobacco, and other drug use; behaviors that result in unintentional and intentional injuries (e.g., violence); dietary behaviors and physical activity; and related risk and protective factors. The survey produces estimates of the prevalence of major adolescent health risk behaviors and provides crucial information to school officials, health professionals, human service agencies, policymakers, and parents as they work together to ensure the optimum health of young people across the state. This report uses the survey results to estimate the current status of these health risk behaviors and examine trends in the behaviors over the past 26 years.

The survey results also serve as important needs assessment data for program planning. They offer insight into the effectiveness of statewide prevention and health promotion initiatives designed to reach a range of education- and health-related goals at the federal and state levels. Federal initiatives of interest to readers of this report include these:

- No Child Left Behind (DOE, 2001), which addresses the importance of school safety.
- High School Graduation Initiative (US DOE, 2002).
- The National Drug Control Strategy (The White House, 2014).
- Substance Abuse Prevention and Mental Health Promotion Five Year Strategic Plan (SAMHSA, 2012).
- The U.S. Department of Health and Human Services' Healthy People 2020 Health Promotion Objectives (U.S. Department of Health and Human Services, 2010).

State initiatives of interest to readers of this report include these:

- The Washington State Board of Health Strategic Plan 2009 (Washington State Board of Health, 2009).
- Graduation: A Team Effort (GATE) Initiative (OSPI, 2011).

The 2014 administration of the Healthy Youth Survey (HYS 2014) meets a wide variety of information needs by producing:

- Empirical needs assessment data necessary for planning substance abuse and other prevention and early intervention programs, including county-level six-year strategic plans.
- Data for studying trends of student substance use and abuse, and associated risk and protective factors.
- Information to support monitoring of the state's block grant for substance abuse prevention and treatment from the Substance Abuse and Mental Health Services Administration.
- Needs assessment, evaluation, and monitoring of federal grants to prevent and reduce substance use such as the Reducing Underage Drinking Initiative and the evaluation of results from the Partnership for Success Grant.
- Information to support the evaluation of prevention and education programs funded under the federal Safe and Drug-Free Schools and Communities Act, the federal Tobacco Settlement, and the state Omnibus Controlled Substance and Alcohol Abuse Act.
- Data to measure the progress toward attainment of the state's goals for substance abuse prevention.
- Information on the progress of programs implemented pursuant to the state's Youth Violence Act (E2SHB 2319).

- Information on sexual education in schools used to help monitor implementation of the Healthy Youth Act.
- Needs assessment data used as part of the Comprehensive Needs Assessment for the Maternal and Child Health Block Grant.
- Data that can contribute information to local community profiles designed to help community stakeholders understand the importance of programs that support youth.
- Data to describe risk and protective factors that can be used by local school and community members as they plan or refine school- and community-based prevention and intervention programs.
- Data to support community and state level grant applications.
- Data to support the Governor’s Results Washington Initiative (<http://www.results.wa.gov/>)

HYS 2014 represents a collaborative effort by the Department of Health; the Office of Superintendent of Public Instruction; the Department of Social and Health Services’ Division of Behavioral Health and Recovery; the Liquor and Cannabis Board; and the survey contractor, Looking Glass Analytics, Inc. Representatives of these agencies served as members of the Healthy Youth Survey Planning Committee, which guided every aspect of the survey development and implementation. In addition, staff members from the University of Washington’s Social Development Research Group provided consultation on the risk and protective factors assessment portion of the survey.

Staff members at the nine Educational Service Districts (ESDs) coordinated local school recruitment efforts and provided technical assistance. Local health jurisdictions, educational agencies, and other local partners provided valuable input into the development and administration of the survey.

HYS 2014 was the 14th administration of a statewide survey among Washington’s students. Ten of the surveys included students in Grades 6, 8, 10, and 12, one survey (1988) included students in Grades 6, 8 and 10, and one survey (1999) included students in Grades 9 through 12. The survey content and methodology have varied over time:

- The first two administrations in 1988 and 1990 included only questions about alcohol, tobacco, and other drug use and associated behaviors (Deck and Nickel, 1989; Gabriel, 1991).
- The 1992 and 1995 surveys asked additional questions that addressed other health risk behaviors (Einspruch and Pollard, 1993; Gabriel, Deck, Einspruch, and Nickel, 1995).
- The 1998 survey focused on alcohol, tobacco, and other drug use and related risk and protective factors (Einspruch, Gabriel, Deck, and Nickel, 1998).
- The 1999 survey (Bensley, VanEenwyk, Schoder, and Tollefsen, 2000) was based on the Centers for Disease Control and Prevention’s Youth Risk Behavior Survey (Grunbaum et al., 2004).
- The 2000 survey was similar to the 1988 survey and focused on alcohol, tobacco, and other drug use and related risk and protective factors (Einspruch, Deck, Nickel, and Hyatt, 2001).
- Surveys since 2002 included items related to health behaviors, substance use, and related risk and protective factors (Einspruch and Hyatt, 2004), (Einspruch, 2005, and 2007)

Copies of prior Analytic Reports are available online at: <http://www.askhys.net/Reports/Additional>.

Organization and Purpose of the Report

This report provides the results of the 2014 administration of the Healthy Youth Survey and results from the earlier Washington State surveys. It is organized in the following sections.

- Chapter 1 describes the purpose of this report.
- Chapter 2 describes the survey methods.
- Chapter 3 presents results related to physical activity and dietary behaviors.
- Chapter 4 presents results related to health status and health care.
- Chapter 5 presents results related to mental health
- Chapter 6 presents results related to sexual behavior
- Chapter 7 presents results related to school climate.
- Chapter 8 presents results related to unintentional injury behaviors.
- Chapter 9 presents results related to intentional injury behaviors.
- Chapter 10 details results related to alcohol, tobacco, and other drug use.
- Chapter 11 details results pertaining to relevant risk and protective factors.
- And the Appendix includes all of the Healthy Youth Survey 2014 state sample results by grade.

Chapters 3 through 11 are organized so that the 2014 results are presented first, followed by comparative analyses to test for differences by grade level and gender. Next, the differences in Washington State survey results over time are presented along with the results of comparative analyses to test for significant differences from 2012 to 2014 and trend analyses for items that have five or more years of data. These comparisons allow readers to view the trends over past years' reports of health risk behaviors among Washington's students at the same grade levels.

Throughout the report, national- and state-level goals, objectives, and benchmarks—such as Healthy People 2020 (U.S. Department of Health and Human Services, 2010)—are included to provide a context in which to review the results.

Participation

The Department of Health selected three simple random samples of schools serving Grade 6, Grade 8, and Grades 10 and 12 to constitute representative samples of Washington's Grade 6, 8, 10, and 12 students. One sample was drawn for Grades 10 and 12 because those grades usually occur together in a high school, whereas Grades 6 and 8 may be together in a middle school, or separate in an elementary school or junior high school. Of those schools asked to participate in the survey, about 87 percent with Grade 6 students, 94 percent with Grade 8 students, 87 percent with Grade 10 students, and 82 percent with Grade 12 students took part in the survey.

Overall response rates are about 79 percent of the Grade 6 students, 79 percent of the Grade 8 students, 67 percent of the Grade 10 students, and 50 percent of the Grade 12 students. These participation rates are based on the October 2014 enrollment in all sampled schools (including non-participating schools). Non-response is both a function of schools choosing not to participate AND students not participating. Student non-participation could come from a number of reasons, including being absent on the day of the survey, they or their parents opted them out, or they did participate but their survey was not valid. Although the Grade 10 and 12 participation rates are below 70 percent, these findings are expected to be representative of Washington youth in public schools, based on an extensive examination of bias conducted for HYS 2002, 2004, 2008, 2010, and 2014.

Looking Glass Analytics' analysis of the survey results included a series of quality controls to remove data that were incomplete, obviously inaccurate, or internally inconsistent (e.g., reporting no lifetime use of a substance and also reporting use of the same substance in the past 30 days). The results presented in this report are not perfect estimates. There are margins of error indicated by the confidence intervals.

A total of 35,292 students in 192 schools contributed data to the statewide results. In addition, 187,910 students in 887 schools participated in the survey as non-sampled schools. Non-sampled schools received reports of their own results, but those results are not included in this statewide report because the schools were not part of the representative statewide sample.

Over the life of the survey, the number of participating students has grown: 10,485 in 1988, 18,375 in 1990, 15,463 in 1992, 20,780 in 1995, 52,332 in 1998, 102,532 in 2000, 137,515 in 2002, 185,095 in 2004, 198,312 in 2006, 210,851 in 2008, 211,331 in 2010, 204,929 in 2012, and 224,224 in 2014. Participation may reflect increasing interest across the state in health-related information and is a tribute to the collaboration and funding effort among sponsoring agencies, schools, and local community members.

Cautions

Readers should bear in mind several cautions when interpreting the survey results presented in this report. This section describes these cautions in detail.

Representativeness

Survey responses are often used to estimate the frequency of behaviors or other characteristics in a population larger than that which actually completed the survey. Thus, the results of the survey are used to characterize all Grade 6, 8, 10, and 12 students in Washington even though only a portion of public school students took the survey. This is possible only if the students who participated in the survey are not different from those who did not participate. If they are different, the survey is considered biased and the results are limited in their ability to be generalized to all students. Bias represents systematic error and is different from the random fluctuation measured by confidence intervals.

The 2014 HYS results are generalizable to the majority of youth in Washington State, but may underrepresent students attending small and non-urban public schools. They also may not be representative of youth who attend private schools, nonpublic tribal schools, home school, or who have dropped out of school. Students in juvenile detention facilities are restricted from participating in the survey.

In previous survey administrations, alternative schools were less likely to participate in HYS. In 2014, alternative schools were just as likely as traditional schools to participate. Very few alternative schools were selected for the 2014 state sample, possibly making it difficult to detect a difference in participation.

Trends

In comparing the results of HYS 2014 survey and earlier surveys, readers should remember that certain factors may influence apparent trends. For example, information about the characteristics of the 1988 and 1990 samples is not readily available. Comparisons with the 1992 survey might be influenced by the inclusion of non-sampled schools in the data from that year, although comparisons between the sampled and non-sampled schools that year revealed similar levels of substance use. In addition, the wording of some of the survey items has changed over the years so that some items are only somewhat

comparable over the years, and some are not comparable at all. A description of changes to substance use survey items over time is available on pages 60 and 65.

Many administration procedures and data processing concepts have, however, been consistent over time, and the Healthy Youth Survey 2002, 2004, 2006, 2008, 2010, 2012, and 2014 administrations were very similar.

Results for every year available are presented in charts and tables throughout the report. Trend analyses only include results from 2002 through 2014. Trends were assessed using Joinpoint software, available at <http://surveillance.cancer.gov/joinpoint/>. Models with at least 7 time points allowed for one joinpoint (change in trend line); however, if the analysis revealed a significant joinpoint, but neither trend achieved significance, the analysis was rerun allowing no joinpoints. This affected 4 trend lines (current chewing tobacco and alcohol use for grade 8 and current cigarette use for grades 10 and 12).

Rounding Differences

Results presented in this Analytic Report were calculated to two decimal points and then rounded to whole numbers. Results presented in the Appendix of this report and in the local reports prepared by Looking Glass Analytics were also calculated to two decimal points and then rounded to one decimal point. If the results ending in 0.5 in the Appendix or local reports were rounded to whole numbers, those rounded results may be 1 percent different from the whole numbers presented in this report. For example, if a result in the Appendix is 8.5 percent, then you would round up to 9 percent. But that 8.5 percent could have originally been 8.49 percent - thus it was rounded down to 8 percent in this report.

School Dropouts

In interpreting differences between survey results for each grade level, readers should remember that some reported behaviors and risk factors may appear more prevalent in Grade 10 compared to Grade 12 because of increased rate of school dropout after age 16 (i.e., prior to Grade 12). It is generally accepted that the results for high school seniors in surveys such as this one are underestimates because many of the youth most likely to engage in risky behaviors may have dropped out of school (Johnston, O'Malley, and Bachman, 1994). Thus the authors recommend interpreting results for high school seniors with some caution, particularly when their prevalence rates differ markedly from those of students in earlier grades.

The school dropout concern is not new and has existed in previous Washington surveys. Unless the characteristics of school dropouts have changed over time, the bias in Grade 12 estimates is likely similar to what it has been in the past. This fact means that although any given year's data on health risk behaviors among Grade 12 students may be an underestimate, the year-to-year comparisons are likely to be less affected by this bias (Johnston et al., 1994).

Developmental Changes

In interpreting differences between grade levels, readers should remember that developmental changes may influence students' perceptions and accuracy of reporting. These factors include the ability to read or accurately interpret the intention of survey questions, to accurately recall events during a specific time frame, or to have developed opinions about different topics.

Self-Report Data

The survey measures self-reports, which may be influenced by factors including problems in remembering, social desirability or the wish to present oneself in a positive manner, reading ability, and developmental changes.

Correlational Data

Interrelationships among the variables should not be interpreted as indicating that one variable caused the other. Although this causal relationship might exist, the direction of the correlation may be reverse of what is expected, or an apparent relationship might be due to some other measured or unmeasured cause.

2. Methods

This chapter details the methodological considerations of HYS 2014. The chapter addresses the topics of sampling, survey administration, the questionnaires, reliability and validity, data preparation and analysis, response rates, non-completion rates, and the characteristics of the students who completed the survey. The survey procedures were approved by the Washington State Institutional Review Board.

Sampling

The statewide results presented in this report are based on a statewide sample of all schools in the public school system serving the surveyed grades, with at least 15 students in each grade. For the statewide sample, Department of Health epidemiology staff members drew three simple random samples of all public schools serving Grade 6, Grade 8, and Grades 10 and 12. This procedure was used because Grades 10 and 12 usually occur together within a single school, whereas Grades 6 and 8 may be together in a middle school or separate in an elementary school and a junior high school. About 28 percent of the schools had fewer than 15 students per grade, but these schools accounted for only 1 percent of the students. Consequently, excluding these schools saves considerable effort in the recruitment and administration phase without biasing the final results.

To obtain a confidence interval of plus or minus 3 percent for statewide results at each grade, based on the intraclass correlations obtained in the 2000 survey, it was estimated that a sample size of about 5,335 students would be needed per grade. The average school enrollments were 131 in Grade 6, 180 in Grade 8, 196 in Grade 10, and 200 in Grade 12. Using estimations of a 50 percent response rate for schools and a 90 percent response rate for students within the participating schools and experience from the 2002, 2004, 2006, 2008, 2010, and 2012 surveys, the sample was drawn to include 99 schools serving Grade 6, 67 schools serving Grade 8, 65 schools serving Grades 10 and 12, and 3 schools serving Grade 12 but not 10. The additional schools for Grade 12 were necessary because they had lower average enrollments than the Grade 10 schools.

Schools not selected for the state sample were offered an opportunity to participate in the survey by “piggybacking” onto the statewide data collection effort. The Department of Health also drew county samples in six large counties where the reduction in the number of schools in a sample compared to a census justified the additional effort associated with drawing and analyzing a sample (Clark, King, Pierce, Snohomish, Spokane and Thurston for Grade 6; King, Pierce, Snohomish, and Spokane for Grade 8; and King, Pierce, and Snohomish for Grades 10 and 12). For county samples, additional schools were added to those already in the state sample. The data from the piggyback schools, including those drawn for the county samples, are not included in the results presented in this report because they were not part of the state sample.

Survey Administration

All Washington public schools, except institutional/correctional schools, serving Grades 6, 8, 10, or 12 were invited to participate in the survey as either a state sampled, county sampled, or piggyback school at the beginning of the 2014 calendar year. Schools that wished to participate registered between February and the end of June 2014.

Each school designated a survey coordinator. The survey contractor and sponsoring agencies offered an on-line training to provide the coordinators with the information necessary to successfully administer

the survey. Materials were made available on the project website, www.AskHYS.net. Coordinators were instructed to train the teachers in their school(s) who were to administer the survey to students (teacher training materials were provided to the coordinators).

The coordinators received detailed written instructions with their survey materials, along with materials used to notify parents and students prior to the survey administration. Parents had an opportunity to decline their child's participation, and students could also choose not to participate. The coordinators distributed the survey materials to the teachers, who in turn distributed them to the students and proctored the survey administration. Students participated on a voluntary and anonymous basis. Students who did not wish to participate were provided with an alternative activity.

Teachers read a standardized set of instructions to the students, informing them of the importance of the survey. The survey was to be administered to all participating students in a single class period during the school day. Students absent that day were not to make up the survey. Students placed their completed answer sheets in an envelope that was sealed, returned to the coordinator, and ultimately returned to Looking Glass Analytics.

Questionnaires

The questions on HYS 2014 were derived primarily from the following sources: the Monitoring the Future survey (Johnston et al., 1994; National Institute on Drug Abuse, 2001), the Youth Risk Behavior Survey (Eaton et al., 2006), the Global Youth Tobacco Survey (Centers for Disease Control and Prevention, 2000), and the Communities that Care Survey (Arthur, Hawkins, Catalano, and Pollard, 1998). In 2014, there were three main survey forms - Form C for primary students and Forms A and B for secondary students. The questions for secondary students were divided into two forms (A and B) because the number of items of interest to the sponsoring agencies was greater than could be answered by a student during the allotted time (one class period).

Form A mainly contained items from the Monitoring the Future survey and the Communities that Care Survey. Form B mainly contained items from the Youth Risk Behavior Survey and the Global Youth Tobacco Survey.

Secondary schools that wanted to ask optional questions could register for an enhanced version of Survey Forms A and B. Form A-enhanced included one question on sexual orientation and Form B-enhanced included four questions on sexual behavior and two on sexual abuse.

Form A had 138 items, Form A-enhanced had 139 items, Form B had 116 items, Form B-enhanced had 122 items. Thirty-five items were common to both forms A/A-enhanced and B/B-enhanced. Students in Grades 8, 10, and 12 completed Forms A/A-enhanced and B/B-enhanced (the forms were alternated when they were packaged by the printer so that in a classroom every other student completed Form A/A-enhanced and every other student completed either Form B/B-enhanced, effectively distributing the two forms randomly among the students). The number of survey questions on each survey form does not match the number of survey items on each form, as some survey items include multiple survey questions.

Form C contained 76 items drawn primarily from Forms A and B (not including the optional questions) and was completed by students in Grade 6.

Translations

The survey was available in English and Spanish. All schools received Spanish-language survey materials. The survey coordinators duplicated the Spanish survey materials locally and provided them to the students as needed. Students read the translated survey but responded on the English answer sheet to preserve anonymity. It is, therefore, impossible to know how many students read a Spanish survey.

Reliability and Validity

A survey item is *valid* if it accurately measures the concept it is intended to measure. A survey item is *reliable* if it consistently produces the same results under the same circumstances. Nearly all HYS 2014 questions were gleaned from four established surveys that have been used throughout the United States—some for more than 25 years. Each of these surveys has been subjected to scientific research regarding reliability and validity, and has been field-tested extensively (Arthur et. al, 1998; Eaton et al., 2006; Johnston et al., 1994). This field testing generally addresses such issues as the content and structure of questions, the ordering of questions, the types and ordering of the response options, and survey length.

Bensley (1997) reviewed the reliability and validity of school-based surveys and found adequate reliability based on a large test–retest study and on studies of interrelationships among the data (such as gender and age differences and differences between dropouts and in-school youth). Bensley found that remaining questions about validity were based on differences among methodologies. School-based, self-administered surveys appeared to yield higher prevalence than either telephone surveys or face-to-face interviews, but lower prevalence than biochemical indicators of substance use or methods that provide even greater anonymity. Biochemical indicators, which provide the most objective comparison data, and low self-reported use of a fictitious drug suggest that most self-reported behaviors on school-based surveys are likely valid but some underreporting may occur. Underreporting of socially disapproved behaviors has been noted for both adults and youth, particularly when the possibility is greater that the responding individual is identifiable.

Data Preparation and Analysis

Looking Glass Analytics, Inc. received, prepared, and scanned the completed answer sheets, then cleaned the data using programs designed to detect dishonest and inconsistent answers. Most data processing and analytic code were written using SAS analytic software.

Looking Glass Analytics, Inc. also used SAS to create local reports with item-level frequency distributions and scale results for the participating schools (unless the school requested at the time of registration that these reports not be sent), districts, counties, and ESDs. In all cases a minimum of 15 valid, completed surveys were required at a given grade level for a grade level report to be produced. In addition, 70 percent or more of the students enrolled at a district, county, or ESD were required to have participated in the survey for a report of results to be produced at that level (if participation was between 40 and 69 percent, a “report of participating schools” was produced). An interpretive guide to aid recipients in reading their report was made available on the project web site, www.AskHYS.net. Statewide results were presented as comparative data in the local reports.

For this Analytic Report, STATA Statistical Software was used for the following additional analyses.

Differences by Grade Level and Gender

A chi-square test of significance was used to compare 2014 results among grade levels and between genders. Comparisons with a p -value less than 0.05 were considered significant differences.

Differences over Time

A chi-square test of significance was used to compare HYS 2012 results to HYS 2014 results. Comparisons with a p -value less than 0.05 were reported as significant differences.

Joinpoint analysis (National Cancer Institute, 2005) was used to examine trends over time back to 2002 for those questions that had been asked on five or more administrations of the survey. Differences in the linear trend of the time span of the question are reported for analyses in which the p -value was less than 0.05. Joinpoint analysis tested both whether there was a significant trend over time and whether there was a change in the trend over time (i.e., a change in inflection). The Joinpoint analysis allowed one change in trend if there were seven time points. The direction of the differences and if there was a significant change in trend, the time spans with significant trends are reported for analyses in which the p -value was less than 0.05.

Washington data presented in this report are from surveys that were implemented in Washington public schools from 1988 to 2014. Results from the 1998 to 2000 surveys are presented in charts when available, but not included in trend analyses.

- **1988: Student Alcohol and Drug Use Survey (SADUS)**—This health risk-focused survey was administered in public schools in the fall of 1988. A total of 10,485 Grade 6, 8, and 10 students in 125 schools participated in the state sample for a state response rate of about 50 percent.
- **1990: Student Alcohol and Drug Use Survey**—SADUS was administered in public schools in the fall of 1990. A total of 18,375 Grade 6, 8, 10, and 12 students in 176 schools participated in the state sample for a state response rate of about 65 percent.
- **1992: Washington State Survey of Adolescent Health Behaviors (WSSAHB)**—This substance use and risk and protective factor-focused survey was administered in public schools in the fall of 1992. Because the state sample response rate was 45 percent, sampled and non-sampled schools were combined for the report (a total of 15,463 Grade 6, 8, 10, and 12 students in 144 schools).
- **1995: Washington State Survey of Adolescent Health Behaviors**—WSSAHB was administered in public schools in the spring of 1995. A total of 8,780 Grade 6, 8, 10, and 12 students in 89 schools participated in the state sample for a state response rate of about 25 percent. An additional 12,060 students participated in the survey voluntarily and contributed to local results.
- **1998: Washington State Survey of Adolescent Health Behaviors**—WSSAHB was administered in public schools in the spring of 1998. A total of 14,601 Grade 6, 8, and 10 students in 102 schools participated in the state sample for a state response rate of about 60 percent. An additional 37,731 students participated in the survey voluntarily and contributed to local results.
- **1999: Washington State Youth Risk Behavior Survey**—This health risk-focused survey was administered in public schools in the spring of 1999. A total of 7,642 Grade 9, 10, 11, and 12 students completed the survey (4,022 from the Seattle region and 3,602 across the state). The overall response rate was about 40 percent.
- **2000: Washington State Survey of Adolescent Health Behaviors**—WSSAHB was administered in public schools in the fall of 2000. A total of 17,780 Grade 6, 8, 10, and 12 students in 98 schools participated in the state sample for a state response rate of about 65 percent. An additional 84,662 students participated in the survey voluntarily and contributed to local results.
- **2002: Healthy Youth Survey**—This health risk and risk and protective factor-focused survey was administered in public schools in the fall of 2002. A total of 24,685 Grade 6, 8, 10, and 12

students in 171 schools participated in the state sample for a state response rate of about 55 percent. An additional 112,650 students participated in the survey voluntarily and contributed to local results.

- **2004: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2004. A total of 30,263 Grade 6, 8, 10, and 12 students in 191 schools participated in the state sample for a state response rate of about 65 percent. An additional 154,832 students participated in the survey voluntarily and contributed to local results.
- **2006: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2006. A total of 32,531 Grade 6, 8, 10, and 12 students in 203 schools participated in the state sample for a state response rate of about 65 percent. An additional 165,781 students participated in the survey voluntarily and contributed to local results.
- **2008: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2008. A total of 30,346 Grade 6, 8, 10, and 12 students in 201 schools participated in the state sample for a state response rate of about 66 percent. An additional 180,505 students participated in the survey voluntarily and contributed to local results.
- **2010: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2010. A total of 34,069 Grade 6, 8, 10, and 12 students in 212 schools participated in the state sample for a state response rate of about 70 percent. An additional 177,262 students participated in the survey voluntarily and contributed to local results.
- **2012: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2012. A total of 33,207 Grade 6, 8, 10, and 12 students in 201 schools participated in the state sample for a state response rate of about 69 percent. An additional 171,659 students participated in the survey voluntarily and contributed to local results.
- **2014: Healthy Youth Survey**—HYS was administered in public schools in the fall of 2014. A total of 35,262 Grade 6, 8, 10, and 12 students in 192 schools participated in the state sample for a state response rate of about 68 percent. An additional 188,962 students participated in the survey voluntarily and contributed to local results.

Chart Formatting

In Analytic Reports before 2012, bar charts detailing specific outcomes by year and grade were presented. As additional years have been added, these bar charts have become increasingly complex, and so since the 2012 Analytic Report, we used line charts for most outcomes. In order to maintain readability we have not presented confidence intervals in the charts. However, confidence intervals for 2014 are available at the end of this report in the Appendix: Healthy Youth Survey 2014 statewide sample results and confidence intervals for previous years are depicted in previous Analytic Reports (available online at www.AskHYS.net). Also, significance tests for changes between 2012 and 2014, and tests of trends between 2002 and 2014 are included in this Analytic Report for each of the outcomes depicted in the charts.

Calculating Confidence Intervals

Reports of results from previous Washington State surveys are available on www.AskHYS.net. Confidence intervals for the 1999, 2002, 2004, 2006, 2008, and 2010 data were obtained by direct analysis using SUDAAN. For 2012 and 2014, confidence intervals were obtained using SAS. Confidence intervals for the 1992, 1995, 1998, and 2000 data were based on estimates provided in the respective reports (and confidence intervals for 1988 and 1990 were based on the 1992 estimates). Confidence

intervals in these years were limited to single estimates that have been applied to all percentages obtained in those years:

- For 1988, 1990, and 1992 percentages near 50 percent, these estimates were plus or minus 1.4 percent for Grade 6, 1.4 percent for Grade 8, 1.7 percent for Grade 10, and 2.0 percent for Grade 12. For 1988, 1990 and 1992 percentages near 10 or 90 percent, these estimates were plus or minus 0.9 percent for Grade 6, 0.8 percent for Grade 8, 1.0 percent for Grade 10, and 1.2 percent for Grade 12. Twenty-five percent was used to divide these two groups of percentages. (The confidence intervals for 1998 and 1990 are based on the estimates provided in 1992.)
- For 1995 these estimates were plus or minus 2 percent for Grade 6, 2 percent for Grade 8, 2 percent for Grade 10, and 4 percent for Grade 12.
- For 1998 these estimates were plus or minus 2 percent for Grade 6, 3 percent for Grade 8, 4 percent for Grade 10, and 4 percent for Grade 12.
- For 2000 these estimates were plus or minus 3 percent for Grade 6, 3 percent for Grade 8, 4 percent for Grade 10, and 4 percent for Grade 12.

Response Rates

The overall response rates (the number of participating students who completed valid surveys divided by the total enrollment in schools asked to participate in the state sample) were 79 percent in Grade 6, 79 percent in Grade 8, 67 percent in Grade 10, and 50 percent in Grade 12. Participation rates presented here are based on the 2014 enrollment data from the Office of Superintendent of Public Instruction’s P-105 October Enrollment Headcount Report for October 2014 (retrieved from <http://www.k12.wa.us/DataAdmin/default.aspx>). Although some of the participation rates are below 70 percent, these findings are expected to be representative of most Washington youth in public schools based on an examination of bias conducted for HYS 2002, 2004, 2008, 2010, and 2014.

Table 1 provides the response rates for schools calculated by dividing the number of participating schools by the number of schools asked to participate. Because some schools were selected for more than one sampled grade, the total number of schools is less than the sum of the number of schools at each grade.

Table 1
State Sample School Response Rates in 2014

Number of Schools			
Grade	School Participated	Schools Asked to Participate	Response Rate
Grade 6	87	99	88%
Grade 8	60	67	90%
Grade 10	54	62	87%
Grade 12	53	65	82%

Table 2 provides the percentage of valid surveys compared to total enrollment in sampled schools asked to participate.

Table 2
Student Response Rates in 2014 (Valid Surveys)

Grade	Number of Valid Surveys	Enrollment in Schools Asked to Participate	Percent of Valid Surveys
Grade 6	9,129	11,511	79%
Grade 8	10,680	13,556	79%
Grade 10	8,833	13,269	67%
Grade 12	6,650	13,239	50%
Total	35,292	51,575	68%

Of the original 239,360 surveys that were submitted from all schools (sampled and “piggyback”), a total of 7,457 were removed during the scanning process because they were unscannable or from an invalid grade level or grade for the school. The remaining 231,903 scanned surveys were screened to detect dishonest and inconsistent answers. A total of 7,679 were dropped during the data cleaning process. This was about 2 percent of Grade 6 surveys, 3 percent of Grade 8 surveys, and 4 percent of Grade 10 and 5 percent of Grade 12 surveys. Another 1,830 surveys were completed by students who used the wrong survey form for their grade. Responses from students who took the wrong form were included in school building results, but excluded from higher aggregations, such as district, county and state results.

Non-completion Rates by Form

HYS 2014 consisted of five forms, one elementary version and four secondary versions. Two of the four secondary versions included optional questions; Form A-enhanced and Form B-enhanced. Figures 1 and 2 illustrate the percentage of Grade 8, 10 and 12 students who did not complete each item on Form A and Form A-enhanced; Figures 3 and 4 illustrate the percentage of Grade 8, 10, and 12 students who did not complete each item on Forms B and Form B-enhanced; and Figure 5 illustrates the percentage of Grade 6 students who did not complete each item on Form C.

Schools could choose to ask the optional questions on Form A-enhanced or Form A-enhanced and Form B-enhanced when they registered for the 2014 HYS.

The rates at which respondents failed to complete the last question on a survey by form type were:

- 22 percent of Grade 8, 14 percent of Grade 10, and 12 percent of Grade 12 students did not complete Form A.
- 18 percent of Grade 8, 17 percent of Grade 10, and 15 percent of Grade 12 students did not complete Form A-enhanced.
- 17 percent of Grade 8, 12 percent of Grade 10, and 9 percent of Grade 12 students did not complete Form B.
- 17 percent of Grade 8, 14 percent of Grade 10, and 12 percent of Grade 12 students did not complete Form B-enhanced.
- 2 percent of Grade 6 students did not complete Form C.

Conversely, 85 percent of students completed at least:

- 109 questions for Grade 8, 132 for Grade 10, and 138 for Grade 12 out of 138 questions on Form A.
- 108 questions for Grade 8, 116 for Grade 10, and 116 for Grade 12 out of 116 questions on Form B.
- 93 out of 93 questions for Form C.

Figure 1: Non-completion Rates for Form A, Grades 8, 10, and 12 in 2014




Figure 2: Non-completion Rates for Form A-enhanced, Grades 8, 10, and 12 in 2014




Figure 3: Non-completion Rates for Form B Grades 8, 10, and 12 in 2014






Figure 4: Non-completion Rates for Form B-enhanced Grades 8, 10, and 12 in 2014



On Form B and B-enhanced, there was a spike at questions 37 and 38 – these questions were about self-report height and weight.

Figure 5: Non-completion Rates for Form C, Grade 6 in 2014



Respondent Characteristics

The findings of HYS 2014 presented in this report are based on the responses of 35,262, students in Grades 6, 8, 10, and 12. These students were selected using a scientific sampling plan intended to represent the full population of public school students at these grade levels across the state. Table 3 provides details about the demographic characteristics of the participating students.

Table 3: Respondent Characteristics in 2014, Percent of Students (and 95% CI)

	6th Grade % (±CI)	8th Grade % (±CI)	10th Grade % (±CI)	12th Grade % (±CI)
Age	(n=9,126)	(n=10677)	(n=8829)	(n=6649)
10 or younger	1.5% (±0.3)	**	**	**
11	76.0% (±1.2)	**	**	**
12	21.8% (±1.2)	1.1% (±0.2)	0.1% (±0.1)	0.1% (±0.1)
13	0.6% (±0.2)	74.8% (±1.4)	0.1% (±0.1)	0.0% (±0.0)
14	0.0% (±0.0)	23.4% (±1.4)	1.4% (±0.3)	0.0% (±0.0)
15	0.1% (±0.1)	0.7% (±0.2)	73.7% (±1.6)	0.0% (±0.0)
16	**	0.0% (±0.0)	24.0% (±1.6)	1.4% (±0.3)
17	**	0.0% (±0.0)	0.7% (±0.2)	72.7% (±1.7)
18	**	0.0% (±0.0)	0.1% (±0.0)	24.1% (±1.5)
19 or older	**	0.0% (±0.0)	0.0% (±0.0)	1.6% (±0.6)
Gender	(n=9118)	(n=10660)	(n=8817)	(n=6635)
Female	50.6% (±1.1)	50.3% (±1.0)	51.6% (±1.0)	49.8% (±1.4)
Male	49.4% (±1.1)	49.7% (±1.0)	48.4% (±1.0)	50.2% (±1.4)
Race - Ethnic Group	(n=8696)	(n=10550)	(n=8794)	(n=6640)
American Indian or Alaskan Native	5.9% (±1.1)	3.3% (±0.7)	2.4% (±0.6)	2.1% (±0.8)
Asian or Asian American	9.7% (±3.4)	9.9% (±3.1)	9.3% (±2.8)	8.6% (±2.8)
Black or African-American	4.8% (±1.4)	4.1% (±1.0)	4.9% (±1.5)	4.9% (±1.6)
Hispanic or Latino/Latina	12.3% (±3.5)	15.1% (±5.1)	11.8% (±4.7)	12.4% (±5.0)
Native Hawaiian or other Pacific Islander	1.6% (±0.4)	1.7% (±0.4)	2.2% (±0.7)	1.9% (±0.8)
White or Caucasian	39.1% (±4.7)	49.3% (±5.5)	56.0% (±6.2)	58.8% (±6.5)
Other	17.4% (±1.8)	8.0% (±0.7)	5.6% (±0.6)	3.9% (±0.5)
More than one race/ethnicity marked	9.3% (±0.8)	8.7% (±1.1)	7.9% (±1.0)	7.4% (±0.9)
Language Spoken at Home	(n=9089)	(n=10431)	(n=8608)	(n=6501)
English	80.7% (±4.4)	79.5% (±4.0)	82.0% (±4.1)	82.8% (±4.1)
Spanish	10.0% (±2.9)	10.2% (±3.5)	7.4% (±3.2)	7.5% (±3.5)
Russian	**	1.0% (±0.3)	1.5% (±0.5)	1.2% (±0.5)
Ukrainian	**	0.8% (±0.3)	0.9% (±0.3)	1.0% (±0.3)
Vietnamese	**	1.4% (±0.9)	1.5% (±0.7)	1.4% (±0.7)
Chinese	**	1.6% (±1.0)	1.1% (±1.0)	1.2% (±0.7)
Korean	**	0.5% (±0.2)	0.6% (±0.4)	0.6% (±0.4)
Japanese	**	0.4% (±0.1)	0.3% (±0.1)	0.3% (±0.2)
Other	9.3% (±3.1)	4.5% (±1.0)	4.6% (±1.0)	3.9% (±1.0)

Notes:

- “**” indicate that the answer choice was not included on the survey.
- Individual race/ethnic groups are reported for students who only select a single race/ethnic group. Students who select more than one group are reported as “more than one race/ethnicity marked”.

3. Physical Activity and Dietary Behavior

Obesity and Overweight

The Healthy People 2020 objective is to reduce the proportion of adolescents ages 12–19 who are obese, determined using a body mass index (BMI) based on height and weight, to 16.1 percent by 2020. The Healthy People 2020 BMI goal is based on measured height and weights and may not be comparable to obesity based on self-reported heights and weights from the Healthy Youth Survey.¹ For adults, self-reported heights and weights tend to underestimate obesity. Results WA aims to increase the percentage of 10th graders with healthy weight from 75% to 76% by 2016.

In 2014, 9 percent of Grade 8, and 11 percent of 10 and 12 students were obese based on their reported BMI. Fourteen percent of Grade 8 and 10 students and 13 percent of Grade 12 students were overweight.

Differences by grade level:


- There were no differences in obesity or overweight by grade level.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to be obese.
- There were no differences in overweight by gender.

Differences over time:

- There were no significant changes in obesity or overweight from 2012 to 2014.
- There were no significant trends for obesity among students in grades 8, 10 and 12 from 2002 through 2014. Among Grade 12 students, there was an increase from 2002 to 2006 and a decrease from 2006 to 2014 for overweight.



Survey Questions:

- How tall are you without your shoes on?
- How much do you weigh without your shoes on?

Note: Findings based on reported Body Mass Index (BMI) ratings calculated from height and weight, see footnote on the bottom of the page.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Obese and overweight are based on age and gender specific growth charts developed by the Centers for Disease Control and Prevention (Kuzmarski, Ogden, Grummer-Strawn, et al., 2000). Body mass index is obtained by dividing a person's weight (in kilograms) by the square of his or her height (in centimeters). Individuals in the top 5 percent for body mass index (based on the growth charts) are considered obese and those in the top 15 percent, but not the top 5 percent, are considered overweight. This is a change from 2006 and earlier years, when these categories were called overweight and at risk for overweight, respectively.

Exercise and Physical Activity

60 Minutes of Exercise Daily

In 2014, 29 percent of Grade 6 students, 31 percent of Grade 8 students, 24 percent of Grade 10 students and 21 percent of Grade 12 students reported that they were physically active for 60 minutes, seven days a week.

The Centers for Disease Control and Prevention recommends that children and adolescents participate in at least 60 minutes of physical activity daily, and muscle strengthening 3 days a week.

Differences by grade level:


- Grade 6 and 8 students were more likely than Grade 10 and 12 students to be physically active for 60 minutes seven days a week.

Differences by gender:

- Grades 6, 8, 10 and 12 males were more likely than females to be physically active for 60 minutes seven days a week.

Differences over time:

- Among Grade 6 and 8 students, there were increases in being physically active for 60 minutes seven days a week from 2012 to 2014.
- Among Grade 8, 10 and 12 students there were increases in being physically active for 60 minutes seven days a week from 2006 through 2014.



Survey Question: In the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increases your heart rate or makes you breathe hard some of the time.)

Note. Percentages represent students who reported they were physically active for 60 minutes on 7 days in an average week.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Physical Education Classes

In 2014, 41 percent of Grade 8 students, 26 percent of Grade 10 students, and 22 percent of Grade 12 students reported that they participated in a physical education class every day during an average school week.

The Healthy People 2020 objective for physical education is that 36.6 percent of adolescents in grade 9 through 12 should participate in daily school physical education classes (five days a week).

Differences by grade level:


- Grade 8 students were more likely than Grade 10 or 12 students to report participation in physical education classes every day during an average school week.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to participate in physical education classes every day during an average school week.

Differences over time:

- There were no significant changes in participation in physical education classes from 2012 to 2014.
- Among Grade 10 students, there was a decrease in participation in physical education classes on five days a week from 2002 to 2014.



Survey Question: In an average week when you are in school, on how many days do you go to physical education (PE) classes?

Note: Percentages represent students who reported they participated in five days of physical education classes in an average week when in school.

Source: YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Time Spent in Physical Education Classes

In 2014, among students who took physical education, 90 percent of Grade 8 students, 91 percent of Grade 10 and 12 students reported spending more than 20 minutes of an average physical education (PE) class actually exercising or playing sports.

Differences by grade level:


- There were no differences in spending an average of 20 minutes of an average PE class exercising by grade level.

Differences by gender:

- There were no differences in spending an average of 20 minutes of an average PE class exercising by gender.

Differences over time:

- There were no significant changes in spending an average of 20 minutes of an average PE class exercising from 2012 to 2014.
- Among Grade 8 and 10 students, there were increases in spending an average of 20 minutes of an average PE class exercising from 2002 through 2014.



Survey Question: During an average PE class, how many minutes do you spend actually exercising or playing sports?

Notes:

- Percentages represent students who reported they participated in physical education and exercised for more than 20 minutes during physical education classes.
- Students who reported that they “do not take PE” were not included in the results.
- The sample sizes for the 2014 results in this figure are: 3,653 Grade 8, 1,894 Grade 10, and 1,300 Grade 12 students.

Source: YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Television Watching and Video Game Playing

In 2014, about 53 percent of Grade 6 students, 42 percent of Grade 8 and Grade 10 students, and 45 percent of Grade 12 students reported restricting television and video viewing and video game playing to two hours or less on a school day.


A Healthy People 2020 objective is that at least 74 percent of students in grades 9 through 12 restrict television and video viewing and video game playing to two hours or less on a school day.

Differences by grade level:

- Grade 6 students more likely than Grade 8, 10 and 12 to restrict television and video viewing and video game playing to two hours or less on a school day.

Differences by gender:

- Grade 6 and 12 males were less likely than females to restrict television and video viewing and video game playing to two hours or less on a school day.



Survey Questions:

- On the average school day, how many hours did you watch TV?
- On an average school day, how many hours do you play video or computer games or use a computer for something that is not school work? (Count time spent on such things as Xbox, PlayStation, an iPod, an iPad or other tablet, a smartphone, YouTube, Facebook or other social networking tools, and the Internet.)

Notes:

- Percentages represented students who reported watching and playing video games for less than two hours on an average school day.
- The question wording for both questions changed in 2014, making changes over time no longer comparable. The questions asked from 2006 to 2012 were:
 - On the average school day, how many hours did you watch television, including videos and DVDs?
 - On an average school day, how many hours do you play video games or use a computer for fun? (Include activities such as Nintendo, Game Boy, Play Station, computer games, and the Internet.)

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Nutrition

Fruit and Vegetable Consumption

Youth need to eat a variety of fruits and vegetables every day to get essential vitamins and minerals, fiber, and other substances that are important for good health and to reduce the risk of obesity and chronic diseases. The 2010 U.S. Dietary Guidelines for Americans recommend eating sufficient amounts of fruits and vegetables within caloric needs rather than the previous recommendation of five servings for all calorie levels. For example, the USDA MyPyramid recommends daily intake of 2-3 cups of vegetables and 1.5-2 cups of fruits for youth. The Healthy Youth Survey does not measure intake of fruits and vegetables relative to caloric need and age but in terms of number of times fruits and vegetables are eaten a day. (U.S. Department of Health and Human Services, 2015)

In 2014, 24 percent of Grade 8 students, and 22 percent of Grade 10 and 12 students ate fruit and vegetables five or more times a day.

Differences by grade level:


- Grade 8 students were more likely than Grade 10 and 12 students to eat fruit and vegetables five or more times a day.

Differences by gender:

- Grade 8 and 10 males were more likely than females to eat fruit and vegetables five or more times a day over the past seven days.

Differences over time:

- Among Grade 10 students, there was a decrease in eating fruit and vegetables five or more times a day over the past seven days from 2012 to 2014.
- There were no significant trends in eating fruit and vegetables five or more times a day over the past seven days among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Questions: During the past 7 days, how many times did you?:

- Drink 100% fruit juice such as orange juice, apple juice or grape juice? (Do not count punch, Kool-Aid, sports drinks, and other fruit-flavored drinks.)
- Eat fruit? (Do not count fruit juice.)
- Eat green salad?
- Eat potatoes? (Do not count French fries, fried potatoes, or potato chips.)
- Eat carrots?
- Eat other vegetables? (Do not count green salad, potatoes, or carrots.)

Source: HYS 2002, 2004, 2006, 2008, 2012, and 2014.

Note. Percentages are calculated from the questions above to represent students who ate fruit or vegetables five or more times a day.

Eating Dinner with Family

In 2014, 75 percent of Grade 6 students, 68 percent of Grade 8 students, 61 percent of Grade 10 students, and 53 percent of Grade 12 students reported eating dinner with their family most of the time or always.

Children and adolescents who eat meals with family are more likely to have healthy eating habits.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was less likely to eat dinner with their family most of the time or always.

Differences by gender:

- Grade 6, 8, and 10 males were more likely than females to eat dinner with their family most of the time or always.

Differences over time:

- There were no significant changes in eating dinner with the family from 2012 to 2014.
- Among Grade 10 students, there was a decrease from 2002 to 2008 and an increase from 2008 to 2014 in eating dinner with the family most of the time or always. Among Grade 12 students there was an increase in eating dinner with the family from 2002 through 2014.



Survey Question: How often do you eat dinner with your family?

Note: Percentages represent students who reported they ate dinner with their family most of the time or always.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Drinking Sweetened Beverages

In 2014, 7 percent of Grade 6 students, 10 percent of Grade 8 students, 11 percent of Grade 10 students, and 12 percent of Grade 12 students reported drinking sweetened beverages two or more times a day.


Drinking sugar-sweetened beverage is associated with obesity and dental caries.

Differences by grade level:

- There were no differences in drinking two or more sweetened beverages daily by gender.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to report drinking sweetened beverages two or more times a day.



Survey Question: During the past 7 days, how many times did you drink regular soda, sports drinks (such as Gatorade) and other flavored sweetened drinks (such as Snapple or SoBe)? Do not include diet drinks.

Note. Percentages represent students who reported they consumed two or more sweetened beverages daily.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Drinking Sweetened Drinks at School

In 2014, 40 percent of Grade 8 students, 49 percent of Grade 10 students, and 44 percent of Grade 12 students reported drinking sweetened drinks at school.

Differences by grade level:


- Grade 10 students were more likely than Grade 8 and 12 students to drink sweetened drinks at school.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to drink sweetened drinks at school.

Differences over time:

- Among Grade 8, 10 and 12 students, there were decreases in drinking sweetened drinks at school in the past week from 2012 to 2014.
- Among Grade 10 and 12 students, there were decreases in drinking sweetened drinks at school in the past week from 2006 through 2014.



Survey Question: During the past 7 days, how many times did you drink regular soda, sports drinks (such as Gatorade) and other flavored sweetened drinks (such as Snapple or SoBe) at school (including any after-school and weekend activities)? Do not include diet drinks.

Notes: Percentages represent students who reported that they drank soda or sweetened drinks at school in the past 7 days.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Buying Sweetened Drinks at School

In 2014, 40 percent of Grade 8 students, 49 percent of Grade 10 students, and 44 percent of Grade 12 students reported drinking sweetened drinks at school. Among those who reported drinking these beverages at school in 2014, 17 percent of Grade 8 and 12 students, and 25 percent of Grade 10 students bought the sweetened drinks at school.

Differences by grade level:


- Among those who drank sweetened beverages at school, Grade 10 students were more likely than Grade 8 and 12 students to buy the sweetened drinks at school.

Differences by gender:

- Among those who drank sweetened beverages at school, Grade 8 females were more likely than males and Grade 12 males were more likely than females to buy the sweetened drinks at school.

Differences over time:

- Among Grade 10 and 12 students who drank sweetened beverages at school, there were decreases in buying sweetened drinks at school.
- Among Grade 8, 10 and 12 students who drank sweetened beverages at school, there were decreases in buying sweetened drinks at school.



Survey Question: During the past 7 days, where did you usually get the soda or other sweetened drinks that you drank at school? (Choose only one answer.)

Notes:

- Percentages represent students who reported that they bought the soft drinks at school.
- Students who reported that they “did not drink sodas, sports drinks, or other flavored drinks at school” in the past, 7 were not included in the source of soda results.
- The sample sizes for the 2014 results in this figure are: 2,091 Grade 8, 2,276 Grade 10, and 1,589 Grade 12 students.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Food Insecurity

In 2014, 11 percent of Grade 8, 13 percent of Grade 10, and 14 percent of Grade 12 students reported food insecurity.

Compared to children from families who are food secure, children from families with food insecurity are more likely to have behavior problems, do poorly in school, need medical care and hospitalization, and to develop chronic diseases (Center on Hunger and Poverty, 2002; Hampton, 2007). Food insecurity may also be associated with poor quality diet and obesity (Lobstein, 2015, Weinreb, 2002).

Differences by grade level:


- Grade 8 students were less likely than Grade 10 and 12 to have cut meal size or skipped meals.

Differences by gender:

- There were no differences in to having cut meal size or skipped meals by gender.

Differences over time:

- Among Grade 8, 10 and 12 students, there were decreases in having to cut meal size or skip meals from 2012 to 2014.
- There were no significant trends in having to cut meal size or skip meals among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: How often in the past 12 months did you or your family have to cut meal size or skip meals because there wasn't enough money for food?

Notes:

- Percentages represent students who reported their family cut meal size or skipped meals in the past year due to lack of money for food.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

4. Health Status and Health Care

Asthma

Lifetime Asthma

Lifetime asthma includes anyone who has ever been told by a doctor or nurse that they have asthma. In 2014, 14 percent of Grade 6 students, 18 percent of Grade 8 students, 22 percent of Grade 10 students and 21 percent of Grade 12 students reported that they had been told they have asthma.

Differences by grade level:


- Grade 6 students were less likely than Grade 8, 10 and 12 students to have been diagnosed with asthma in their lifetime.
- Grade 8 students were less likely than Grade 10 and 12 students to have been diagnosed with asthma in their lifetime.

Differences by gender:

- There were no differences in ever being diagnosed with asthma in their lifetime by gender.

Differences over time:

- Among Grade 6 and 12 students, there were decreases in having been diagnosed with asthma in their lifetime from 2012 to 2014.
- There were no significant trends in having been diagnosed with asthma in their lifetime for any grades from 2002 through 2014.



Survey Question: Has a doctor or nurse ever told you that you have asthma?

Note. Percentages represent students who reported were ever told they had asthma by a doctor or nurse in their life.

Source: YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Current Asthma

Current asthma includes anyone who had ever been told they have asthma by a doctor or a nurse and also reports that they still have asthma. In 2014, 7 percent of Grade 6 students, 8 percent of Grade 8 and Grade 12 students, and 10 percent of Grade 10 students reported that they were told they had asthma and that they still have asthma.

Differences by grade level:


- Grade 6 students were less likely than Grade 10 and 12 students to have current asthma.
- Grade 8 students were less likely than Grade 10 students to have current asthma.
- Grade 12 students were less likely than Grade 10 students to have current asthma.

Differences by gender:

- Grade 10 and 12 females were more likely than males to have current asthma.

Differences over time:

- Among Grade 6, 8 and 12 students, there were decreases in current asthma from 2012 to 2014.



Survey Questions:

- Has a doctor or nurse ever told you that you have asthma?
- Do you still have asthma?

Notes:

- Percentages represent students who reported they were ever told they had asthma and those who were ever told they have asthma and still have asthma.
- The definition of current asthma changed in 2008, so previous results for current asthma are not comparable. In the past current asthma was defined as being diagnosed by a doctor and having an asthma attack in the past year.

Source: HYS 2008, 2010, 2012 and 2014.

Diabetes

Diabetes has lifelong implications for health and wellbeing. In 2014, 3 percent of Grade 8, 10 and 12 students reported having been told they have diabetes.

Differences by grade level:


- There were no differences in having diabetes by grade level.

Differences by gender:

- There were no differences in having diabetes by gender.

Differences over time:

- Among Grade 6, 8 and 12 students, there were decreases in diabetes from 2010 to 2014.
- There were no significant trends in diabetes among students in grades 8, 10 and 12 from 2004 through 2014.



Survey Question: Have you ever been told by a doctor or other health professional that you have diabetes?

Note: Percentages represent students who reported they were ever told they had diabetes.

Source: HYS 2004, 2006, 2008, 2010, and 2014.

Access to Care

Access to medical and dental care is an important component in creating a healthy adolescent and adult.

Access to a Doctor

In 2014, 65 percent of Grade 8 students, 66 percent of Grade 10 students, and 61 percent of Grade 12 students saw a doctor for a checkup or physical exam in the past 12 months when they were not sick or injured.

Differences by grade level:


- Grade 12 students were less likely than Grade 8 and Grade 10 students to have seen a doctor for a checkup or physical exam in the past year.

Differences by gender:

- There were no differences in seeing a doctor for a checkup or physical exam in the past year by gender.

Differences over time:

- Among Grade 8, 10, and 12 students, there were increases in seeing a doctor for a checkup or physical exam in the past year from 2010 to 2014.
- There were no significant trends in seeing a doctor for a checkup or physical exam in the past year among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: When was the last time you saw a doctor or health care provider for a check-up or physical exam when you were not sick or injured?

Note: Percentages represent students who reported they saw a doctor or health care provider when they were not sick or injured in the past year.

Source: WSSAHB 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, and 2014.

Access to a Dentist

In 2014, 77 percent of Grade 8 and 12 students, 79 percent of Grade 10 students, and 76 percent of Grade 12 saw a dentist in the past 12 months.

Differences by grade level:


- There were no differences in seeing a dentist in the past year by grade level.

Differences by gender:

- Grade 8 females were more likely than males to have seen a dentist in the past 12 months.

Differences over time:

- There were no significant changes in seeing a dentist in the past year from 2012 to 2014.
- Among Grade 8 students, there was an increase in seeing a dentist in the past year from 2002 through 2014.



Survey Question: When was the last time you saw a dentist for a check-up, exam, teeth cleaning, or other dental work?

Note: Percentages represent students who reported they saw a dentist in the past year.

Source: WSSAHB 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

5. Mental Health

Depressive Feelings

Students were asked, “During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?” Although this question is not sufficient to diagnose depression, it can be used as a surrogate measure for experiencing symptoms of depression (Merikangas, 2009).

In 2014, 27 percent of students in Grade 8, 35 percent of students in Grades 10, and 34 percent of students in Grade 12 reported experiencing depressive feelings during the past year.

Differences by grade level:


- Grade 10 and 12 students were more likely than Grade 8 students to experience depressive feelings.

Differences by gender:

- Grade 8, 10 and 12 females were more likely than males to experience depressive feelings.

Differences over time:

- Among Grade 10 and 12 students, there were increases in experiencing depressive feelings from 2012 to 2014.
- There were no significant trends in experiencing depressive feelings among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?

Note: Percentages represent students who reported, yes, they felt sad or hopeless for two weeks or more in the past year.


Source: YRBS 1999, WSSAHB 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Suicide

Suicide attempts and suicide ideation are associated with substance abuse, sexual activity, and physical fighting (King, 2001). Attempted suicide also heightens the risk of eventual suicide.

In 2014, students reported the following suicide-related behaviors:

- Seriously considered attempting suicide in the past year: 16 percent of Grade 8, and 20 percent of Grade 10 students, and 18 percent of Grade 12 students.
- Made a plan about how to attempt suicide in the past year: 14 percent of Grade 8 and 12 students, and 16% of Grade 10 students.
- Actually attempted suicide: 9 percent of Grade 8, 10 percent of Grade 10 students, and 8 percent of Grade 12 students.
- Felt that they did not have an adult to turn to for help when feeling sad or hopeless: 8 percent of Grade 6, 13 percent of Grade 8, 15 percent of Grade 10, and 13 percent of Grade 12 students.



Survey Questions:

- During the past 12 months, did you ever seriously consider attempting suicide?
- During the past 12 months, did you make a plan about how you would attempt suicide?
- During the past 12 months, how many times did you actually attempt suicide?
- When you feel sad or hopeless, are there adults that you can turn to for help?

Notes:

- Percentages represent students who seriously considered suicide, who made a plan to attempt suicide, or who actually attempted suicide any time in the past 12 months.
- Percentages for “no adult to turn to when sad or hopeless” represent students who said “no”. In past versions of the Analytic Reports, students who said they “never feel sad or hopeless” were not included in the results.

Source: HYS 2014.

Suicide Attempts

In 2014, 9 percent of Grade 8, 10 percent of Grade 10 students, and 8 percent of Grade 12 students actually attempted suicide in the past year.

The Healthy People 2020 objective is to reduce the percentage of adolescents in grades 9 through 12 who attempt suicide from 1.9 suicide attempts per 100,000 population to 1.7 suicide attempts per 100,000 population.

Differences by grade level:


- Grade 10 students were more likely than Grade 12 students to have attempted suicide in the past year.

Differences by gender:

- Grade 8, 10 and 12 females were more likely than males to have attempted suicide in the past year.

Differences over time:

- Among Grade 10 students, there was an increase in attempting suicide in the past 12 months from 2012 to 2014.
- There were no significant trends in attempting suicide in the past 12 months among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Questions: During the past 12 months, how many times did you actually attempt suicide?

Notes:

- Percentages represent students who reported attempted suicide any time in the past 12 months.
- In 2006, the survey response options were changed from the number of times of attempted suicide to “yes” or “no” attempted suicide.

Source: WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

6. Sexual Behavior

Sexual Behavior

About 35% of schools included in the state sample elected to administer Form B-enhanced which included six optional questions on sexual orientation, behavior and abuse.

Engaging in sexual activities can result in unintended pregnancy and sexually transmitted diseases including HIV.

Lifetime Sexual Intercourse

In 2014, 9 percent of Grade 8 students, 27 percent in Grade 10 and 52 percent in Grade 12 students reported they ever had sexual intercourse. Other results for Washington youth in 2014:

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to have ever had sexual intercourse.

Differences by gender:

- Grade 8 and 10 males were more likely than females to ever have sexual intercourse.

Differences over time:

- Among Grade 8 students, there was a decrease in having sexual intercourse from 2012 to 2014.



Survey Question: How old were you when you had sexual intercourse for the first time?

Note: Percentages represent students who had ever had sexual intercourse.

Source: HYS 2010, 2012, and 2014.

Sexual Behaviors among Those who have had Sex

Sexual behaviors were reported among those who have ever had sexual intercourse:

- 14 percent of Grade 10 students and 8 percent of Grade 12 had sexual intercourse before the age 13.
- About 20 percent of Grade 10 students and 28 percent of Grade 12 students had sexual intercourse with four or more partners in their lifetime.
- About 62 percent of Grade 10 students and 54 percent of Grade 12 students used a condom the last time they had sexual intercourse.

Differences by grade level:


- Grade 10 students were more likely than Grade 12 students to have had sexual intercourse before age 13, and more likely to have used a condom the last time they had sexual intercourse.
- Grade 12 students were more likely than Grade 10 students to have had four or more partners.

Differences by gender:

- Grade 10 males were more likely than females to have had sexual intercourse before age 13.
- Grade 10 and 12 males were more likely than females to have used a condom the last time they had sexual intercourse.

Differences over time:

- There were no significant changes in having sexual intercourse, having sexual intercourse before age 13, having four or more partners, or using a condom among students in grades 10 and 12 from 2010 to 2014.



Survey Questions:

- Have you ever had sexual intercourse?
- How old were you when you had sexual intercourse for the first time?
- With how many people have you ever had sexual intercourse?
- The last time you had sexual intercourse; did you or your partner use a condom?

Notes:

- Percentages represent students who had ever had sexual intercourse, who had sex before age 13, or who had 4 or more sexual partners.
- Students who reported that they had not had sexual intercourse in their lifetime were not included in the results. The sample sizes for the 2014 results in this chart are 402 Grade 10 and 580 Grade 12 students.
- The results for Grade 8 are not reported.

Source: HYS 2010, 2012, and 2014.

Sexual Orientation


Sexual orientation was added to HYS in 2014. About 1 percent of Grade 8 students, and 2 percent of Grade 10 and 12 students identified as gay or lesbian. About 5 percent of Grade 8 students and 6 percent of Grade 10 and 12 students identified as bisexual.

Differences by grade level:

- Grade 12 students were more likely than Grade 8 students to identify as gay or lesbian.
- Grade 10 students were more likely than Grade 8 students to identify as bisexual.

Differences by gender:

- There were no differences in identifying as gay or lesbian by gender.
- Grade 8, 10 and 12 females were more likely than males to identify as bisexual.



Survey Questions: Which of the following best describes you?

Note: Percentages represent students who identified as heterosexual (straight), gay or lesbian, bisexual or were not sure.

Source: HYS 2014.

Sex of Sexual Contacts

Sex of sexual contacts was added to HYS in 2014. Among those who had sexual intercourse:


- About 1 percent of Grade 8 students, 2 percent of Grade 10 students and 4 percent of Grade 12 students reported having same-sex contact.
- About 2 percent of Grade 8 students, 6 percent of Grade 10 students and 7 percent of Grade 12 students reported having contact with both females and males.

Differences by grade level:

- Grade 12 students were more likely than Grade 8 and 10 students to report same-sex contact.
- Grade 10 and 12 students were more likely than Grade 8 students to report having contact with both females and males.

Differences by gender:

- There were no differences in reporting same sex contact by gender*.
- Grade 10 and 12 females were more likely than males to report having contact with both females and males.



Survey Questions: During your life, with whom have you had sexual contact?

Notes:

- Percentages represent students who never had sexual contact, females that had contact with males/males that had contact with females (heterosexual contact), females that had contact with females/males that had contact with males (same-sex contact) and students who had contact with both females and males.
- *Grade 8 comparisons by gender were suppressed due to small numbers.

Source: HYS 2014.

7. School Climate

School Safety, Bullying, and Harassment

RCW 28A.320.185 requires all public school districts and public schools to have current safety plans and procedures in place. State legislators, the Governor, the state education agency, local schools and communities, and parents recognize that students must feel safe at school to be successful learners. Effective school safety plans that include bullying and harassment prevention programs challenge traditional cultural norms that might condone bullying as a normal part of growing up.

Feeling Safe at School

When students feel safe at school, they are more likely to make better grades compared to those students who do not feel safe at school (Dilley 2009). In 2014, 89 percent of Grade 6 students, 86 percent of Grade 8 students, 85 percent of Grade 10 students, and 87 percent of Grade 12 students felt safe at school.

Differences by grade level:


- Grade 6 students were more likely than Grade 8 and 10 to feel safe at school.

Differences by gender:

- Grades 6 and 12 females were more likely than males to feel safe at school.

Differences over time:

- There were no significant changes in feeling safe at school from 2012 to 2014.
- Among Grade 8 students, there was an increase in feeling safe at school from 2002 through 2014.



Survey Question: I feel safe at my school.

Notes:

- Survey forms A and B have different response options.
- Percentages represent students who reported yes or mostly true, or, YES! or definitely true, that they felt safe at school.

Source: WSSAHB 1995, 1998 and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Bullying

Bullying is defined as a student or group of students saying or doing nasty or unpleasant things to another student. Under this definition bullying includes teasing a student repeatedly in a way he or she does not like but does not include two students of about the same strength quarreling or fighting. The definition of bullying includes electronic forms of bullying, that is, cyberbullying.

Students who are bullied at school are more likely to get lower grades compared to those who are not bullied. Creating a safe environment is critical for students' academic achievement. Research has identified best practice support programs that address school harassment and bullying and build positive school culture (Smith, Pepler, and Rigby, 2004). In 2014, 31 percent of Grade 6 students, 28 percent of Grade 8 students, 23 percent of Grade 10 students, and 16 percent of Grade 12 students were bullied at school.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was less to be bullied.

Differences by gender:

- Grade 6, 8, 10 and 12 females were more likely than males to have been bullied.

Differences over time:

- Among Grade 8, 10 and 12 students, there were decreases in bullying from 2012 to 2014.
- Among Grade 12 students, there was an increase in bullying from 2002 through 2014.



Survey Question: A student is being bullied when another student, or group of students, say or do nasty or unpleasant things to him or her. It is also bullying when a student is teased repeatedly in a way he or she doesn't like. It is NOT bullying when two students of about the same strength argue or fight. In the last 30 days, how often have you been bullied?

Note: Percentages of students who reported they were bullied on any days in the last 30 days.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Harassment by Computer or Cell Phone

Many schools have modified their policy and procedures to specifically address harassment. In addition, computer or cell phone harassment is also included in bullying and harassment policies and procedures.

In 2014, 13 percent of Grade 8, 12 percent of Grade 10 students and 11 percent of Grade 12 students reported being harassed with a computer or cell phone in the past 30 days.

Differences by grade level:


- Grade 8 students were more likely than Grade 10 students to be harassed with a computer or cell phone.

Differences by gender:

- Grade 8, 10 and 12 females were more likely than males to be harassed with a computer or cell phone.

Differences over time:

- Among Grade 8 students, there was an increase in being harassed with a computer or cell phone from 2012 to 2014.
- There were no significant trends in being harassed with a computer or cell phone among students in grades 8, 10 and 12 from 2006 through 2014.



Survey Question: In the past 30 days, has someone used the computer or a cell phone to bully, harass or intimidate you?

Notes: Percentages represent students who were harassed with a computer or cell phone on any days.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Harassment Due to Perceived Sexual Orientation

In 2014, 11 percent of Grade 8, 9 percent of Grade 10, and 7 percent of Grade 12 students reported being harassed because someone thought they were gay, lesbian or bisexual

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was less likely to be harassed due to perceived sexual orientation.

Differences by gender:

- There were no differences in being harassed due to perceived sexual orientation by gender.

Differences over time:

- There were no significant changes in being harassed due to perceived sexual orientation from 2012 to 2014.
- Among Grade 10 students, there was a decrease in being harassed due to perceived sexual orientation from 2006 through 2014.



Survey Question: In the past 30 days, how often were you bullied, harassed, or intimidated at school or on your way to or from school because someone thought you were gay, lesbian or bisexual (whether you are or are not)?

Notes: Percentages represent students who reported being harassed due to their perceived sexual orientation on any days.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Weapon Carrying at School

School safety requires the commitment of staff members, students, parents and the community. Creating a safe and supportive learning environment is critical for student academic success (Dilley, 2009). In 2014, 3 percent of Grade 6, 4 percent of Grade 8, and 6 percent of Grade 10 and Grade 12 students reported weapon carrying at school in the past 30 days.

Differences by grade level:


- Grade 6 students were less likely than Grade 8, 10, and 12 students to carry a weapon at school in the past 30 days.
- Grade 8 students were less likely than Grade 10, and 12 students to carry a weapon at school in the past 30 days.

Differences by gender:

- Grade 6, 8, 10 and 12 males were more likely than females to carry a weapon at school in the past 30 days.

Differences over time:

- There were no significant changes in weapon carrying at school from 2012 to 2014.
- This estimate has been stable since 2002 among 6th graders, but has decreased among older adults. Among Grade 8 students there was a decrease from 2008 to 2014 in weapon carrying at school. Among Grade 10 students, there was a decrease from 2006 to 2014 in weapon carrying at school. Among Grade 12 students, there was an increase in weapon carrying at school from 2002 to 2006, and a decrease from 2006 to 2014.



Survey Question: During the past 30 days, did you carry a weapon such as a gun, knife, or club on school property?

Notes:

- Percentages represent students who reported any weapon carrying at school in the past 30 days.
- Grade 6 students were asked if they carried a weapon at school, “yes” or “no.”
- Grade 8, 10 and 12 students were asked the number of times they carried a weapon.
- In 2006, the response options were reduced from 5 different numbers of times options to 3 different numbers of times.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Substance Use at School

The use of substances at school significantly affects student learning and compromises the school environment. Substance use and abuse are closely correlated with violent behavior (Office of National Drug Control Policy, 2007). Prevention, early intervention, treatment, and other related efforts that reduce the number of students engaging in these behaviors and coming to school high or drunk enhances school safety and increases student potential for academic success.

Alcohol or Other Drug Use on School Property

The National Center for Education Statistics (2013) track alcohol and marijuana use as Indicators of School Crime and Safety relying on data collected from the Youth Risk Behavior Surveillance System (YRBS). In 2011, 6 percent of students surveyed in grades 9–12 reported using marijuana on school property while three percent reported using alcohol on school property during the past 30 days.

In 2014, 6 percent of Grade 8 students, 14 percent of Grade 10 students, and 18 percent of Grade 12 students reported being drunk or high at school in the past year.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to be drunk or high at school in the past year.

Differences by gender:

- Grade 12 males were more likely than females to report being drunk or high at school in the past year.

Differences over time:

- Among Grade 8 students, there was a decrease in being drunk or high at school from 2012 to 2014.
- There were no significant trends in being drunk or high at school among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: How many times in the past year (12 months) have you been drunk or high at school?

Note: Percentages represent students who reported being drunk or high on school property on any days in the past year.

Source: WSSAHB 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Alcohol Drinking on School Property


In 2014, 3 percent of Grade 8 students, 6 percent of Grade 10 students and Grade 12 students reported drinking alcohol at school in the past 30 days. Among those who reported drinking alcohol on any days in the past 30 days, 21 percent of Grade 8 students, 20 percent of Grade 10 students, and 14 percent of Grade 12 students reported drinking alcohol at school in the past 30 days

Differences by grade level:

- Grade 10 and 12 students were more likely than Grade 8 students to drink alcohol at school in the past 30 days.

Differences by gender:

- Grade 10 and 12 males were more likely than females to drink alcohol at school in the past 30 days.



Survey Question: During the past 30 days, on how many days did you have at least one drink of alcohol on school property?

Note: Percentages represent students who reported drinking alcohol on school property on any days in the past 30 days.

Source: HYS 2014.

Marijuana Use on School Property


In 2014, 3 percent of Grade 8 students, 7 percent of Grade 10 students, and 8 percent of Grade 12 students reported using marijuana at school in the past 30 days. Among those who reported using marijuana on any days in the past 30 days, 29 percent of Grade 8 students, 30 percent of Grade 10 students, and 24 percent of Grade 12 students reported using marijuana at school in the past 30 days

Differences by grade level:

- Grade 10 and 12 students were more likely than Grade 8 students to use marijuana at school in the past 30 days.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to use marijuana at school in the past 30 days.



Survey Question: During the past 30 days, on how many days did you use marijuana on school property?

Note: Percentages represent students who reported using marijuana on school property on any days in the past 30 days.

Source: HYS 2014.

Perceived Availability of School Staff to Discuss Substance-Related Problems

Students who have opportunities for interaction with school staff, especially in times of crisis, are more likely to be connected to school and academically successful (Catalano, Haggerty, Oesterle, Fleming, and Hawkins, 2004). In 2014, 65 percent of Grade 8, and 56 percent of Grade 10 and Grade 12 students reported having someone at school with whom to discuss substance-related problems.

Differences by grade level:


- Grade 8 students were more likely than Grade 10 and 12 students to have someone at school to discuss substance-related problems.

Differences by gender:

- Grade 8 females were more likely than males to have someone at school to discuss substance-related problems.

Differences over time:

- There were no significant changes in having someone at school to discuss substance-related problems from 2012 to 2014.
- Among Grade 8, 10 and 12 students, there were decreases in having someone at school to discuss substance-related problems from 2002 through 2014.



Survey Question: Does your school provide a counselor, intervention specialist, or other school staff member for students to discuss problems with alcohol, tobacco, or other drugs?

Note: Percentages represent students who were aware of having someone at school with whom they could discuss substance-related problems. Those who answered "I'm not sure" were considered not aware.

Source: WSSAHB 1995 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

School Attendance

A significant portion of young people’s lives is spent attending school. When youth enjoy school and attend regularly, they are more likely to achieve academically and are at much less risk of engaging in a variety of at-risk behaviors.

Skipping or Cutting School

In 2014, 17 percent of Grade 6 students, 15 percent of Grade 8 students, 18 percent of Grade 10 students, and 28 percent of Grade 12 students reporting skipping or cutting at least one day of school in the past 30 days.

Differences by grade level:


- Grade 10 students were more likely than Grade 8 students to skip or cut a whole day of school in the past 30 days.
- Grade 12 students were more likely than Grade 6, 8 and 10 students to skip or cut a whole day of school in the past 30 days.

Differences by gender:

- Grade 6 males were more likely than females to skip or cut a whole day of school in the past 30 days.

Differences over time:

- For Grade 12, there was an increase in skipping school from 2012 to 2014.
- There were no significant trends in skipping school for any grades from 2002 through 2014.



Survey Question: During the LAST 4 WEEKS, how many whole days of school have you missed because you skipped or “cut”?

Notes:

- Percentages represent students who reported they skipped or cut any days of school in the past 30 days.
- This question was not asked of Grade 6 students in 2002, 2004 and 2006, but was added back on the survey in 2008.

Source: WSSAHB 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Enjoying School

Students who report a positive attitude toward school are more likely to be academically successful (Catalano, Haggerty, Oesterle, Fleming, and Hawkins, 2004). In 2014, 31 percent of Grade 6, 18 percent of Grade 8, and 11 percent of Grade 10, and 10 percent of Grade 12 students reported almost always enjoying school over the past year.

Differences by grade level:


- Grade 6 students are more likely than Grade 8, 10 and 12 students to almost always enjoy school.
- Grade 8 students are more likely than Grade 10 and 12 students to almost always enjoy school.

Differences by gender:

- Grade 6 and 8 females were more likely than males to almost always enjoy school.

Differences over time:

- Among Grade 8, 10, and 12 there were decreases in almost always enjoying school from 2012 to 2014.
- There were no significant trends in almost always enjoying school for any grades from 2002 through 2014.



Survey Question: Think back over the past year in school. How often did you: Enjoy being in school?

Note: Percentages represent students who reported they almost always enjoy school.

Source: WSSAHB 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

8. Unintentional Injury Behaviors

Motor Vehicle Safety

Riding with a Drinking Driver

Impaired driving is a strong risk factor for a fatal crash. At all levels of blood alcohol concentration (BAC), the risk of involvement in a motor vehicle crash is greater for teens than for older drivers. Among drivers between 15 and 20 years of age who were involved in fatal crashes in 2012, 23% had been drinking (National Highway Traffic Safety Administration, 2012). The Healthy People 2020 objective is to reduce the percentage of adolescents in grades 9 through 12 who report riding with a driver who has been drinking alcohol from 28.3 to 25.5 percent.

In 2014, 6 percent of Grade 6 students, 17 percent of Grade 8 students, 18 percent of Grade 10 and 17 percent of Grade 12 students reported riding in a car driven by someone who had been drinking alcohol.

Differences by grade level:


- Grade 6 students were less likely than Grade 8, 10 and 12 students to ride in a vehicle driven by someone who had been drinking alcohol.

Differences by gender:

- Grade 8 and 10 females were more likely than males to ride in a vehicle driven by someone who had been drinking alcohol.

Differences over time:

- Among Grade 12 students, there was a decrease in riding in a vehicle driven by someone who had been drinking alcohol from 2012 to 2014.
- Among Grade 10 students, there was a decrease from 2008 to 2014 in riding in a vehicle driven by someone who had been drinking alcohol. Among Grade 12 students, there was a decrease in riding in a vehicle driven by someone who had been drinking alcohol from 2002 through 2014.



Survey Questions:

- Secondary: During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been drinking alcohol?
- Primary: In the last 30 days, have you ridden in a car driven by someone who had been drinking alcohol?

Note: Percentages represent students who reported that they rode in a vehicle in the past 30 days whose driver had been drinking alcohol.

Source: WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Drinking and Driving

In 2014, 5 percent of Grade 10 students and 9 percent of Grade 12 students reported drinking alcohol and driving in the past 30 days. Among those who drank alcohol on any days in the past 30 days, 20 percent of Grade 10 students and 17 percent of Grade 12 students report driving after drinking alcohol.

Differences by grade level:


- Grade 12 students were more likely than Grade 10 students to report driving a vehicle after drinking alcohol.

Differences by gender:

- Grade 10 and 12 males were more likely than females to report driving a vehicle after drinking alcohol.

Differences over time:

- Among Grade 12, there was a decrease in driving a vehicle after drinking alcohol from 2012 to 2014.
- Among Grade 12 students, there was a decrease in driving a vehicle after drinking alcohol from 2002 through 2014.



Survey Question: During the past 30 days, how many times did you drive a car or other vehicle when you had been drinking alcohol?

Notes:

- Percentages represent students who reported driving and drinking alcohol any times in the past 30 days.
- The results for Grade 8 students are not reported due to the fact that most are not old enough to drive.

Source: WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Marijuana Use – Riding and Driving

Research indicates impairment in driving after recent smoking or with blood THC serum concentrations 2-5 ng/mL, particularly for infrequent users of cannabis (Skopp, 2003; Hartman, 2013; Hammond, 2014). In addition, there is a higher risk of auto accidents for drivers under the influence of both alcohol and marijuana than under the influence of one substance alone (Dubois, 2015).

In 2014, 10 percent of Grade 8 students, 19 percent of Grade 10 students, and 26 percent of Grade 12 students report riding in a vehicle with someone who used marijuana in the past 30 days.


In 2014, 9 percent of Grade 10 students and 17 percent of Grade 12 students report driving within three hours after using marijuana. Among those who used marijuana on any days in the past 30 days, 35 percent of Grade 10 students and 52 percent of Grade 12 students report driving within three hours after using marijuana.

Differences by grade level:

- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to ride with a driver who used marijuana.
- Grade 12 students were more likely than Grade 10 students to drive within three hours after using marijuana.

Differences by gender:

- Grade 8 females were more likely than males to report riding with a driver who used marijuana.
- Grade 10 and 12 males were more likely than females to report driving within three hours after using marijuana.



Survey Questions:

- During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been using marijuana?
- During the past 30 days, how many times did you drive a car or other vehicle within three hours after using marijuana?

Notes:

- Percentages represent students who reported driving after using marijuana any times within three hours in the past 30 days
- Percentages represent students who reported riding in a vehicle driven by someone who used marijuana.
- The results for Grade 8 students are not reported due to the fact that most are not old enough to drive (NR, not reported).

Source: HYS 2014.

Texting or Emailing – Riding and Driving

In 2014, 27 percent of Grade 6, 47 percent of Grade 8 students, 57 percent of Grade 10 students, and 59 percent of Grade 12 students report riding in a vehicle with someone who was texting or emailing while driving in the past 30 days.


In 2014, 8 percent of Grade 10 students and 43 percent of Grade 12 students report driving while texting or emailing in the past 30 days.

Differences by grade level:

- Grade 8, 10 and 12 students were more likely than Grade 6 students to ride with someone texting or emailing.
- Grade 10 and 12 students were more likely than Grade 8 students to ride with someone texting or emailing.
- Grade 12 students were more likely than Grade 10 students to drive while texting or emailing.

Differences by gender:

- Grade 8, 10 and 12 females were more likely than males to ride with someone texting or emailing.
- Grade 10 males were more likely than females, and Grade 12 females were more likely than males to drive while texting or emailing.



Survey Questions:

- For Grades 8, 10, and 12: During the past 30 days, how many days did you ride in a car or other vehicle driven by someone who was texting or emailing? For Grade 6: During the past 30 days, did you ride in a car or other vehicle driven by someone who was texting or emailing?
- During the past 30 days, how many days did you text or email while driving a car or other vehicle?

Notes:

- Percentages represent students who rode with a texting or emailing driver in the past 30 days.
- Percentages represent students who were driving while texting or emailing in the past 30 days.
- The results for Grade 8 students are not reported due to the fact that most are not old enough to drive (NR, not reported).

Source: HYS 2014.

Swimming Safety

There is one study that shows swimming lessons reduce drowning risk (Brenner, 2009).


In 2014, 56 percent of Grade 6 students, 59 percent of Grade 8 and 10 students, and 64 percent of Grade 12 students had taken formal swimming lessons. In 2014, 85 percent of Grade 6 students, 87 percent of Grade 8, students, and 86 percent of Grade 10 and 12 students felt comfortable playing or swimming in water over their head.

Differences by grade level:

- Grade 12 students were more likely than Grade 6 students to have taken formal swim lessons.
- There were no differences in feeling comfortable playing or swimming in deep water by grade level.

Differences by gender:

- Grade 6 and 12 females were more likely than males to have taken formal swim lessons.
- Grade 10 and 12 males were more likely than females to report feeling comfortable playing or swimming in deep water.



Survey Questions:

- Have you ever taken formal swimming lessons?
- I am comfortable playing and swimming in water over my head.

Notes:

- Percentages represent students who reported “yes” they had formal swim lessons.
- Percentages represent students who “Agree” or “Strongly agree” they feel comfortable playing or swimming in water over their head.

Source: HYS 2014.

Boat Safety

Drowning is the second leading cause of unintentional injury death for children in Washington. Most Washington State drownings occur in open water such as lakes, rivers, and the ocean. However, less than half of teens wear life vests while riding in small boats. In 2014, 42 percent of the Grade 8 students, 32 percent of the Grade 10 students, and 27 percent of the Grade 12 students who go boating reported always wearing a life vest.

Differences by grade level:


- Among Grade 8, 10 and 12 students who boat, as grade levels increase, each grade was less likely to always wear a life vest when boating.

Differences by gender:

- Grade 12 males who boat were less likely than females to always wear a life vest when boating.

Differences over time:

- Among Grade 8, 10 and 12 students who boat, there were decreases in wearing a life vest while boating from 2012 to 2014.
- There were no significant trends in wearing a life vest while boating among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: How often do you wear a life vest when you're in a small boat like a canoe, raft, or small motorboat?

Notes:

- Percentages represent students who boat and reported always wearing a life vest when in a small boat such as a canoe, raft, or small motor boat.
- Students who reported that they "never go boating" were not included in the results. The sample sizes for the 2014 results in this chart are 4,122 Grade 8; 3,602 Grade 10; and 2,732 Grade 12 students.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

9. Intentional Injury Behaviors

Physical Fighting

Physical fighting, a common form of interpersonal violence among teens, is a public health concern both because of the potential for fight-related injuries and its association with participation in many other health risk behaviors.

In 2014, 23 percent of Grade 6 students, 27 percent of Grade 8 students, 22 percent of Grade 10 students, and 16 percent of Grade 12 students reported being in a physical fight in the past year.

The Healthy People 2020 objective is to reduce physical fighting in the past year among adolescents in grades 9 through 12 to 28.4 percent.

Differences by grade level:


- Grade 6 students were more likely than Grade 12 students to be in a physical fight.
- Grade 8 students were more likely than Grade 6, 10 and 12 students to be in a physical fight.
- Grade 10 students were more likely than Grade 12 students to be in a physical fight.

Differences by gender:

- Grade 6, 8, 10 and 12 males were more likely than females to be in a physical fight in the past year.

Differences over time:

- Among Grade 8 and 12 students, there were decreases in physical fighting from 2012 to 2014.
- Among Grade 6 and 8 students, there were decreases in physical fighting from 2002 through 2014. Among Grade 10 students, there was a decrease in physical fighting from 2008 to 2014.



Survey Question:
During the past 12 months, how many times were you in a physical fight?

Note: Percentages represent students who reported being in a physical fight in the past year.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Gangs

Youth gangs are responsible for a substantial portion of serious violence in the United States and commit a disproportionate share of offenses (Egley et al., 2012). In schools and neighborhoods where gangs are active, gangs create a climate of fear, and increase the amount of violence and criminal behavior.


In 2014, 6 percent of Grade 8 and 10 students, and 5 percent of Grade 12 students reported being in a gang in the past year. In 2014, 11 percent of Grade 8, 19 percent of Grade 10 and 17 percent of Grade 12 students reported that there are gangs at their school.

Differences by grade level:

- Grade 10 students were more likely than Grade 12 students to have been a gang membership in the past year.
- Grade 8 students were less likely than Grade 10 and 12 students to report gangs at their school.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to have been a gang member in the past year.
- Grade 10 and 12 males were less likely than females to report gangs at their school.



Survey Questions:

- During the past 12 months, have you been a member of a gang?
- Are there gangs at your school?

Notes:

- Percentages represent students who reported “yes” they were a member of a gang in the past 12 months. Trend results for gang membership are not reported because the introduction to the gang membership question changed in 2014 to include “A gang is a group of people with a leader who act together often for violent or illegal activities.”
- Percentages represent students who reported “yes” there are gangs at their school.

Source: HYS 2014.

10. Alcohol, Tobacco, and Other Drug Use

Current Substance Use

Student responses to questions about substance use in the past 30 days are indicators of their current substance use. This section presents current (30-day) prevalence results by grade from 1988 to 2014 (see Tables 4 through 7). Binge drinking in the past 2 weeks is also included in these tables. Detailed results for individual substances appear in subsequent sections.

The prevalence of current use for some substances has been assessed differently as survey instruments have changed over time. Superscripts in the tables describe any changes to survey questions or responses. Therefore, readers should use caution when making strong conclusions about changes over time for these substances.

In addition, it is important to recognize that these results are based on responses from students attending public schools. Rates of substance use may be different in other educational settings, and are likely higher among youth who have dropped out of school.

Description of Superscript Notes for Tables 4 through 7

How the question was asked and changes over time:

1. Question asked as “how often did you use . . .”
2. Question asked as “during the past 30 days, how many times have you . . .”
3. Question asked as “during the past 30 days, on how many days did you . . .”
4. Question asked as “think back over the past two weeks, how many times have you . . .”
5. Question asked as “during the past 30 days, how many cigarettes have you smoked . . .”
6. Question asked as “which describes your use of cocaine (coke, crack or freebase) . . .”

Other changes in question format and wording over time:

- a. In 1990, 1992, 1995, and 1998 question worded as “used alcohol,” in 1999 worded as “have at least one drink,” and in 2000, 2002 and 2004 worded as “drink a glass, bottle, or can.”
- b. The description of chewing tobacco has changed over time; from “smokeless tobacco (chew, plug, snuff)” in 1995 and 1998, to “chewing tobacco or snuff, such as Redman, Levi Garret, Beechnut, Skoal, Skoal Bandits or Copenhagen” in 1999, to “chew tobacco or use snuff” in 2000 and 2002, and then to “chewing tobacco, snuff, or dip” in 2004.
- c. The term “hallucinogens” was used in 1990, 1992, 1995 and 1998 and then changed to “psychedelics” in 2000.
- d. In 1995, 1998, 2000 and 2002 the description of inhalants included only “things you sniff to get high.” In 1999 it included “sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high.”
- e. In 2010, the Ritalin question changed from “Some kids take a medicine prescribed by their doctor to help with hyperactivity or focus (ADD),” to a more inclusive list of drugs, “Some names for this medicine are Ritalin, Adderall, or Concerta.”
- f. In 2014, the description of marijuana was changed from “grass, hash, pot” to “weed, hash, pot”.
- g. In 2014, the question for electronic cigarettes, e-cigs added the language “or vape pens”

Table 4: Current (30-Day) Substance Use by Year, Grade 6

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	–	11.8 ^{1,a}	12.8	12.2 ²	13.8	–	6.6 ^{3,a}	3.8	4.4	4.3	3.5	3.8	2.5	2.1	-0.4
Binge drinking*	–	4.0 ⁴	4.7	6.2	7.6	–	4.7	–	–	–	3.0	3.7	2.4	2.3	-0.1
Cigarettes	–	2.4 ¹	2.8	4.4 ⁵	4.7	–	4.0 ³	2.2	2.0	1.9	1.4	1.7	1.2	1.1	-0.1
Tobacco, chewing	–	–	–	3.6 ^{2,b}	3.5	–	0.8 ^{3,b}	1.0 ^b	1.0	1.2	1.1	1.0	1.0	1.2	0.2
Cigars	–	–	–	–	–	–	1.5 ³	–	–	–	–	–	–	–	–
Tobacco in pipe	–	–	–	–	–	–	0.6 ³	–	–	–	–	–	–	–	–
Bidis	–	–	–	–	–	–	1.0 ³	–	–	–	–	–	–	–	–
Marijuana	–	1.3 ¹	1.3	3.1 ²	3.4	–	1.5 ³	1.3	1.7	1.5	1.2	1.6	1.2	1.3 ^f	0.1
Other illegal drugs** (not alcohol, tobacco or marijuana)	–	–	–	–	–	–	–	–	–	–	–	0.9	0.8	0.6	-0.2
Hallucinogens	–	–	–	–	1.3 ^{2,c}	–	0.6 ^{3,c}	–	–	–	–	–	–	–	–
Inhalants	–	–	–	2.7 ²	3.2	–	1.4 ³	–	–	–	–	–	–	–	–
Cocaine	–	–	–	1.0	1.1	–	–	–	–	–	–	–	–	–	–
Heroin	–	–	–	–	0.6	–	–	–	–	–	–	–	–	–	–
Amphetamines	–	–	–	–	1.4	–	–	–	–	–	–	–	–	–	–
Methamphetamines	–	–	–	–	0.9	–	–	–	–	–	–	–	–	–	–
Party Drugs	–	–	–	–	–	–	0.7	–	–	–	–	–	–	–	–

Notes:

- * Binge drinking in the past two weeks (not in the past 30 days)
- ** Other illegal drugs do not include alcohol, tobacco or marijuana.
- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 60.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 5: Current (30-Day) Substance Use by Year, Grade 8

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	–	29.1 ^{1,a}	24.0	30.1 ²	31.0	–	22.3 ^{3,a}	17.8	18.0	15.4	16.1	14.4	11.9	8.1	-3.8
Binge Drinking*	15.0 ⁴	12.8	10.7	17.1	18.3	–	14.9	10.0	10.2	8.6	9.1	8.1	7.1	4.5	-2.6
Cigarettes	–	12.1 ¹	10.3	18.8 ⁵	15.2	–	12.5 ³	9.2	7.8	6.4	7.3	6.6	5.1	4.0	-1.1
Tobacco, chewing	–	–	–	11.5 ^{2,b}	6.7	–	2.1 ^{3,b}	2.7 ^b	2.8	2.8	3.4	3.0	2.6	1.3	-1.3
Cigars	–	–	–	–	–	–	4.3 ³	8.3	6.4	6.9	8.3	4.3	2.9	1.9	-1.0
Tobacco in a pipe	–	–	–	–	–	–	2.1 ³	5.6	4.0	3.7	5.1	–	–	–	–
Bidis	–	–	–	–	–	–	3.3 ³	6.8	5.3	4.5	6.3	–	–	–	–
Cloves	–	–	–	–	–	–	–	5.0 ³	3.5	3.2	4.0	–	–	–	–
Tobacco in a hookah	–	–	–	–	–	–	–	–	–	–	6.1	–	4.1	4.7	0.6
E-cigarettes	–	–	–	–	–	–	–	–	–	–	–	–	1.7	8.5 ^B	–
Candy flavor tobacco	–	–	–	–	–	–	–	–	–	–	–	5.0	4.1	4.2	0.1
Marijuana	–	7.6 ¹	6.1	16.2 ²	16.5	–	12.0 ³	10.4	9.2	7.0	8.3	9.5	9.4	7.3 ^f	-2.1
Other illegal drugs** (not alcohol, tobacco or marijuana)	–	–	–	–	–	–	–	–	3.3 ³	3.0	3.4	3.0	2.8	1.9	-0.9
Hallucinogens (psychedelics)	–	–	–	–	3.8 ^{2,c}	–	3.1 ^{3,c}	3.0	–	–	–	–	–	–	–
Inhalants	–	–	–	7.3 ²	6.6	–	4.9 ³	5.0	–	5.0	6.4	–	–	–	–
Cocaine	–	3.1 ¹	2.0	3.6 ²	2.5	–	1.5 ³	2.4	3.1	–	–	–	–	–	–
Heroin	–	–	–	–	1.3 ²	–	0.8 ³	–	–	–	–	–	–	–	–
Amphetamines	–	–	–	–	3.9 ²	–	2.7 ³	–	–	–	–	–	–	–	–
Methamphetamines	–	–	–	–	2.3 ²	–	1.2 ³	2.1	1.9	1.3	2.1	–	–	–	–
Party drugs	–	–	–	–	–	–	3.4 ³	–	–	–	–	–	–	–	–
Ecstasy	–	–	–	–	–	–	–	2.4 ³	2.1	–	–	–	–	–	–
Ritalin	–	–	–	–	–	–	–	–	2.8	2.0	2.8	–	1.6	–	–
Pain killers	–	–	–	–	–	–	–	–	–	3.6	4.3	4.3	3.2	2.3	-0.9
Prescription, not prescribed	–	–	–	–	–	–	–	–	–	–	–	–	–	4.2	–

Notes:

- * Binge drinking in the past two weeks (not in the past 30 days)
- ** Other illegal drugs do not include alcohol, tobacco or marijuana.
- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 60.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 6: Current (30-Day) Substance Use by Year, Grade 10

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	–	44.0 ^{1,a}	40.0	37.0 ²	44.9	45.3 ^{3,a}	37.6 ^a	29.3	32.6	32.8	31.7	27.7	23.3	20.6	-2.7
Binge Drinking*	24.5 ⁴	20.2	17.9	22.2	27.7	–	23.2	18.7	18.7	19.6	18.4	16.2	14.3	10.6	-3.7
Cigarettes	–	15.5 ¹	17.1	20.9 ⁵	21.8	25.0 ³	19.8	15.0	13.0	14.9	14.4	12.7	9.5	7.9	-1.6
Tobacco, chewing	–	–	–	15.3 ^{2,b}	9.6	10.5 ^{3,b}	4.6 ^b	4.8 ^b	4.9	6.4	6.7	6.2	4.6	3.7	-0.9
Cigars	–	–	–	–	–	15.4 ³	7.9	11.4	11.4	16.8	16.0	8.5	6.9	5.1	-1.8
Tobacco in a pipe	–	–	–	–	–	–	1.9 ³	5.9	5.6	10.1	7.1	–	–	–	–
Bidis	–	–	–	–	–	–	4.6 ³	8.0	8.1	12.7	10.4	–	–	–	–
Cloves	–	–	–	–	–	–	–	6.3 ³	5.5	9.5	6.7	–	–	–	–
Tobacco in a hookah	–	–	–	–	–	–	–	–	–	–	10.0	–	8.9	10.0	1.1
E-cigarettes	–	–	–	–	–	–	–	–	–	–	–	–	3.9	18.0 ⁶	–
Candy flavor tobacco	–	–	–	–	–	–	–	–	–	–	–	10.6	9.4	9.1	-0.3
Marijuana	–	10.6 ¹	13.2	23.0 ²	26.6	24.3	21.9 ³	18.3	17.1	18.3	19.1	20.0	19.3	18.1 ^f	-1.2
Other illegal drugs** (not alcohol, tobacco or marijuana)	–	–	–	–	–	–	–	–	5.7 ³	7.2	7.0	6.5	5.1	4.4	-0.7
Hallucinogens (psychedelics)	–	–	–	–	5.8 ^{2,c}	–	5.8 ^{3,c}	4.0	–	–	–	–	–	–	–
Inhalants	–	–	–	5.4 ^{2,d}	3.9	5.7 ^d	3.6 ^{3,d}	3.8	–	5.7	5.6	–	–	–	–
Cocaine	–	2.1 ¹	2.1	3.2 ²	3.2	2.6 ⁶	2.6 ³	2.7	–	–	–	–	–	–	–
Heroin	–	–	–	–	1.3 ²	–	1.0 ³	–	–	–	–	–	–	–	–
Amphetamines	–	–	–	–	5.6 ²	–	4.5 ³	–	–	–	–	–	–	–	–
Methamphetamines	–	–	–	–	3.8 ²	–	2.6 ³	2.9	2.9	2.9	3.6	–	–	–	–
Party drugs	–	–	–	–	–	–	6.2 ³	–	–	–	–	–	–	–	–
Ecstasy	–	–	–	–	–	–	–	3.2 ³	2.7	–	–	–	–	–	–
Ritalin	–	–	–	–	–	–	–	–	4.2	5.0	4.9	–	2.8	–	–
Pain killers	–	–	–	–	–	–	–	–	–	10.0	9.5	8.3	6.0	4.6	-1.4
Prescription, not prescribed	–	–	–	–	–	–	–	–	–	–	–	–	–	7.6	–

Notes:

- * Binge drinking in the past two weeks (not in the past 30 days)
- ** Other illegal drugs do not include alcohol, tobacco or marijuana.
- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 60.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 7: Current (30-Day) Substance Use by Year, Grade 12

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	–	52.0 ^{1,a}	51.8	44.8 ²	52.0	49.0 ^{3,a}	46.8 ^a	42.8	42.6	42.1	40.8	40.0	36.1	32.9	-3.2
Binge Drinking*	–	27.8 ⁴	27.3	26.6	32.7	–	31.8	27.3	25.8	26.1	25.9	24.9	21.8	19.2	-2.6
Cigarettes	–	20.7 ¹	22.3	24.0 ⁵	28.6	35.2 ³	27.6	22.7	19.7	20.0	20.0	19.6	15.6	13.0	-2.6
Tobacco, chewing	–	–	–	18.2 ^{2,b}	12.4	11.1 ^{3,b}	8.8 ^b	7.5 ^b	7.6	8.9	8.6	8.9	7.6	5.1	-2.5
Cigars	–	–	–	–	–	21.2 ³	13.1	15.2	18.3	24.3	20.9	17.4	13.7	10.2	-3.5
Tobacco in a pipe	–	–	–	–	–	–	1.7 ³	5.0	5.0	9.1	6.8	–	–	–	–
Bidis	–	–	–	–	–	–	6.5 ³	8.3	8.3	11.8	10.1	–	–	–	–
Cloves	–	–	–	–	–	–	–	5.5 ³	5.5	8.9	7.0	–	–	–	–
Tobacco in a hookah	–	–	–	–	–	–	–	–	–	–	13.1	–	16.7	14.8	-1.9
E-cigarettes	–	–	–	–	–	–	–	–	–	–	–	–	6.7	23.1 ⁸	–
Candy flavor tobacco	–	–	–	–	–	–	–	–	–	–	–	18.8	16.1	15.1	-1
Marijuana	–	15.9 ¹	17.3	23.3 ²	28.7	28.0	24.4 ³	24.7	19.5	21.6	23.4	26.3	26.7 ^f	26.7 ^f	0
Other illegal drugs** (not alcohol, tobacco or marijuana)	–	–	–	–	–	–	–	–	6.8 ³	8.6	8.1	7.5	7.3	6.6	-0.7
Hallucinogens (psychedelics)	–	–	–	–	6.0 ^{2,c}	–	6.5 ^{3,c}	5.1	–	–	–	–	–	–	–
Inhalants	–	–	–	2.7 ^{2,d}	2.3	6.3 ^d	2.4 ^{3,d}	3.0	–	3.5	4.5	–	–	–	–
Cocaine	–	2.6 ¹	2.0	1.9 ²	2.7	2.7 ⁶	2.8 ³	4.4	–	–	–	–	–	–	–
Heroin	–	–	–	–	0.7 ²	–	0.8 ³	–	–	–	–	–	–	–	–
Amphetamines	–	–	–	–	3.6 ²	–	4.0 ³	–	–	–	–	–	–	–	–
Methamphetamines	–	–	–	–	2.9 ²	–	2.9 ³	3.3	2.7	2.7	3.8	–	–	–	–
Party drugs	–	–	–	–	–	–	6.8 ³	–	–	–	–	–	–	–	–
Ecstasy	–	–	–	–	–	–	–	3.6 ³	2.7	–	–	–	–	–	–
Ritalin	–	–	–	–	–	–	–	–	3.6 ³	5.2	5.4	–	4.9	–	–
Pain killers	–	–	–	–	–	–	–	–	–	11.6	12.0	7.9	7.5	5.6	-1.9
Prescription, not prescribed	–	–	–	–	–	–	–	–	–	–	–	–	–	9.0	–

Notes:

- * Binge drinking in the past two weeks (not in the past 30 days)
- ** Other illegal drugs do not include alcohol, tobacco or marijuana.
- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 60.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Lifetime Substance Use

Lifetime prevalence is the percentage of students who had ever tried a substance, *even if on only one occasion*. This section presents lifetime substance use results by grade from 1988 to 2010 (see Tables 8 through 11). Lifetime prevalence trends reflect experimental use, and thus are especially relevant to efforts that aim to delay youths' initiation of substance use.

The prevalence of lifetime use for some substances has been assessed differently as survey instruments have changed over time. Superscripts in the tables describe any changes to survey questions or responses. Therefore, readers should use caution when making strong conclusions about changes over time for these substances.

Description of Superscript Notes for Tables 8 through 11

How the question was asked and changes over time:

1. Question asked as "how often did you use . . ."
2. Question asked as "have you ever in your life, even once, used . . ."
3. Question asked as "how old were you when you first used . . ."
4. Question asked as "how many times have you . . ."

Other changes in question format and wording over time:

- a. In 1988 and 1990 three questions were combined to create an alcohol estimate (how often did you use: beer, wine or wine coolers, hard liquor). In 1992, four questions were combined (beer, wine, wine coolers, hard liquor). In 1995 only one question was asked about alcohol (beer, wine, wine coolers, liquor). In 2000 the language changed to specify "more than a sip or two."
- b. The description of chewing tobacco has changed from "chewing tobacco" in 1988 to "smokeless tobacco (chew, plug, snuff)" in 1990. In 1995, "spit" was added, then changed to "(chew, dip or snuff)" in 2000, and to "chewing tobacco, snuff, or dip" in 2002.
- c. The term "hallucinogens" was used in 1990, 1992, 1995 and 1998 and then changed to "psychedelics" in 2000.
- d. In 1988 the inhaled substance question included "glue, gasoline, paint thinner, spray cans, and white out." In 1990, "snappers, poppers, and rush" were added. In 2002 the question was simplified to say only "things you sniff to get high."
- e. In 1990 and 1992 the over-the-counter question included "drugs purchased from the drug store to get high (diet pills like Dexatrim, stay awake pills like NoDoz and Vivarin, pep pills, Nyquil or other coffee medicine)." In 1995 it was shortened to "drugs you can get from the drug store to get high."
- f. In 1999, 2002, and 2004 "without a doctor's prescription" was added to the steroids question.
- g. In 1990 the methamphetamine question was for "crystal methamphetamine (crystal meth, ice)." In 1998 and 2000 the question was "methamphetamine, specifically (meth, crystal meth, ice, crank)." In 2002 and 2004 a statement was added, "do not include other types of amphetamines."
- h. In 2010, the description "(coke, rock, snow)" was dropped from the cocaine question.

Table 8: Lifetime Substance Use by Year, Grade 6

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	51.4 ^{1,a}	33.0	33.0 ^a	33.4 ^{2,a}	39.8	–	21.2 ^a	32.7	30.3	30.9	29.2	26.3	23.0	21.2	-1.8
Cigarette (even just a puff)	–	–	–	26.7 ³	26.5	–	15.1	–	–	–	–	–	–	–	–
Cigarette (whole)	–	–	–	–	–	–	7.2 ³	6.2	5.4	4.9	3.8	–	–	–	–
Tobacco, chewing	9.5 ^{1,b}	5.4 ^b	5.5	7.1 ^{2,b}	7.8	–	1.8 ^{3,b}	^b –	–	–	–	–	–	–	–
Marijuana	3.6 ¹	1.7	1.9	4.9 ²	7.0	–	2.2 ³	3.4 ²	3.0	3.2	2.7	3.9	2.9	3.1	0.2
Hallucinogens	1.5 ^{1,c}	0.8	1.2	1.1 ²	2.6	–	0.8 ^c	–	–	–	–	–	–	–	–
Inhalants	13.0 ^{1,d}	7.5 ^d	7.7	3.9 ²	7.0	–	2.5	3.6 ^d	3.7	3.7	2.9	3.5	2.4	2.3	-0.1
Over-the-counter	–	7.0 ^{1,e}	7.8	2.0 ^{2,e}	–	–	–	–	–	–	–	–	–	–	–
Cocaine	0.8 ¹	0.9	1.1	1.3 ²	2.3	–	–	–	–	–	–	–	–	–	–
Steroids	1.7 ¹	1.2	1.1	1.2 ²	2.6	–	–	–	–	–	–	–	–	–	–
Other illegal drugs	–	–	–	–	–	–	–	3.3 ²	2.9	3.3	3.8	3.3	2.0	2.4	0.4
Heroin	–	–	–	–	1.7 ²	–	–	–	–	–	–	–	–	–	–
Illegal injection drugs	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Amphetamines	–	–	–	–	3.4 ²	–	–	–	–	–	–	–	–	–	–
Methamphetamines	–	0.9 ^{1,g}	–	–	2.3 ^{2,g}	–	–	–	–	–	–	–	–	–	–
Party drugs	–	–	–	–	–	–	0.9 ²	–	–	–	–	–	–	–	–

Notes:

- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 65.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 9: Lifetime Substance Use by Year, Grade 8

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	68.9 ^{1,a}	60.2	55.3 ^a	61.4 ^{2,a}	62.7	–	45.7 ^{3,a}	44.2	42.0	37.6	39.4	39.0	35.4	29.0	-6.4
Cigarette (even just a puff)	–	–	–	53.3 ³	49.1	–	37.1	28.6	23.9	19.8	20.1	17.6	14.7	11.8	-2.9
Cigarette (whole)	–	–	–	–	–	–	25.3 ³	19.8	15.8	12.7	13.2	–	10.2	–	–
Tobacco, chewing	16.6 ^{1,b}	13.9 ^b	13.1	22.9 ^{2,b}	14.8	–	5.2 ^{3,b}	8.0 ^b	7.3	–	–	–	–	–	–
Marijuana	14.4 ¹	11.2	9.7	27.2 ²	28.2	–	19.7	15.7	14.0	10.7	11.9	13.2	13.7	10.4	-3.3
Hallucinogens (psychedelics)	4.1 ^{1,c}	5.7	5.6	9.3 ²	8.7	–	4.7 ^c	–	–	–	–	–	–	–	–
Inhalants	17.3 ^{1,d}	17.1 ^d	17.4	14.5 ^{2,d}	14.3	–	9.6	–	5.3	5.7	6.1	5.8	6.1	4.5	-1.6
Over-the-counter	–	23.2 ^{1,e}	18.4	12.3 ^{2,e}	–	–	–	–	–	–	–	–	–	–	–
Cocaine	2.8 ¹	3.4	2.6	5.5 ²	5.2	–	3.3 ²	3.0	3.4	2.4 ⁴	3.2	2.6 ^h	3.8	2.9	-0.9
Steroids	3.3 ^{1,f}	2.7	1.9	2.5 ²	2.6	–	2.2 ^f	3.1	1.6	1.9	–	2.4	3.0	2.3	-0.7
Other illegal drugs	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Heroin	–	–	–	–	2.6 ²	–	1.4	–	–	1.6	–	2.2	3.0	2.6	-0.4
Illegal injection drugs	–	–	–	–	–	–	1.0 ²	1.6	1.4	1.7	–	–	–	–	–
Amphetamines	–	–	–	–	8.4 ²	–	4.3	–	–	–	–	–	–	–	–
Methamphetamines	–	3.0 ^{1,g}	–	–	4.6 ^{2,g}	–	2.0	2.5	3.3	1.9	2.8	2.4	3.3	2.5	-0.8
Party drugs	–	–	–	–	–	–	4.8 ²	–	–	–	–	–	–	–	–

Notes:

- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 65.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 10: Lifetime Substance Use by Year, Grade 10

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	84.1 ^{1,a}	75.7	70.3 ^a	73.0 ^{2,a}	79.7	68.9 ^{3,a}	65.0	60.0	60.4	61.2	60.6	57.1	52.2	50.1	-2.1
Cigarette (even just a puff)	–	–	–	59.8 ³	64.1	–	52.2	38.9	35.1	35.5	33.0	29.2	23.9	22.0	-1.9
Cigarette (whole)	–	–	–	–	–	50.1 ³	40.9	29.6	26.3	26.6	25.2	–	18.5	–	–
Tobacco, chewing	21.5 ^{1,b}	22.1 ^b	23.2	30.7 ^{2,b}	25.8	–	14.3 ^{3,b}	13.1 ^b	11.6	–	–	–	–	–	–
Marijuana	32.7 ¹	21.5	22.8	39.1 ²	49.5	42.4	37.6	32.4	29.5	30.8	30.8	30.9	29.3	29.4	0.1
Hallucinogens (psychedelics)	12.1 ^{1,c}	9.1	11.1	15.4 ²	18.8	–	10.7 ^c	–	–	–	–	–	–	–	–
Inhalants	19.5 ^{1,d}	17.7 ^d	15.6	12.3 ^{2,d}	15.3	–	11.9	–	6.6	10.7	8.9	9.2	9.2	7.6	-1.6
Over-the-counter	–	27.2 ^{1,e}	22.3	10.4 ^{2,e}	–	–	–	–	–	–	–	–	–	–	–
Cocaine	8.1 ¹	4.3	3.5	7.4 ²	9.4	7.7 ⁴	6.0 ²	5.4	6.0	7.3 ⁵	7.0	6.1 ^h	6.1	4.2	-1.9
Steroids	4.9 ^{1,f}	3.0	2.2	2.1 ²	3.1	3.6 ^{4,f}	2.9 ²	2.9 ⁴	2.7	3.2	–	3.5	4.2	3.2	-1.0
Other illegal drugs	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Heroin	–	–	–	–	3.9 ²	6.3 ⁴	1.9 ²	–	–	4.7	–	3.5	4.2	3.4	-0.8
Illegal injection drugs	–	–	–	–	–	2.8 ⁴	1.3 ¹	2.1	1.8	2.5	–	–	–	–	–
Amphetamines	–	–	–	–	14.6 ²	–	8.4	–	–	–	–	–	–	–	–
Methamphetamines	–	3.1 ^{1,g}	–	–	9.8 ^{2,g}	–	5.3	4.5 ^g	5.1	5.9	4.7	4.8	5.2	4.1	-1.1
Party drugs	–	–	–	–	–	–	9.3 ²	–	–	–	–	–	–	–	–

Notes:

- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 65.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Table 11: Lifetime Substance Use by Year, Grade 12

Substance	Percent of Students														Change
	1988	1990	1992	1995	1998	1999	2000	2002	2004	2006	2008	2010	2012	2014	
Alcohol	–	83.0 ^{1,a}	79.8 ^a	81.9 ^{2,a}	84.2	75.9 ^{3,a}	76.0	74.9	72.6	72.2	72.4	70.6	68.0	66.2	-1.8
Cigarette (even just a puff)	–	–	–	67.6 ³	67.4	–	60.9	52.1	47.5	45.0	44.3	40.8	36.6	31.5	-5.1
Cigarette (whole)	–	–	–	–	–	59.6 ³	52.0	42.5	36.8	35.5	34.3	–	28.7	–	–
Tobacco, chewing	–	28.5 ^{1,b}	27.9	37.7 ^{2,b}	35.0	–	24.8 ^{3,b}	20.0 ^b	17.6	–	–	–	–	–	–
Marijuana	–	34.0 ¹	32.9	43.5 ²	55.1	57.3	50.5	48.0	41.1	43.1	44.6	45.7	45.6	45.7	0.1
Hallucinogens (psychedelics)	–	13.7 ^{1,c}	16.8	18.7 ²	23.8	–	15.1 ^c	–	–	–	–	–	–	–	–
Inhalants	–	16.4 ^{1,d}	13.1	11.0 ^{2,d}	13.3	–	13.1	–	7.1	9.4	9.7	10.7	9.7	8.0	-1.7
Over-the-counter	–	27.2 ^{1,e}	22.3	10.4 ^{2,e}	–	–	–	–	–	–	–	–	–	–	–
Cocaine	–	7.8 ¹	4.6	7.6 ²	9.7	13.1 ⁴	9.2 ²	8.2	8.3	9.8 ⁵	10.5	8.9 ^h	8.1	6.5	-1.6
Steroids	–	3.2 ^{1,f}	2.4	2.4 ²	3.0	2.6 ^{4,f}	2.9 ²	4.2 ⁴	2.5	3.9	–	3.5	4.5	3.2	-1.3
Other illegal drugs	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Heroin	–	–	–	–	3.6 ²	4.6 ⁴	2.4 ²	–	–	4.7	–	4.1	5.1	3.2	-1.9
Illegal injection drugs	–	–	–	–	–	3.0 ⁴	1.5 ¹	2.1	1.8	2.9	–	–	–	–	–
Amphetamines	–	–	–	–	14.9 ²	–	10.0	–	–	–	–	–	–	–	–
Methamphetamines	–	4.3 ^{1,g}	–	–	11.0 ^{2,g}	–	7.5	7.2 ^g	6.3	7.1	5.6	4.8	5.6	3.8	-1.8
Party drugs	–	–	–	–	–	–	13.5 ²	–	–	–	–	–	–	–	–

Notes:

- Dashes (–) indicate a substance was not represented on that particular year’s survey.
- Change column provides the percentage point change from 2012 to 2014. Changes that are statistically significant at the 95 percent confidence level are bolded.
- The superscript numbers and letters are used to describe the changes in questions over time. Details are available on page 65.

Source: 1988 and 1990 SADUS, WSSAHB 1992 and 1995, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Alcohol Use

Alcohol has been consistently reported as the substance most frequently used by Washington’s youth. However, the prevalence of past 30-day use of alcohol has steadily declined nationally (Monitoring the Future, 2015) and in Washington State since 2000. As age-specific survey data illustrate, the number of youth using alcohol increases sharply with each grade. The number of Grade 6 and 8 students who report any lifetime use is of particular concern because of the strong association between age of initiation and subsequent alcohol abuse and dependence.

Lifetime Alcohol Use

In 2014, 21 percent of Grade 6 students, 29 percent of Grade 8 students, 50 percent of Grade 10 students, and 66 percent of Grade 12 students reported having tried more than a sip or two of alcohol sometime in their lives (lifetime use).

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to report they drank more than a sip or two of alcohol in their lifetime.

Differences by gender:

- Grade 6 males were more likely than females to report they drank more than a sip or two of alcohol in their lifetime.
- Grade 10 females were more likely than males to report they drank more than a sip or two of alcohol in their lifetime.

Differences over time:

- Among Grade 8 students, there was a decrease in lifetime alcohol use from 2012 to 2014.
- Among Grade 6 and 10 students, there were decreases from 2008 to 2014 in lifetime alcohol use. Among Grade 8 and 12 students, there were decreases in lifetime alcohol use from 2002 through 2014.



Survey Questions:

- How old were you the first time you: Had more than a sip or two of beer, wine, or hard liquor (for example: vodka, whiskey, or gin)?
- Have you ever, even once in your lifetime: Had more than a sip or two of beer, wine, or hard liquor (for example: vodka, whiskey, or gin)?

Note: Percentage represents students who had ever had more than a sip of alcohol at any age in their life (Grades 8, 10 and 12) or had ever had a sip of alcohol in their life (Grade 6).

Source: SADUS 1988 and 1990, WSSAHB 1992, 1995, 1998, and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014

30-Day Alcohol Use

In 2014, 2 percent of Grade 6 students, 8 percent of Grade 8 students, 21 percent of Grade 10 students, and 33 percent of Grade 12 students reported drinking alcohol in the past 30 days.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to use alcohol in the past 30 days.

Differences by gender:

- Grade 6 males were more likely than females to use alcohol in the past 30 days.
- Grade 10 females were more likely than males to use alcohol in the past 30 days.

Differences over time:

- Among Grade 8 and 10 students, there were decreases in 30-day alcohol use from 2012 to 2014.
- Among Grade 6 and 8 students, there were decreases in 30-day alcohol use from 2002 through 2014. Among Grade 10 students, there was a decrease in 30-day alcohol use from 2008 to 2014. Among Grade 12 students there was a decrease in 30-day alcohol use from 2010 to 2014.



Survey Question: During the past 30 days, on how many days did you: Drink a glass, can or bottle of alcohol (beer, wine, wine coolers, hard liquor)?

Note: Percentages represent students who reported that they drank alcohol on any days in the past 30 days.

Source: SADUS 1990, WSSAHB 1992, 1995, 1998, and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Binge Drinking

The survey question on binge drinking (framed as five or more drinks in a row during the previous two weeks) may underestimate excessive alcohol consumption. Low-weight and inexperienced drinkers suffer negative effects from fewer than five drinks, and students may underestimate the amount of alcohol they consume in a “drink.” In addition, the new recommended measurement of binge drinking for women is 4 drinks or more in one occasion (Chavez, 2011).

In 2014, 2 percent of Grade 6 students, 5 percent of Grade 8 students, 11 percent of Grade 10 students, and 19 percent of Grade 12 students reported binge drinking in the past two weeks.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to report binge drinking.

Differences by gender:

- Grade 12 males were more likely than females to report binge drinking.

Differences over time:

- Among Grade 8 and 10 students, there were decreases in binge drinking from 2012 to 2014.
- Among Grade 8 students, there was a decrease in binge drinking from 2002 through 2014. Among Grade 12 students, there was a decrease from 2010 to 2014 in binge drinking.



Survey Question: Think back over the last 2 weeks. How many times have you had five or more drinks in a row? (A drink is a glass of wine, a bottle of beer, a shot glass of liquor, or a mixed drink.)

Note: Percentages represent students who reported that they had five or more drinks in a row in the past two weeks.

Source: SADUS 1988 and 1990, WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Average Age of First Alcohol Use

Some youth begin experimenting with alcohol and other drugs at an early age. Early (age 12-14) and late (age 15-17) adolescence initiation and use of alcohol are associated with alcohol related problems in adulthood (Buchman, 2009; McCambridge, 2011); the younger the age of drinking onset, the greater the chance that an individual will develop a clinically defined alcohol disorder at some point in life (National Center on Addiction and Substance Abuse, 2011).

The following summarizes the average age when students first tried more than a sip or two of alcohol and the average age when students began drinking regularly in 2014:

- Grade 10 students, on average, first had more than a sip or two of beer, wine, or hard liquor at 12.8 years of age.
- Grade 10 students, on average, began drinking alcoholic beverages at least once or twice a month at 13.8 years of age.
- These results are similar to the results from previous years.

Table 12
Average Age of First Use and Regular Use of Alcohol in 2014

Behavior	Mean Age of First Reported Use		
	Grade 8	Grade 10	Grade 12
Had more than a sip of beer, wine, or hard liquor	11.3 (± 0.04)	12.8 (± 0.06)	14.1 (± 0.1)
Began drinking regularly, at least once or twice a month	12.1 (± 0.2)	13.8 (± 0.2)	15.3 (± 0.2)

Survey Questions:

- How old were you the first time you had more than a sip or two of beer, wine, or hard liquor (for example: vodka, whiskey, or gin)?
- How old were you the first time you began drinking alcoholic beverages regularly, that is, at least once or twice a month?

Notes:

- Age of first use is calculated by excluding students who responded “Never had,” drank alcohol and calculating the mean age of use among those who drank at any age.
- Age of first use is calculated by excluding students who responded “Never had,” drank alcohol regularly and calculating the mean age of use among those who drank regularly at any age.

Source: HYS 2014.

Levels of Problem Drinking: Composite Scale

The level of drinking is an important consideration in the design of prevention and intervention strategies. The definitions of experimental, problem and heavy drinking combine frequency of drinking with episodes of binge drinking (see Notes below) (Courtney et al., 2009). Students reported the following levels of drinking in 2014:

- Experimental drinking: 4 percent of Grade 8, 9 percent of Grade 10, and 12 percent of Grade 12 students.
- Problem drinking: 3 percent of Grade 8, 6 percent of Grade 10, and 11 percent of Grade 12 students.
- Heavy drinking: 3 percent of Grade 8, 7 percent of Grade 10, and 12 percent of Grade 12 students.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to report experimental drinking, problem drinking and heavy drinking.

Differences by gender:

- Grade 10 and 12 females were more likely than males to report experimental drinking.
- Grade 10 and 12 females were more likely than males to report problem drinking.
- Grade 10 and 12 males were more likely than females to report heavy drinking.

Differences over time:

- Among 8 students, there was a decrease in experimental drinking from 2012 to 2014. Among 8 and 10 students, there were decreases in problem drinking from 2012 to 2014.
- Among Grade 8, 10 and 12 students, there were decreases in experimental drinking, problem drinking and heavy drinking from 2010 to 2014. Among Grade 10 students, there was a decrease in problem drinking from 2006 through 2014. Among Grade 10 and 12 students, there were decreases in heavy drinking from 2005 through 2014.



Survey Questions:

- During the past 30 days, on how many days did you: Drink a glass, can or bottle of alcohol?
- Think back over the last 2 weeks. How many times have you had five or more drinks in a row?

Notes:

- Experimental drinking represents drinking 1–2 times in the past 30 days and no binge drinking in the past two weeks.
- Problem drinking represents drinking 3–5 times in the past 30 days and/or binge drinking in the past two weeks.
- Heavy drinking represents drinking 6 or more times in the past 30 days and/or binge drinking 2 or more times in the past two weeks.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Perception of Access to Alcohol

In spite of the laws that seek to prevent underage drinking, a high percentage of youth do not find it hard to obtain alcohol. The perception of easy access to alcohol is lower among Washington State youth than the national average (Johnston, 2015).

In 2014, 73 percent of Grade 6 students, 48 percent of Grade 8 students, 22 percent of Grade 10 students, and 14 percent of Grade 12 students reported that alcohol would be “very hard” to get.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was less likely to perceive that alcohol would be very hard to get.

Differences by gender:

- Grade 6 males were less likely than females to perceive that alcohol would be very hard to get.
- Grade 10 females were less likely than males to perceive that alcohol would be very hard to get.

Differences over time:

- Among Grade 10 students, there was an increase in the perception that alcohol would be very hard to get from 2012 to 2014.
- Among Grade 6 students, there was an increase from 2008 to 2014 in the perception that alcohol would be very hard to get. Among Grade 8, 10, and 12 students, there were increases in the perception that alcohol would be very hard to get from 2002 through 2014.



Survey Question: If you wanted to get some beer, wine, or hard liquor (for example: vodka, whiskey, or gin), how easy would it be for you to get some?

Note: Percentages represent students who reported it would be very hard to get alcohol if they wanted some.

Source: WSSAHB 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Usual Sources of Alcohol

Younger students are more likely to get alcohol from home, while older students are more likely to get alcohol from friends, at parties, and to give money to someone to buy it for them. The following chart represents where they usually obtained alcohol, among students who used alcohol in the past 30 days.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to give someone money to get alcohol.
- Grade 8 and 10 students were more likely than Grade 12 students to get alcohol at home without parental permission.
- Grade 10 and 12 students were more likely than Grade 8 students to get alcohol from friends, at a party, and less likely to get alcohol from some other way.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to steal alcohol from a store.
- Grade 8, 10 and 12 females were more likely than males to get alcohol from friends
- Grade 10 and 12 males were more likely than females to buy alcohol from a store.
- Grade 10 males were more likely than females to get alcohol from some other way, and Grade 10 females were more likely than males to get alcohol and from home without parental permission.
- Grade 8 males were more likely than females to get alcohol and from home with parental permission.

Differences over time:

- Among Grade 8 students, there was a decrease in getting alcohol at a party, and a significant increase in getting alcohol from some other way from 2012 to 2014.
- Among Grade 10 students, there was an increase in getting alcohol from friends from 2012 to 2014.



Survey Question: During the past 30 days, how did you usually get alcohol (beer, wine, or hard liquor)? Choose all that apply.

Notes: Students could check multiple responses. Stealing alcohol from a store was not asked in 2012.

- Students who reported “did not get alcohol in the past 30 days” were not included in the results.
- The sample sizes for the 2014 results in this figure are: 382 Grade 8; 755 Grade 10; and 921 Grade 12 students.

Source: HYS 2014.

Perception of Risk from Daily Alcohol Consumption

Because alcohol use is so widely accepted in our culture, it is not surprising that youth do not appreciate the possible harmful effects of alcohol consumption.

In 2014, 33 percent of Grade 6 students, 43 percent of Grade 8, 44 percent of Grade 10 students and 38 percent of Grade 12 students perceived “great risk” in having one or two drinks of alcohol every day.

Differences by grade level:


- Grade 6 students were less likely than students in Grades 8, 10 and 12 to perceive great risk in having one or two drinks of alcohol every day.
- Grade 8 and 12 were less likely than students in Grade 10 to perceive great risk in having one or two drinks of alcohol every day.

Differences by gender:

- Grade 8, 10 and 12 males were less likely than males to perceive great risk in having more than one or two drinks of alcohol every day.

Differences over time:

- Among Grade 6 and 8 students, there were increases in the perception of great risk in having one or two drinks of alcohol every day from 2012 to 2014.
- Among Grade 8 and 10 students there were increases in the perception of great risk in having one or two drinks of alcohol every day from 2002 through 2014. Among Grade 12 students, there was an increase from 2002 to 2010 in the perception of great risk in having one or two drinks of alcohol every day.



Survey Question: How much do you think people risk harming themselves if they take one or two drinks of an alcoholic beverage (wine, beer, a shot of liquor) nearly every day?

Note: Percentages represent students who reported that there is great risk from daily alcohol consumption.

Source: WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Tobacco Use

Historically, cigarettes have been the most popular tobacco product used by youth. Youth cigarette smoking rates peaked in the late 1990s but have dropped significantly since. Recently, youth have been experimenting with other types of tobacco including smokeless, hookah, and cigars. Use of e-cigarettes or vapor products is also common among youth in our state and nationally.

Lifetime Cigarette Smoking

In 2014, 12 percent of Grade 8 students, 22 percent of Grade 10 students, and 31 percent of Grade 12 students reported ever having smoked a cigarette, even just a puff.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to have ever smoked a cigarette, even just a puff.

Differences by gender:

- Grade 12 males were more likely than females to have ever smoked a cigarette, even just a puff.

Differences over time:

- Among Grade 8 and 12 students, there were decreases in ever smoking cigarettes from 2012 to 2014.
- Among Grade 8, 10 and 12 students there were decreases in ever smoking a cigarette from 2002 through 2014.



Survey Question: How old were you the first time you smoked a cigarette, even just a puff?

Note: Lifetime percentage represents students who had ever smoked a whole cigarette at any age in their life.

Source: WSSAHB 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

30-Day Cigarette Smoking

In 2014, 1 percent of Grade 6 students, 4 percent of Grade 8 students, 8 percent of Grade 10 students, and 13 percent of Grade 12 students reported smoking a cigarette in the past 30 days.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to have smoked cigarettes in the past 30 days.

Differences by gender:

- Grade 6 and 12 males were more likely than females to have smoked cigarettes in the past 30 days.

Differences over time:

- Among Grade 8 and 12 students, there were decreases in 30-day cigarette smoking from 2012 to 2014.
- Among Grade 6, 8, 10 and 12 students, there were decreases in 30-day use in cigarette smoking from 2002 through 2014.



Survey Question: During the past 30 days, on how many days did you: Smoke cigarettes?

Note: Percentages represent students who smoked cigarettes on any days in the past 30 days.

Source: SADUS 1990, WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010 and 2012.

Average Age of First Cigarette Smoking

Table 13 shows the average age of first use for students who had ever tried a cigarette, even just a puff.

The earlier youth begin smoking cigarettes, the more likely they are to become strongly addicted to nicotine. About nine out of ten adult smokers began smoking when they were teens or earlier (Surgeon General Report 2012).

Grade 10 students, on average, first smoked a puff of a cigarette at 12.7 years of age. These results are similar to those from previous Healthy Youth Survey administrations.

Table 13
Average Age of First Cigarette Use in 2014

Behavior	Mean Age of First Reported Use		
	Grade 8	Grade 10	Grade 12
Smoked a cigarette, even just a puff	11.5 (\pm 0.1)	12.7 (\pm 0.2)	13.9 (\pm 0.2)

Survey Question: How old were you the first time you smoked a cigarette, even just a puff?

Note: Age of first use is calculated by excluding students who responded "Never had," smoked a puff of a cigarette and calculating the mean age of use among those who smoked at any age.

Source: HYS 2014.

30-Day Chewing Tobacco Use

Using chewing tobacco represents a significant health risk and is not a safe substitute for smoking cigarettes. Chewing tobacco causes cancers of the mouth, pharynx and esophagus; gum recession; and an increased risk for heart disease and stroke. Youth chewing tobacco use can lead to a lifetime of addiction to nicotine, and frequently leads to cigarette smoking (U.S. Department of Health and Human Services, 1994; National Cancer Institute, 1992; World Health Organization, 2007; and Tomar, 2003).

In 2014, use of chewing tobacco in the past 30 days was reported by 1 percent of Grade 6 students and Grade 8 students, 4 percent of Grade 10 students, and 5 percent of Grade 12 students.

Differences by grade level:


- Grade 6 and Grade 8 students were less likely than Grade 10 and 12 students to use chewing tobacco in the past 30 days.
- Grade 10 students were less likely than Grade 12 students to use chewing tobacco in the past 30 days.

Differences by gender:

- Grade 6, 8, 10 and 12 males were more likely than females to use chewing tobacco in the past 30 days.

Differences over time:

- Among Grade 8 and 12 students, there were decreases in 30-day chewing tobacco use from 2012 to 2014.
- There were no significant trends in 30-day chewing tobacco use for any grades from 2002 through 2014.



Survey Question: During the past 30 days, on how many days did you: Use chewing tobacco, snuff, or dip?

Note: Percentages represent students who reported that they had used chewing tobacco on any days in the past 30 days.

Source: WSSAHB 1995, 1998 and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

30-Day Cigar, Cigarillo or Little Cigar Smoking

In 2014, cigar smoking in the past 30 days was reported by 2 percent of Grade 8 students, 5 percent of Grade 10 students, and 10 percent of Grade 12 students.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to have smoked cigars in the past 30 days

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to have smoked cigars in the past 30 days.

Differences over time:

- Among Grade 8, 10, and 12 students, there were decreases in 30-day cigar smoking from 2012 to 2014.
- There were no significant trends in 30-day cigar smoking for any grades from 2002 through 2014.



Survey Question: During the past 30 days, on how many days did you: Smoke cigars, cigarillos or little cigars?

Note: Percentages represent students who reported that they had smoked cigars on any days in the past 30 days.

Source: YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

30-Day Electronic Cigarettes, E-cigs or Vape Pen Use


In 2014, e-cig or vape pen use in the past 30 days was reported by 8 percent of Grade 8 students, 18 percent of Grade 10 students, and 23 percent of Grade 12 students.

Differences by grade level:

- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to have used an e-cig or vape pen in the past 30 days

Differences by gender:

- Grade 10 and 12 males were more likely than females to have used an e-cig or vape pen in the past 30 days.



Survey Question: During the past 30 days, on how many days did you: Use an electronic cigarette, also called e-cigs, or vape pens?

Notes:

- Percentages represent students who reported that they had used an electronic cigarette, also called e-cigs, or vape pens on any days in the past 30 days.
- A question about 30-day electronic cigarettes and e-cigs was asked in 2012, but the question did not include vape pens.

Source: HYS 2014.

Secondhand Smoke Exposure

Secondhand smoke exposure causes disease and premature death in children and adults who do not smoke. Scientific evidence indicates that there is no risk-free level of exposure to secondhand smoke (U.S. Department of Health and Human Services, 2006c).

In 2014, 19 percent of Grade 6 students, 22 percent of Grade 8 students, 28 percent of Grade 10 students, and 30 percent of Grade 12 students reported being exposed to secondhand smoke in a room in the past week.

Differences by grade level:


- Grade 6 students were less likely than Grade 8, 10 and 12 students to be exposed to secondhand smoke in a room in the past week.
- Grade 8 students were less likely than Grade 10 and 12 students to be exposed to secondhand smoke in a room in the past week.

Differences by gender:

- Grade 6 males were more likely than females to be exposed to secondhand smoke in a room in the past week.

Differences over time:

- Among Grade 8, 10 and 12 students, there were decreases in exposure to secondhand smoke in a room in the past week from 2012 to 2014.
- There were no significant trends in exposure to secondhand smoke in a room in the past week for any grades from 2002 through 2014.



Survey Question: During the past 7 days, on how many days were you in the same room with someone who was smoking cigarettes?

Note: Percentages represent students who reported they had been exposed to secondhand smoke in a room in the past week.

Source: WSSAHB 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Perception of Access to Cigarettes

There is strong evidence that community mobilization, along with additional interventions such as strong local laws for tobacco retailers, active enforcement of retailer sales laws, and retailer education with reinforcement are effective in reducing youth tobacco use and access to tobacco products from commercial sources (Task Force on Community Preventive Services, 2005).

In 2014, 77 percent of Grade 6 students, 60 percent of Grade 8 students, 34 percent of Grade 10 students, and 18 percent of Grade 12 students reported that it would be very hard to get cigarettes.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was less likely to perceive that cigarettes would be very hard to get.

Differences by gender:

- Grade 8 males were more likely than females to perceive that cigarettes are very hard to get.
- Grade 12 females were more likely than males to perceive that cigarettes are very hard to get.

Differences over time:

- Among Grade 8, 10 and 12 students, there were increases in the perception that it would be very hard to get cigarettes from 2012 to 2014.
- Among Grade 6, 8, 10 and 12 students, there were increases in the perception that it would be very hard to get cigarettes from 2002 through 2014.



Survey Question: If you wanted to get some cigarettes, how easy would it be for you to get some?

Note: Percentages represent students who reported it would be very hard to get cigarettes if they wanted some.

Source: WSSAHB 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Perception of Risk from Heavy Cigarette Smoking (Pack or More Daily)

In 2014, 62 percent of Grade 6 students, 73 percent of Grade 8 students, 78 percent of Grade 10 students, and 75 percent of Grade 12 students reported there was great risk in smoking a pack or more of cigarettes a day.

Differences by grade level:


- Grade 6 students were less likely than Grade 8, 10, and 12 students to perceive great risk in smoking a pack or more of cigarettes a day.
- Grade 8 students were less likely than Grade 10 and 12 students to perceive great risk in smoking a pack or more of cigarettes a day.

Differences by gender:

- Grade 6 and 10 males were less likely than females to perceive great risk in smoking a pack or more of cigarettes a day.

Differences over time:

- Among Grade 12, there was an increase in the perception of great risk from smoking a pack of cigarettes or more a day from 2012 to 2014.
- Among Grade 6 students, there was a decrease in the perception of great risk from smoking a pack of cigarettes or more a day from 2002 through 2014. Among Grade 10 students, there was an increase from 2006 to 2014 in the perception of great risk from smoking a pack of cigarettes or more a day.



Survey Question: How much do you think people risk harming themselves if they: Smoke one or more packs of cigarettes per day?

Note: Percentages represent students who reported there is great risk from smoking a pack or more of cigarettes a day.

Source: WSSAHB 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Marijuana Use

Marijuana has been the most widely used drug since the state’s first survey of youth substance use in 1988. It is also by far the primary drug used by youth entering treatment. National trends in use have been associated with youth perception of the risk of marijuana use—that is, as perception of risk declined during the 1990s, the prevalence of marijuana use grew.

Lifetime Marijuana Use

In 2014, 3 percent of Grade 6 students, 10 percent of Grade 8 students, 29 percent of Grade 10 students, and 46 percent of Grade 12 students reported having used marijuana at some time in their life.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to have used marijuana in their lifetime.

Differences by gender:

- There were no differences in lifetime marijuana use by gender.

Differences over time:

- Among Grade 8 students, there was a decrease in lifetime marijuana use from 2012 to 2014.
- There were no significant trends in lifetime marijuana use for any grades from 2002 through 2014.



Survey Question: How old were you the first time you smoked marijuana?

- How old were you the first time you: Used marijuana?
- Have you ever, even once in your lifetime: Used marijuana?

Notes:

- Percentages represent students who had ever used marijuana at any age in their life (Grades 8, 10 and 12) or had ever used marijuana in their life (Grade 6).
- For both questions the word “smoked” was changed to “used” in 2014.

Source: SADUS 1988 and 1990, WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

30-Day Marijuana Use

In 2014, 1 percent of Grade 6 students, 7 percent of Grade 8 students, 18 percent of Grade 10 students, and 27 percent of Grade 12 students reported using marijuana in the past 30 days.

Differences by grade level:


- Among Grade 6, 8, 10, and 12 students, as grade levels increase, each grade was more likely to have used marijuana in the past 30 days.

Differences by gender:

- Grade 6 and 12 males were more likely than females to have used marijuana in the past 30 days.

Differences over time:

- Among Grade 8 students there was a decrease in 30-day marijuana use from 2012 to 2014.
- There were no significant trends in 30-day marijuana use for any grades from 2002 through 2014.



Survey Question: During the past 30 days, on how many days did you: Use marijuana or hashish (weed, hash, pot)?

Notes:

- Percentages represent students who used marijuana on any days in the past 30 days.
- The description of marijuana changed from “grass, hash, pot” to “weed, hash, pot” in 2014.

Source: SADUS 1990, WSSAHB 1992, 1995, 1998 and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Levels of Marijuana Use

Students reported the following levels of marijuana use in 2014:

- Experimental use: about 3 percent of Grade 8, 7 percent of Grade 10, and 9 percent of Grade 12 students.
- Occasional use: about 2 percent of Grade 8, 4 percent of Grade 10, and 5 percent of Grade 12 students.
- Regular use: about 3 percent of Grade 8, 8 percent of Grade 10, and 13 percent of Grade 12 students.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to report experimental, occasional, and regular marijuana use.

Differences by gender:

- Grade 10 females were more likely than males to report occasional marijuana use.
- Grade 8, 10 and 12 males were more likely than females to report regular marijuana use.

Differences over time:

- There were no significant changes in experimental, occasional, and regular marijuana use from 2012 to 2014.
- Among Grade 10 students, there was an increase in occasional marijuana use from 2002 through 2014.



Survey Question: During the past 30 days, on how many days did you: Use marijuana or hashish (weed, hash, pot)?

Notes:

- Experimental marijuana use represents 1-2 days of use, occasional marijuana use represents 3-5 days, and regular marijuana use represents 6 or more days in the past 30 days.
- The description of marijuana changed from “grass, hash, pot” to “weed, hash, pot” in 2014.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Average Age of First Marijuana Use

Some students begin experimenting with marijuana at an early age. Early (12-14 years of age) initiation of drug use, such as marijuana, is associated with a greater risk of developing an addiction and drug abuse than initiation during adulthood (Chen, 2009).

Among Grade 10 students who reported lifetime use of marijuana, the average age they first used marijuana was at 13.3 years of age.

Table 14: Average Age of First Marijuana Use in 2014

Behavior	Mean Age of First Reported Use		
	Grade 8	Grade 10	Grade 12
Used marijuana	12.0 (± 0.08)	13.3 (± 0.1)	14.5 (± 0.1)

Survey Question: How old were you the first time you used marijuana?

Notes:

- Age of first use is calculated by excluding students who responded “Never had,” used marijuana and calculating the mean age of use among those who used marijuana at any age.
- The word “smoked” was changed to “used” in 2014.

Source: HYS 2014.

Perception of Access to Marijuana

A study based on a national survey (Caulkins and Pacula, 2006) found that among people of all ages, most marijuana users obtain the drug for free (59 percent), from a friend or relative (88 percent), and through indoor transactions (87 percent). Only 6 percent reported purchasing marijuana from a stranger. The perceived ease of availability of marijuana among Washington State youth has been consistently below the national average (Monitoring the Future, 2013).

HYS asks students how hard it would be for them to get marijuana if they wanted it. In 2014, 86 percent of Grade 6 students, 66 percent of Grade 8 students, 29 percent of Grade 10 students, and 17 percent of Grade 12 students reported that it would be very hard to get marijuana.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was less likely to perceive that marijuana would be very hard to get.

Differences by gender:

- Grade 6 females were more likely than males to perceive that marijuana would be very hard to get.
- Grade 8 males were more likely than females to perceive that marijuana would be very hard to get.

Differences over time:

- Among Grade 8 students, there was an increase in the perception that getting marijuana would be very hard from 2012 to 2014.
- There were no significant trends in the perception that getting marijuana would be very hard for any grades from 2002 through 2014.



Survey Question: If you wanted to get some marijuana, how easy would it be for you to get some?

Note: Percentages represent students who reported that it would be very hard to get marijuana if they wanted some.

Source: WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Usual Type of Marijuana

In 2014, students were asked if they used marijuana, and how they usually used it. Among those who said they used marijuana:


- 65 percent of Grade 8 students, and 74 percent of Grade 10 and 12 students smoked it.
- 15 percent of Grade 8 and 10 students, and 12 percent of Grade 12 students ate it.
- 7 percent of Grade 8 students, and 3 percent of Grade 10 and 12 students drank it.
- 7 percent of Grade 8 and 12 students, and 5 percent of Grade 10 students vaporized it.
- 6 percent of Grade 8 students, 3 percent of Grade 10, and 4 percent of Grade 12 used it some other way.

Differences by grade level:

- Grade 10 and 12 students were more likely than Grade 8 students to usually smoke it.
- Grade 8 students were more likely than Grade 10 and 12 students to usually drink it.
- Grade 8 students were more likely than Grade 10 students to use it some other way.

Differences by gender:

- Grade 10 and 12 females were more likely than males to smoke it.
- Grade 10 and 12 males were more likely than females to drink it.
- Grade 12 males were more likely than females to vaporize it.
- Grade 12 males were more likely than females to use it some other way.



Survey Question: During the past 30 days, if you used marijuana, how did you usually use it?

Notes:

- Percentages represent students who reported that they used marijuana in one of the specified ways.
- Students who reported “did not use marijuana in the past 30 days” were not included in the results.
- The sample sizes for the 2014 results in this figure are: 333 Grade 8; 688 Grade 10; and 791 Grade 12 students.

Source: HYS 2014.

Perception of Risk from Regular Marijuana Use

Long-term trend data from Monitoring the Future suggests that perceived risk of marijuana use is a leading indicator of actual use. That is, during the 1970s, and again in the 1990s, as the perception of risk fell, the use of marijuana rose (Johnston, O'Malley, Bachman, and Schulenberg, 2007).

In 2014, 50 percent of Grade 6 students, 53 percent of Grade 8 students, 36 percent of Grade 10 students, and 26 percent of Grade 12 students reported there was great risk in using marijuana regularly.

Differences by grade:


- Grade 6 and 8 students were more likely than Grade 10 and 12 students to perceive great risk in regular marijuana use.
- Grade 10 students were more likely than Grade 12 students to perceive great risk in regular marijuana use.

Differences by gender:

- Grade 8, 10 and 12 females were more likely than males to perceive great risk in regular marijuana use.

Differences over time:

- Among Grade 6, 10 and 12 students, there were decreases in the perception of great risk from using marijuana regularly from 2012 to 2014.
- Among Grade 6 and 10 students there were decreases in the perception of great risk from using marijuana regularly from 2002 through 2014. Among Grade 8 and 12 students there were decreases from 2006 to 2014 in the perception of great risk from using marijuana regularly.



Survey Question: How much do you think people risk harming themselves if they: Use marijuana regularly? (at least once or twice a week)

Notes:

- Percentages represent students who reported there is great risk from regular marijuana use.
- The word "smoke" was changed to "use" in 2014.

Source: WSSAHB 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Other Drugs Not Including Alcohol, Tobacco, or Marijuana

The Healthy Youth Survey also tracks drugs that are less common than alcohol, tobacco, and marijuana. The drugs that are included in the survey can change over time. For instance, early surveys included prescription drugs, but they were eliminated as concerns about party drugs grew. In 2014, several new questions regarding prescription drugs abuse and misuse were added in response to heightened national and local awareness of this issue among youth.

30-Day Other Drug Use (Not Including Alcohol, Tobacco, or Marijuana)

In 2014, 1 percent of Grade 6 students, 2 percent of Grade 8 students, 4 percent of Grade 10 students, and 7 percent of Grade 12 students reported using an illegal drug other than alcohol, tobacco or marijuana in the past 30 days.

Differences by grade level:


- Among Grade 6, 8, 10 and 12 students, as grade levels increase, each grade was more likely to use other illegal drugs in the past 30 days.

Differences by gender:

- Grade 6, 10 and 12 males were more likely than females to use other illegal drugs in the past 30 days.

Differences over time:

- Among Grade 8 students, there was a decrease in the 30-day other illegal drugs use from 2012 to 2014.
- There were no significant trends in 30-day other illegal drug use for any grades from 2004 through 2014.



Survey Question: During the past 30 days, on how many days did you: Not counting alcohol, tobacco, or marijuana, use another illegal drug?

Note: Percentages represent students who used other illegal drugs on any days in the past 30 days.

Source: HYS 2004, 2006, 2008, 2010, 2012 and 2014.

Prescription Opiate (Painkiller) Use

Awareness and concern are growing regarding the use of prescription drugs that young people are using to get high.

In 2014, painkiller use “to get high” in the past 30 days was reported by 2 percent of Grade 8 students, 5 percent of Grade 10, and 6 percent of Grade 12 students.

Differences by grade level:


- Grade 8 students were less likely than Grade 10 and 12 students to use painkillers to get high in the past 30 days.

Differences by gender:

- Grade 12 males were more likely than females to use painkillers to get high in the past 30 days.

Differences over time:

- Among Grade 8, 10, and 12 students, there were decreases in using painkillers to get high in the past 30 days from 2012 to 2014.
- Among Grade 10 students, there was a decrease in using painkillers to get high in the past 30 days from 2006 through 2014.



Survey Question: During the past 30 days, on how many days did you: Use a pain killer to get high, like Vicodin, OxyContin (sometimes called Oxy or OC) or Percocet (sometimes called Percs)?

Note: Percentages represent students who reported using painkillers to get high on any days in the past 30 days.

Source: HYS 2006, 2008, 2010, 2012 and 2014.

Prescription Drug Misuse


In 2014, using non-subscribed prescription drugs in the past 30 days was reported by 4 percent of Grade 8 students, 8 percent of Grade 10, and 9 percent of Grade 12 students.

Differences by grade level:

- Grade 8 students were less likely than Grade 10 and 12 students to use non-subscribed prescription drugs in the past 30 days.

Differences by gender:

- Grade 8 females were more likely than males to use non-subscribed prescription drugs in the past 30 days.



Survey Question: During the past 30 days, on how many days did you: Use prescription drugs not prescribed to you
Note: Percentages represent students who reported using non-subscribed prescription drugs on any days in the past 30 days.

Source: HYS 2014.

Lifetime Methamphetamine Use

Methamphetamine, a subclass of amphetamines, was at one time called “speed.” During the past several years, media reports have sometimes referred to methamphetamine use as an epidemic. This reflects the environmental and familial consequences of methamphetamine production. Nationally, methamphetamine use has been declining, including most recently among young adults (Substance Abuse and Mental Health Services Administration, 2009).

In 2014, Lifetime methamphetamine use was reported by 3 percent of Grade 8 students, and 4 percent of Grade 10 and Grade 12 students.

Differences by grade level:


- Grade 10 and 12 were more likely than Grade 8 students to have used methamphetamines in their lifetime.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to have used methamphetamines in their lifetime.

Differences over time:

- Among Grade 12 students, there was a decrease in lifetime methamphetamine use from 2012 to 2014.
- Among Grade 12 students, there was a decrease in lifetime methamphetamine use from 2002 through 2014.



Survey Question: Have you ever, even once in your lifetime, used any of the following drugs? Methamphetamines (meth, crystal meth, ice, crank) Do not include other types of amphetamines.

Notes: Percentages represent students who had ever used methamphetamines in their life.

Source: WSSAHB 1990 and 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Lifetime Inhalant Use

Inhalants are fumes or gases that can be inhaled for the purpose of getting high. Inhalants include common household products such as glue, gasoline, solvents such as nail polish remover, and propellants in certain products such as whipped cream dispensers.

In 2014, 2 percent of Grade 6 students, 4 percent of Grade 8 students, and 8 percent of Grade 10 and Grade 12 students reported ever using inhalants.

Differences by grade level:


- Grade 8, 10 and 12 students were more likely than Grade 6 students to use inhalants in their lifetime.
- Grade 10 and 12 students were more likely than Grade 8 students to use inhalants in their lifetime.

Differences by gender:

- Grade 10 and 12 males were more likely than females to report to use inhalants in their lifetime.

Differences over time:

- Among Grade 8 and 10 students, there was a decrease in lifetime inhalant use from 2012 to 2014.
- Among Grade 6 students, there was a decrease in lifetime inhalant use from 2004 through 2014.



Survey Questions:

- How old were you the first time you: Used inhalants?
- Have you ever, even once in your lifetime, used inhalants (things you sniff to get high)?

Note: Percentages represent students who had ever used inhalants at any age in their life (Grades 8, 10 and 12) or had ever used inhalants once in their life (Grade 6).

Source: SADUS 1988 and 1990, WSSAHB 1992, 1995, 1998 and 2000, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Lifetime Heroin Use

In 2014, lifetime heroin use was reported by 3 percent of Grade 8, 10 and 12 students.

Differences by grade level:


- There were no differences in lifetime heroin use by grade level.

Differences by gender:

- Grade 8 and 10 males were more likely than females to use heroin in their lifetime.

Differences over time:

- Among Grade 12 students, there was a decrease in lifetime heroin use from 2012 to 2014.
- There were no significant trends in lifetime heroin use for any grades from 2002 through 2014.



Survey Question: Have you ever, even once in your lifetime, used any of the following drugs: Heroin?

Note: Percentages represent students who had ever used heroin in their life.

Source: WSSAHB 1998 and 2000, YRBS 1999, HYS 2006, 2008, 2010, 2012, and 2014.

Lifetime Cocaine Use

In 2014, lifetime cocaine use was reported by 3 percent of Grade 8 students, 4 percent of Grade 10, and 7 percent of Grade 12 students.

Differences by grade level:


- Among Grade 8, 10 and 12 students, as grade levels increase, each grade was more likely to use cocaine in their lifetime.

Differences by gender:

- Grade 8, 10 and 12 males were more likely than females to use cocaine in their lifetime.

Differences over time:

- Among Grade 10 students, there was a decrease in cocaine use from 2012 to 2014.
- There were no significant trends in lifetime cocaine use among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: Have you ever, even once in your lifetime, used any of the following drugs: Cocaine?

Note: Percentages represent students who had ever used cocaine in their life.

Source: WSSAHB 1988, 1990, 1992, 1995, 1998 and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Lifetime Steroid Use

In 2014, lifetime steroid use was reported by 2 percent of Grade 8 students, and 3 percent of Grade 10 and Grade 12 students.

Differences by grade level:


- Grade 10 and 12 students were more likely than Grade 8 students to use steroids in their lifetime.

Differences by gender:

- Grade 10 and 12 males were more likely than females to use steroids in their lifetime.

Differences over time:

- Among Grade 10 and 12 students, there were decreases in lifetime steroid use from 2012 to 2014.
- There were no significant trends in lifetime steroid use among students in grades 8, 10 and 12 from 2002 through 2014.



Survey Question: Have you ever, even once in your life, used steroids (muscle builders) without a doctor's prescription?

Note: Percentages represent students who had ever used steroids, without a doctor's prescription, in their life.

Source: WSSAHB 1988, 1990, 1992, 1995, 1998 and 2000, YRBS 1999, HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

11. Risk and Protective Factors

This chapter covers a broad set of questions about health behaviors, and about the risk factors and protective factors associated with them. Risk factors are characteristics of individuals and their families, schools, and communities that make them more vulnerable to ill health and poor lifestyle choices. Similarly, protective factors exert a positive influence or buffer against the negative influence of risk in these social environments. The Healthy Youth Survey includes many questions directly related to health, but most of the risk and protective factors measured in the survey are associated with behaviors such as substance use, violence, and dropping out of school. The presence of multiple risk factors predicts an increased likelihood that an individual will engage in these behaviors, whereas the presence of protective factors helps to buffer the effect of risk factors and increase resilience.

Research over several decades has identified risk factors that are associated with increased likelihood of health risk behaviors including alcohol, tobacco, and other drug abuse (Dryfoos, 1991; Hawkins et al., 1992; Kandel, Daview, Karus, and Yamagucchi, 1986); violence and delinquent behaviors (Bensley, Spieker, VanEenwyk, and Schoder, 1999; Brewer, Hawkins, Catalano, and Neckerman, 1995; Herrenkohl, Chung, and Catalano, 2004; Wasserman et al., 2003); and driving after drinking (Sabel, Bensley, and VanEenwyk, 2004).

Another body of research has focused on young people's ability to overcome the odds that challenge them (Werner and Smith, 1989) and to succeed in spite of a preponderance of risk in their environments. Benard (1991) summarized this literature on protective factors, citing the longitudinal research of Werner and Smith and Rutter (1979) in the formulation of a construct termed resilience. Resnick et al. (1997) found that parent-family connectedness and perceived school connectedness were protective against every health risk behavior measured in their study except history of pregnancy. Parental expectations regarding school achievement and school connectedness were also associated with lower levels of health risk behaviors (except in the case of suicide, in which only parent-family connectedness was protective).

Using these multiple strands of research, Hawkins and Catalano at the University of Washington's Social Development Research Group developed a theoretical framework based on a model of social development which hypothesizes that strong bonds serve as protective factors against behaviors that violate socially accepted standards. Attachment (a positive emotional link) and commitment (a personal investment) are the components of the social bond. The theory hypothesizes that when social groups produce strong bonds of attachment and commitment in members and promote clear standards for behavior, these groups increase behavior consistent with those standards and prevent behavior that violates them (Hawkins, Guo, Hill, Battin-Pearson, and Abbott, 2001).

By addressing risk and protective factors, families, schools, and communities can help promote positive social development. Early and sustained intervention through the elementary grades should put children on a developmental trajectory leading to more positive outcomes and fewer problem behaviors over the long term. These risk and protective factors represent promising inputs for prevention and intervention programs and policies.

The data presented in this chapter represent Washington State as a whole. The level of these indicators of risk and protection likely vary by community. Communities can compare community-level data to state-level data—and to county-level data where available—to determine which risk and protective factors are priorities for their communities to address. Communities can then implement prevention services for specific populations or geographical areas where risk exposure is high and protection is low.

The 1995, 1998, 2000, 2002, 2004, 2006, 2008, 2010, 2012 and 2014 survey administrations in Washington included substantial coverage of risk and protective factors using standardized assessment tools developed by the Social Development Research Group (Arthur et al., 1998; Arthur, Hawkins, Pollard, Catalano, and Baglioni, 2002) and published in their *Communities That Care* survey. These risk and protective factors are organized into four domains of influence: community, family, school, and peer-individual.

More information on the risk and protective factors used in the HYS is available at: <http://www.askhys.net/Reports/Additional>

HYS 2014 assessed six risk factors among students in Grade 6 and ten risk factors among students in Grades 8, 10, and 12 (see Table 15).

Table 15
Risk Factors Included in 2014

Domain	Risk Factor
Community	Laws and norms favorable toward drug use
	Perceived availability of drugs
School	Academic failure
	Low commitment to school
Peer-Individual	Perceived risk of drug use
	Early initiation of drug use ^S
	Favorable attitudes toward drug use
	Friends' use of drugs ^S
Family	Poor family management ^S
	Parental attitudes favorable towards drug use ^S

^S Included only on the secondary version of the survey (Grades 8, 10 and 12).

The HYS 2014 administration also assessed five protective factors among students in Grade 6 and seven protective factors among students in Grades 8, 10, and 12 (see Table 16).

Table 16
Protective Factors Included in 2014

Domain	Protective Factor
Community	Opportunities for prosocial involvement ^S
	Rewards for prosocial involvement ^E
School	Opportunities for prosocial involvement ^S
	Rewards for prosocial involvement
Peer-Individual	Social skills ^S
	Belief in the moral order ^S
	Interaction with prosocial peers ^S
	Prosocial involvement ^E
Family	Opportunities for prosocial involvement
	Rewards for prosocial involvement ^E


^S Included only on the secondary version of the survey (Grades 8, 10 and 12).

^E Included only on the elementary version of the survey (Grade 6).

This chapter presents HYS 2014 results for the assessment of risk and protection at each grade level in the community, family, school, and peer-individual domains. The relationships between risk and protective factors and the major health risk behaviors of substance use and violent and delinquent behavior are also presented. Readers should remember that all results are based on student self-report and therefore represent perceptions of risk and protection, which might not be accurate. Furthermore, the statistical relationships between risk and protective factors and health risk behaviors are not necessarily causal. Rather, the statistical relationships indicate an association or co-occurrence of these factors and behaviors. Both the risk factor and the behavior may be associated with a third factor such as poverty or other factors that were not addressed in this study. Each risk and protective factor scale is calculated as the average score of the students' responses to one or more questions. Students whose scores placed them above a cut point, determined by the Social Development Research Group's research, were considered at risk on a given risk factor or resilient on a given protective factor.


Research has also suggested a cumulative effect in the influence of risk and protection on these health risk behaviors (Bry, McKeon, and Pandina, 1982; Newcomb, Maddahian, and Skager, 1987; Werner and Smith, 1989). In addition to examining the specific influence of a given risk or protective factor, examining the relationship between multiple risk or protective factors and these behaviors is important. This examination helps illustrate whether students who are at high risk on more risk factors are more likely to engage in health risk behaviors than students who are at high risk on fewer factors. An examination of the relationship between multiple risk or protective factors and health risk behaviors also helps show whether students who are well protected are less likely to engage in these behaviors than students who are less protected.

The following chart shows the relationship between the number of risk factors present and the use of alcohol, cigarettes, and marijuana for students in Grade 8. This relationship also holds true for grades 10 and 12. The most obvious interpretation is the clear, linear relationship between the number of risk factors present and the prevalence of lifetime and 30-day alcohol, cigarette, and marijuana use. As the number of risk factors for individual students increased, the more likely they were to use alcohol, cigarettes and marijuana. These findings are consistent with the findings from the 1995, 1998, 2000, 2002, 2004, 2006, 2008, 2010, and 2012 survey administrations.



Note: Percentages represent students who reported using alcohol, cigarettes, or marijuana in their lifetime or in the past 30 days according to each number or risk factors (0 through 10).
Source: HYS 2014.

The following chart is a similar display relating the presence of protective factors to the use of alcohol, cigarettes, and marijuana for students in Grade 8. This relationship also holds true for grades 10 and 12. Again, the overall relationship is strong: increased levels of protection were clearly associated with lower rates of alcohol, cigarette, and marijuana use. Protective factors have also been found to have a buffering effect on the presence of risk factors (DeWit, Silverman, Goodstadt, and Stoduto, 1995; Gabriel, Deck, Einspruch, and Nickel, 1997; Jessor, Van den Bos, Vanderryn, Costa, and Trubin, 1995). These findings are consistent with the findings from the 1995, 1998, 2000, 2002, 2004, 2006, 2008, 2010, and 2012 survey administrations.



Note: Percentages represent students who reported using alcohol, cigarettes, or marijuana in their lifetime or in the past 30 days according to each number of protective factors (0 through 7).

Source: HYS 2014.

Community Domain: Risk Factors

HYS 2014 assessed two risk factors in the community domain:

- *Laws and norms favorable toward drug use.* The policies a community holds in relation to health and problem behaviors are communicated through laws, social practices, and expectations, and are related to use.
 - Among Grade 8, 10 and 12 students, there were decreases from 2002 through 2014.
- *Perceived availability of drugs.* Perceptions of the availability of alcohol and other drugs have been shown to predict use of these substances.
 - Among Grade 10 and 12 students, there were decreases from 2002 through 2014.

Table 17
Profile of Community Risk Factors,
Percent of Youth at Risk, Grades 6, 8, 10, and 12 from 2002–2014

Risk Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Laws and norms favorable toward drug use	37.1	37.1	37.0	35.9	36.5	36.5	33.9	33.0	29.8	28.2	28.3	27.7	26.4	23.1	38.7	40.1	39.1	36.7	34.5	31.4	31.7	39.3	37.3	35.8	34.4	32.5	32.4	31.2
Perceived availability of drugs	23.6	22.5	24.6	23.5	22.6	19.5	18.7	29.3	23.0	20.9	24.8	24.1	22.8	17.1	35.5	31.8	32.7	34.2	34.4	28.4	26.5	45.2	40.5	38.1	39.4	38.1	36.2	31.7

Notes:

- Percentages represent students who are at risk based upon their risk factor scale scores.
- Dashes (–) indicate that the risk factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the risk factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Community Domain: Protective Factors

HYS 2014 assessed two protective factors in the community domain (only one for Grade 6 and only one for Grades 8, 10 and 12).

- *Opportunities for prosocial involvement.* Youth need opportunities to participate meaningfully in activities in the community. Note: in 2002 the items in this scale were modified for the Healthy Youth Survey and are therefore different than those used by the Social Development Research Group.
 - There were no significant trends for any grades from 2002 through 2014 in opportunities for prosocial involvement.
- *Rewards for prosocial involvement.* Youth need rewards for positive participation in prosocial activities.
 - Among Grade 6 students, there was a decrease from 2002 through 2014.

Table 18
Profile of Community Protective Factors,
Percent of Youth Protected, Grades 6, 8, 10, and 12 from 2002–2014

Protective Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Opportunities for prosocial involvement	25.8	–	–	–	–	–	–	50.7	72.3	69.2	66.6	67.5	73.2	75.3	46.6	72.4	66.1	69.2	71.1	75.2	75.4	42.7	70.9	69.3	71.3	76.0	76.3	77.7
Rewards for prosocial involvement	48.0	38.6	37.9	36.4	35.9	37.4	36.9	54.9	56.6	54.0	54.0	–	–	–	60.3	60.4	56.2	62.2	–	–	–	55.1	56.6	56.8	62.0	–	–	–

Notes:

- Percentages represent students who are protected based upon their protective factor scale scores.
- Dashes (–) indicate that the protective factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the protective factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

School Domain: Risk Factors

HYS 2014 assessed two risk factors in the school domain. Readers should note that the items used to create the low commitment to school risk factor changed slightly in 2002.

- *Academic failure.* Children fail in school for many reasons, but research indicates that the very experience of failure—regardless of whether the failure is linked to the students’ abilities—places them at higher risk for negative behavior.
 - There were no significant trends for any grades from 2002 through 2014 in academic failure.
- *Low commitment to school.* When young people cease to see the school role as viable, they are at higher risk of engaging in the health risk behaviors.
 - There were no significant trends for any grades from 2002 through 2014 in low commitment to school.

Table 19
Profile of School Risk Factors,
Percent of Youth at Risk, Grades 6, 8, 10, and 12 from 2002–2014

Risk Factor	Grade 6								Grade 8								Grade 10								Grade 12							
	2002	2004	2006	2008	2010	2012	2014		2002	2004	2006	2008	2010	2012	2014		2002	2004	2006	2008	2010	2012	2014		2002	2004	2006	2008	2010	2012	2014	
Academic failure	41.2	40.6	41.5	42.4	41.9	37.8	39.5		47.3	48.2	45.9	47.5	46.8	45.3	43.9		46.8	47.2	50.6	48.2	47.7	45.3	46.4		48.5	46.6	50.1	51.4	49.1	47.5	49.2	
Low commitment to school	40.5	44.4	52.0	43.0	38.9	36.8	38.1		34.4	37.1	36.2	38.6	35.6	31.8	31.9		37.3	40.7	39.9	38.2	37.8	33.1	38.3		37.6	42.2	40.8	41.4	36.5	36.1	40.4	

Notes:

- Percentages represent students who are at risk based upon their risk factor scale scores.
- Dashes (–) indicate that the risk factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the risk factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

School Domain: Protective Factors

HYS 2014 assessed two protective factors in the school domain (only one for Grade 6).

- *Opportunities for prosocial involvement.* When young people are given more opportunities to participate meaningfully in important activities at school, they are less likely to engage in problem behaviors.
 - Among Grade 8 students, there was an increase from 2008 to 2014 in opportunities for prosocial involvement. Among Grade 10 students, there was an increase in opportunities for prosocial involvement from 2002 through 2014.

- *Rewards for prosocial involvement.* When young people are recognized and rewarded for their contributions at school, they are less likely to be involved in health risk behaviors.
 - Among Grade 6 students, there was a decrease in rewards for prosocial involvement from 2002 through 2014.

Table 20
Profile of School Protective Factors,
Percent of Youth Protected, Grades 6, 8, 10, and 12 from 2002–2014

Protective Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Opportunities for prosocial involvement	–	–	–	–	–	–	–	62.6	62.2	64.0	59.8	62.6	65.7	70.0	59.6	58.5	57.7	59.0	61.8	66.5	65.2	63.5	61.2	61.6	60.7	64.0	65.5	68.3
Rewards for prosocial involvement	50.5	52.3	52.8	49.8	49.5	49.6	44.9	52.1	53.4	56.5	53.1	49.0	51.1	52.8	61.4	61.2	61.1	63.5	58.4	60.1	57.5	45.8	44.6	45.4	46.8	45.3	46.2	43.2

Notes:

- Percentages represent students who are protected based upon their protective factor scale scores.
- Dashes (–) indicate that the protective factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the protective factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Peer-Individual Domain: Risk Factors

HYS 2014 assessed four risk factors in the peer-individual domain (only two for Grade 6).

- *Early initiation of drug use.* Research shows that the earlier an individual begins using alcohol, tobacco, and other drugs, the more likely he or she is to develop drug use problems as an adult.
 - Among Grade 8 students, there was a decrease in early initiation of drug use from 2002 through 2014. Among Grade 10 and 12 students, there were decreases in early initiation of drug use from 2002 through 2014.
- *Favorable attitudes toward drug use.* Young people who have positive or accepting attitudes toward drug use are more likely to engage in a variety of health risk behaviors.
 - Among Grade 6 students, there was a decrease in favorable attitudes toward drug use from 2002 through 2014. Among Grade 12 students, there was an increase from 2006 to 2014 in favorable attitudes toward drug use.
- *Perceived risk of use.* Young people who do not perceive a risk in using alcohol, tobacco, and other drugs are at higher risk of engaging in substance use.
 - Among Grade 6, 10 and 12 students, there were increases in the percentage of students at risk due to decreased perception of risk in substance use from 2002 through 2014.
- *Friends' use of drugs.* Young people whose friends use drugs are more likely to engage in health risk behaviors.
 - Among Grade 8 and 12 students, there were decreases in friends' use of drugs from 2002 through 2014.

Table 21
Profile of Peer-Individual Risk Factors,
Percent of Youth at Risk, Grades 6, 8, 10, and 12 from 2000–2014

Risk Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Early initiation of drug use	–	–	–	–	–	–	–	27.4	24.6	19.8	20.8	20.1	18.2	13.7	32.5	29.2	31.4	29.3	26.6	22.2	20.5	37.5	33.0	32.9	32.3	27.9	26.4	22.8
Favorable attitudes towards drug use	22.6	22.2	21.4	20.9	20.9	18.3	19.6	27.8	27.2	22.9	24.8	24.5	26.6	23.8	37.6	35.0	37.2	37.2	36.7	37.0	41.0	40.8	36.7	34.8	37.7	37.9	40.0	39.9
Perceived risk of drug use	32.3	30.3	32.7	31.9	40.3	37.7	40.7	38.3	35.0	33.0	33.9	37.7	39.2	37.9	34.8	33.7	35.0	35.6	39.0	38.1	41.2	43.4	38.4	40.6	43.3	48.0	49.4	52.8
Friends' use of drugs								28.5	27.2	22.8	25.6	24.1	23.2	15.3	30.7	27.6	29.7	28.8	29.0	25.1	23.0	36.9	25.9	26.5	27.2	28.5	25.5	22.5

Notes:

- Percentages represent students who are at risk based upon their risk factor scale scores.
- Dashes (–) indicate that the risk factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- * Items in the risk factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Peer-Individual Domain: Protective Factors

HYS 2014 assessed four protective factors in the peer-individual domain (only one for Grade 6).

- *Social skills.* Young people who are socially competent and engage in positive interpersonal relations with their peers are less likely to participate in negative health risk behaviors.
 - Among Grade 12 students, there was a decrease in social skills from 2002 through 2014.
- *Belief in the moral order.* Young people who have a belief in what is right or wrong are at lower risk for engaging in problem behaviors.
 - Among Grade 8 students, there was an increase from 2008 to 2014 in belief in the moral order.
- *Interaction with prosocial peers.* Young people who interact with peers who are a positive influence are at lower risk for engaging in problem behaviors.
 - Among Grade 8 students, there was an increase in interaction with prosocial peers from 2004 through 2012.
- *Prosocial involvement.* Young people who are engaged in positive social activities are at lower risk for engaging in problem behaviors.
 - There was no significant trend from 2004 through 2012 (6th Grade only).

Table 22
Profile of Peer-Individual Protective Factors,
Percent of Youth Protected, Grades 6, 8, 10, and 12 from 2002–2014

Protective Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Social skills	–	–	–	–	–	–	–	69.2	70.7	71.1	68.8	65.0	66.9	73.8	64.0	60.8	56.9	58.0	53.8	58.8	61.1	67.2	70.3	67.1	68.4	47.5	53.0	54.9
Believe in the moral order	–	–	–	–	–	–	–	66.1	64.2	65.5	61.2	64.1	67.2	71.8	71.4	68.6	65.5	66.8	69.5	74.7	72.9	55.7	55.4	53.2	53.2	54.0	57.9	55.4
Interaction with prosocial peers	–	48.4	46.2	43.2	–	–	–	–	54.7	55.8	57.0	57.0	59.4	64.5	–	56.9	55.3	55.0	56.6	60.5	57.7	–	54.1	52.1	52.6	55.0	55.7	50.9
Prosocial involvement	–	43.3	43.6	40.4	39.7	43.7	40.9	–	40.0	54.0	–	–	–	–	–	45.1	54.3	–	–	–	–	–	43.3	49.7	–	–	–	–

Notes:

- Percentages represent students who are protected based upon their protective factor scale scores.
- Dashes (–) indicate that the protective factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the protective factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012, and 2014.

Family Domain: Risk Factors

HYS 2014 assessed two risk factors in the family domain (both only for Grade 8, 10 and 12).

- *Poor family management.*
 - Among Grade 8 students, there was a decrease from 2008 to 2014 in poor family management.

- *Parental attitudes favorable towards drug use.*
 - There were no significant trends among students in grades 8, 10 and 12 from 2004 through 2014.

Table 23
Profile of Family Risk Factors,
Percent of Youth at Risk, Grades 6, 8, 10, and 12 from 2002–2014

Risk Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Poor family management	-	-	-	-	-	-	-	39.2	38.4	37.4	39.2	36.0	33.7	30.6	36.6	38.7	42.5	42.8	39.3	32.3	32.8	43.8	42.6	43.4	43.5	38.8	38.2	34.4
Parental attitudes favorable towards drug use	-	-	-	-	-	-	-	-	31.2	-	26.6	21.5	23.9	22.2	-	41.8	-	44.4	36.8	37.1	40.5	-	41.7	-	44.2	36.4	41.2	41.3

Notes:

- The family domain was measured on an optional tear-off page prior to the 2014 HYS, not all of the participating schools asked these questions and the number of students who answered the questions in this domain was smaller than the numbers of respondents for the other domains from 2002 to 2012.
- Percentages represent students who are at risk based upon their risk factor scale scores.
- Dashes (-) indicate that the risk factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the risk factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

Family Domain: Protective Factors

HYS 2014 assessed two protective factors in the family domain.

- *Opportunities for prosocial involvement.*
 - Among Grade 6 students, there was a decrease in opportunities for prosocial involvement from 2002 through 2014.
- *Rewards for prosocial involvement.*
 - Among Grade 6 students, there was a decrease in rewards for prosocial involvement from 2002 through 2014.

Table 24
Profile of Family Protective Factors,
Percent of Youth Protected, Grades 6, 8, 10, and 12 from 2002–2014

Protective Factor	Grade 6							Grade 8							Grade 10							Grade 12						
	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014	2002	2004	2006	2008	2010	2012	2014
Opportunities for prosocial involvement	58.1	58.5	54.2	54.8	53.2	55.6	52.6	63.4	–	66.6	61.3	63.6	66.3	68.5	56.7	–	53.5	51.7	55.6	58.7	57.8	56.7	–	53.6	53.4	53.7	55.8	57.8
Rewards for prosocial involvement	62.2	62.5	58.2	58.6	56.8	57.4	53.6	66.0	–	69.6	60.6	62.3	61.6	–	60.3	–	54.9	51.5	52.7	54.9	–	57.1	–	52.7	52.3	49.7	50.8	–

Notes:

- The family domain was measured on an optional tear-off page prior to the 2014 HYS, not all of the participating schools asked these questions and the number of students who answered the questions in this domain was smaller than the numbers of respondents for the other domains from 2002 to 2012.
- Percentages represent students who are protected based upon their protective factor scale scores.
- Dashes (–) indicate that the protective factor was not included in the survey that year.
- Changes that are statistically significant at the 95 percent confidence level from the previous year are bolded.
- ^x Items in the protective factor changed over time; the result is not comparable.

Source: HYS 2002, 2004, 2006, 2008, 2010, 2012 and 2014.

References

- Arlen Egley, Jr., James C. Howell, and Meena Harris. Juvenile Justice Fact Sheet 4: Highlights of the 2012 National Youth Gang Survey. Accessed December 2015:
[https://www.nationalgangcenter.gov/PublicationsArthur, M.W., Hawkins, J.D., Catalano, R.F., and Pollard, J.A. \(1998\). *Student survey of risk and protective factors and prevalence of alcohol, tobacco, and other drug use*. Seattle, WA: Social Development Research Group.](https://www.nationalgangcenter.gov/PublicationsArthur, M.W., Hawkins, J.D., Catalano, R.F., and Pollard, J.A. (1998). Student survey of risk and protective factors and prevalence of alcohol, tobacco, and other drug use. Seattle, WA: Social Development Research Group.)
- Arthur, M.W., Hawkins, J.D., Pollard, J.A., Catalano, R.F., and Baglioni, A.J. (2002). *Measuring risk and protective factors for substance use, delinquency and other adolescent problem behaviors: The Communities That Care Youth Survey*. *Evaluation Review*, 26(2), 575–601.
- Benard, B.L. (1991). *Fostering resiliency in kids: Protective factors in the family, school, and community*. San Francisco, CA: Far West Laboratory for Educational Research and Development.
- Bensley, L. (1997, August). *Reliability and validity of the Youth Risk Behavior Survey: Draft briefing paper*. Olympia, WA: Washington State Department of Health Office of Epidemiology.
- Bensley, L., VanEenwyk, J., Schoder, J., and Tollefsen, P. (2000). *Washington State Youth Risk Behavior Survey: 1999*. Olympia, WA: Washington State Department of Health.
- Brewer, D.D., Hawkins, J.D., Catalano, R.F., and Neckerman, H.J. (1995). *Preventing serious, violent, and chronic juvenile offending*. In Howell, J.C., Krisberg, B., Hawkins, J.D., and Wilson, J.J. A sourcebook: Serious, violent, and chronic juvenile offenders. Thousand Oaks, CA: Sage, 61–141.
- Brenner RA, Taneja GS, Haynie DL, Trumble AC, Qian C, Klinger RM, Klebanoff MA. (2009). *Association between swimming lessons and drowning in childhood: a case-control study*. *Arch Pediatr Adolesc Med*. Mar;163(3):203-10. doi: 10.1001/archpediatrics.2008.563
- Bry, B.H., McKeon, P., and Pandina, R.J. (1982). *Extent of drug use as a function of number of risk factors*. *Journal of Abnormal Psychology*. 91, 273–279.
- Buchmann, A.F., et al. (2009). *Impact of age at first drink on vulnerability to alcohol-related problems: testing the marker hypothesis in a prospective study of young adults*. *J. Psychiatr. Res.* 43, 1205–1212.
- Catalano R.F., Haggerty, K.P., Oesterle, S., Fleming, C.B., and Hawkins, J.D. (2004). *The Importance of Bonding to School for Healthy Development: Findings from the Social Development Research Group*. *Journal of School Health*, 74(7), 252-61.
- Caulkins, J. and Pacula, R. (2006). *Marijuana markets: Inferences from reports by the household population*. *Journal of Drug Issues*, 36(1), 173–200.
- Center on Hunger and Poverty (2002). *The Consequences of Hunger and Food Insecurity for Children: Evidence from Recent Scientific Studies*. Waltham, M.A.: Center on Hunger and Poverty, Heller School for Social Policy and Management, Brandeis University.
- Centers for Disease Control and Prevention. (2000). *Youth tobacco surveillance: United States, 1998–1999*. Retrieved from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss4910a1.htm>

- Centers for Disease Control and Prevention. (2009). *Youth Risk Behavior Surveillance—United States, 2007*. National Center for Chronic Disease Prevention and Health Promotion. Retrieved from <http://www.cdc.gov/healthyyouth/yrbs/index.htm>
- Centers for Disease Control and Prevention. (2010). *Injury Prevention and Control: Motor Vehicle Safety factsheet*. National Center for Chronic Disease Prevention and Health Promotion. Retrieved from http://www.cdc.gov/Motorvehiclesafety/teen_drivers/teendriv_ers_factsheet.html
- Chapman D.P., Perry G.S., and Strine, T.W. (2005). *The vital link between chronic disease and depressive disorders*. Preventing Chronic Disease. Retrieved from: http://www.cdc.gov/pcd/issues/2005/jan/04_0066.htm
- Chavez, P.R., Nelson, D.E., Naimi, T.S., Brewer, R.D. (2011). *Impact of a new gender-specific definition for binge drinking on prevalence estimates for women*. *Am J Prev Med*. Apr;40(4):468-71. doi: 10.1016/j.amepre.2010.12.008.
- Chen, C.-Y., Storr, C. L., & Anthony, J. C. (2009). *Early-onset drug use and risk for drug dependence problems*. *Addictive Behaviors*, 34(3),319-322. doi: 10.1016/j.addbeh.2008.10.021
- Courtney KE, Polich J. Binge Drinking in Young Adults: Data, Definitions, and Determinants. *Psychological bulletin*. 2009;135(1):142-156. Doi:10.1037/a0014414.
- Deck, D.D. and Nickel, P.N. (1989). *Substance abuse among public school students in Washington*. Olympia, WA: Office of Superintendent of Public Instruction.
- DeWit, D.J., Silverman, G., Goodstadt, M., and Stoduto, G. (1995). *The construction of risk and protective factor indices for adolescent alcohol and other drug use*. *Journal of Drug Issue*, 25(4), 837–863.
- Dilley, J. (2009). *School-based Health Interventions and Academic Achievement. Healthy Students, Successful Students*. Partnership Committee, Washington State Board of Health, Washington State Office of Superintendent of Public Instruction, Washington State Department of Health.
- Distefan, J., et al. (1998). *Parental influences predict adolescent smoking in the United States, 1989-1993*. *Journal of Adolescent Health*, 22, 466-74.
- Dye B.A., Tan S., Smith V., Lewis, B.G., Barker, L.K., Thornton-Evans G., et al. (2007). *Trends in oral health status: United States, 1988–1994 and 1999–2004*. National Center for Health Statistics. *Vital Health Stat* 11(248).
- Dryfoos, J.G. (1991). *Adolescents at risk: A summation of work in the field: Programs and policies*. *Journal of Adolescent Health*, 12(8), 630–637.
- Dubois S, Mullen N, Weaver B, Bédard M. (2015). *The combined effects of alcohol and cannabis on driving: Impact on crash risk*. *Forensic Sci Int*. Mar;248:94-100. doi:10.1016/j.forsciint.2014.12.018. Epub 2014 Dec 25. PubMed PMID: 25612879
- Eaton, D.K., Kann, L., Kinchen, S., Ross, J., Hawkins, J., Harris, W.A., et al. (2006). *Youth risk behavioral surveillance United States 2005: Surveillance summaries*. (MMWR 2006:55 No.SS-5). Atlanta, GA: Centers for Disease Control and Prevention.
- Einspruch, E.L. (2005). *Washington State Healthy Youth Survey 2004: Analytic report*. Olympia, WA: Office of Superintendent of Public Instruction.
- Einspruch, E.L., Deck, D.D., Nickel, P.R., and Hyatt, G. (2001). *Washington State Survey of Adolescent Health Behaviors 2000: Analytic report*. Olympia, WA: Office of Superintendent of Public Instruction.

- Einspruch, E.L., Gabriel, R.M., Deck, D.D., and Nickel, P.N. (1998). *Washington State Survey of Adolescent Health Behaviors 1998: Analytic report*. Olympia, WA: Office of Superintendent of Public Instruction.
- Einspruch, E.L. and Hyatt, G. (2004). *Washington State Survey of Adolescent Health Behaviors 2002: Analytic report*. Olympia, WA: Office of Superintendent of Public Instruction.
- Einspruch, E.L. and Pollard, J.P. (1993). *Washington State Survey of Adolescent Health Behaviors: 1988–1990*. Olympia, WA: Office of Superintendent of Public Instruction.
- Food Research and Action Center and Center on Hunger and Poverty (2003) *The Paradox of Hunger and Obesity in America*. Retrieved from: <http://www.frac.org/html/news>
- Gabriel, R.M. (1991). *Substance abuse among public school students in Washington State: 1988–1990*. Olympia, WA: Office of Superintendent of Public Instruction.
- Gabriel, R.M., Deck, D.D., Einspruch, E.L., and Nickel, P.N. (1995). *The findings of the Washington State Survey of Adolescent Health Behaviors: Analytic report*. Olympia, WA: Office of Superintendent of Public Instruction.
- Gabriel, R.M., Deck, D.D., Einspruch, E.L., and Nickel, P.N. (1997). *Risk and protective factors associated with alcohol, tobacco, and other drug use and violence*. Olympia, WA: Office of Superintendent of Public Instruction.
- Hampton, T. (2007). *Food insecurity harms health, well-being of millions in the United States*. JAMA, 298, 1851-1853.
- Hartman RL, Huestis MA. (2013). *Cannabis effects on driving skills*. Clin Chem. Mar;59(3):478-92. doi: 10.1373/clinchem.2012.194381. Epub 2012 Dec 7. Review.PubMed PMID: 23220273; PubMed Central PMCID: PMC3836260.
- Hawkins, J.D., Catalano, R.F., Jr., Barnard, K.E., Gottfredson, G.D., Holmes, A.B., and Miller, J.Y. (1992). *Communities that care: Action for abuse prevention*. San Francisco, CA: Jossey Bass.
- Hawkins, J.D., Catalano, R.F., and Miller, J.Y. (1992). *Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention*. Psychological Bulletin, 112(1), 64–105.
- Hawkins, D., Guo, J., Hill, K., Battin-Pearson, S., and Abbott, R. (2001). *Long-term effect of the Seattle social development intervention on school bonding trajectories*. Applied Developmental Science, 5(4), 225–236.
- Johnston, L.D., O'Malley, P.M., and Bachman, J.G. (1994). *National survey results on drug use: The Monitoring the Future Study 1975–1993. Volume I: Secondary students*. Rockville, MD: National Institute on Drug Abuse.
- Johnston, L.D., O'Malley, P.M., Bachman, J.G., and Schulenberg, J.E. (2007). *Monitoring the Future national results on adolescent drug use: Overview of key findings, 2006*. (NIH Publication No. 07-6202). Bethesda, MD: National Institute on Drug Abuse.
- Johnston, L. D., O'Malley, P. M., Miech, R. A., Bachman, J. G., Schulenberg, J. E. (2015). *Monitoring the Future national survey results on drug use: 1975-2014: Overview, key findings on adolescent drug use*. Ann Arbor: Institute for Social Research, The University of Michigan, 90pp.

- Hammond, C.J., et al. (2014). *Journal of Behavioral Addictions. An exploratory examination of marijuana use, problem-gambling severity, and health correlates among adolescents*. Jun;3(2):90-101. doi: 10.1556/JBA.3.2014.009.
- Kandel, D.B., Daview, M., Karus, D. and Yamagucchi, K. (1986). *The consequences in young adulthood of adolescent drug involvement: An overview*. *Archives of General Psychiatry*, 43, 746–754.
- Lobstein, T., et al. (2015). *Child and adolescent obesity: part of a bigger picture*. *The Lancet*, 385(9986), 2510–2520.
- Merikangas KR, Avenevoli S, Costello EJ, Koretz D, Kessler RC. (2009). The National Comorbidity Survey Adolescent Supplement (NCS-A): I. Background and Measures. *Journal of the American Academy of Child and Adolescent Psychiatry*, 48(4):367-369. doi:10.1097/CHI.0b013e31819996f1.
- McCambridge, J., McAlaney, J., Rowe, R. (2011) *Adult Consequences of Late Adolescent Alcohol Consumption: A Systematic Review of Cohort Studies*. *PLoS Med* 8(2): e1000413. doi:10.1371/journal.pmed.1000413.
- National Cancer Institute. (1992). *Smokeless Tobacco or Health: An International Perspective*. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute.
- National Center for Education Statistics. (2013). *Indicators of School Crime and Safety: 2013. Indicator 16: Students' Use of Marijuana on School Property and Anywhere*. Retrieved from: https://nces.ed.gov/programs/crimeindicators/crimeindicators2013/ind_16.asp, June 20015.
- National Cancer Institute. (2005). *Joinpoint regression program*. Retrieved from: <http://surveillance.cancer.gov/joinpoint>, June 20015.
- The National Center on Addiction and Substance Abuse, Columbia University. (2011). *Adolescent Substance Use: America's #1 Public Health Problem*. June 2011. Retrieved from: <http://www.casacolumbia.org/addiction-research/reports/adolescent-substance-use>
- National Highway Traffic Safety Administration (NHTSA), Dept. of Transportation (US). *Traffic safety facts 2012: Young Drivers*. Washington (DC): NHTSA. Retrieved from: <http://www-nrd.nhtsa.dot.gov/Pubs/812019.pdf>, April 2014.
- National Institute on Drug Abuse. (2001). *Monitoring the future: A continuing study of American youth*. Retrieved from: <http://www.monitoringthefuture.org>
- Newcomb, M.D., Maddahian, E., and Skager, R. (1987). *Substance abuse and psychosocial risk factors among teenagers: Associations with sex, age, ethnicity, and type of school*. *American Journal of Drug and Alcohol Abuse*, 13, 413–433.
- Newman, I., et al. (1989). *The influence of parental attitude and behavior on early adolescent cigarette smoking*. *Journal of School Health* 59(4), 150-152.
- Office of National Drug Control Policy. (2007). *Teens, drugs violence: A special report*. Accessed from: <https://www.hsd.org/?view&did=477440>, May 2015.
- Pierce, J.P., Gilpin, E.A., Farkas, A.J., and Merritt, R.K. (1996). *Validation of susceptibility as a predictor of which adolescents take up smoking in the United States*. *Health Psychology*, 15(5), 355–361.
- Resnick, M., Bearman, P.S., Blum, R.W., Bauman, K.E., Harris, K.K., Jones, J., et al. (1997). *Protecting adolescents from harm: Findings from the National Longitudinal Study on Adolescent Health*. *Journal of the American Medical Association*, 278(10), 823–832.

- Rutter, M. (1979). *Protective factors in children's responses to stress and disadvantage*. In M.W. Kent and J.E. Rolf (Eds.), *Primary Prevention of Psychopathology, Vol. 3. Social competence in children*. Hanover, NH: University Press of New England, 49–74.
- Sabel, J., Bensley, L., and VanEenwyk, J. (2004). *Associations between adolescent drinking and driving involvement and self-reported risk and protective factors in students in public schools in Washington State*. *Journal of Studies on Alcohol*, 65, 213–216.
- Skopp G, Richter B, Pötsch L. *Serum cannabinoid levels 24 to 48 hours after cannabis smoking*. *Arch Kriminol*. 2003 Sep-Oct;212(3-4):83-95. German. PubMed PMID: 14639811.
- Smith P.K., Pepler, D., Rigby, K. (2004). *Bullying in Schools: How Successful Can Interventions Be?* Cambridge University Press, 2004. Serum cannabinoid levels 24 to 48 hours after cannabis smoking. *Arch Kriminol*. 2003 Sep-Oct;212(3-4):83-95.
- Starr, G., Rogers, T., Schooley, M., Porter, S., Wiesen, E., and Jamison, N. (2005). *Key outcome indicators for evaluation compressive tobacco control programs*. Atlanta, GA: Centers for Disease Control and Prevention, 46.
- Substance Abuse and Mental Health Services Administration. (2009). *Results from the 2008 National Survey on Drug Use and Health: National Findings*. Office of Applied Studies, NSDUH Series H-36, HHS Publication No. SMA 09-4434. Rockville, MD.
- Substance Abuse and Mental Health Services Administration. (2012). *Substance Abuse Prevention and Mental Health Promotion Five Year Strategic Plan*. Center for Substance Use and Prevention. Rockville, MD.
- Task Force on Community Preventive Services. Tobacco. (2005). In : Zaza, S., Briss, P.A., Harris, K.W., (eds). *The Guide to Community Preventive Services: What Works to Promote Health?* Atlanta (GA): Oxford University Press;2005:3-79. The White House. National drug control strategy. Washington DC.
- Tomar, S. (2003). *Is use of smokeless tobacco a risk factor for cigarette smoking? The U.S. experience*. *Nicotine & Tobacco Research*, 5(4), 561-569.
- U.S. Department of Education, Office of Elementary and Secondary Education. (2001). *No Child Left Behind: A desktop reference*. Washington, DC.
- U.S. Department of Education, Office of Elementary and Secondary Education. (2002). *High School Graduation Initiative, also known as School Dropout Prevention Program*. Washington, DC.
- U.S. Department of Health and Human Services. (1994). *Preventing tobacco use among young people: A report of the Surgeon General*. Atlanta, GA: National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- U.S. Department of Health and Human Services. (2010). *Healthy People 2020*. Retrieved from <http://www.healthypeople.gov/2020/default.aspx> , December 3, 2012.
- U.S. Department of Health and Human Services. (2015). *Dietary Guidelines*. Retrieved from <http://www.health.gov/dietaryguidelines>, June 22, 2015.
- U.S. Department of Health and Human Services.(2006c). *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Children are Hurt by Secondhand Smoke*. Retrieved from: www.surgeongeneral.gov/library/secondhandsmoke/factsheets/factsheet2.html

- U.S. Department of Health and Human Services. (2008). *Physical Activity Guidelines for Americans*. Retrieved from: <http://www.health.gov/PAGuidelines/>
- Washington State Board of Health. (2009). *2009 Washington State Board of Health Strategic Plan*. Olympia, WA.
- Wasserman, G.A., Keenan, K., Tremblay, R., Coie, J.D., Merrenkohl, T.I., Loeber, R. and Petechuk, D. (2003). *Risk and protective factors of child delinquency*. Child Delinquency Bulletin. Retrieved from: <http://www.ncjrs.org/html/ojjdp/193409/contents.html>
- Weinreb, L., et al. (2002). *Hunger: its impact on children's health and mental health*. Pediatrics, 110(4); 41.
- Werner, E. and Smith, R. (1989). *Vulnerable but invincible: A longitudinal study of resilient children and youth*. New York: Adams, Bannister, and Cox.
- World Health Organization. (2007). *Smokeless Tobacco and Some Tobacco-Specific N-Nitrosamines*. International Agency for Research on Cancer Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France. Vol. 89.



Healthy Youth Survey 2014

Report of Results

Statewide Results
Grades 6, 8, 10 and 12

April 23, 2015

Looking Glass Analytics
215 Legion Way SW
Olympia, WA 98502

More information about the Healthy Youth Survey is available at www.AskHYS.net
(Note: this is an updated version of this report. Please see Errata at www.askhys.net/errata for details)

Report Contents

Introduction and Overview	2
Survey Participation in this Report	2
Caution about Participation Rates and Bias	2
Small Numbers: Caution about Number of Students Participating	2
Key to the Notes	2
Highlights of the Local Results	3
Understanding Your Report	4
N's	4
Confidence Intervals	5
More Information	5
Individual Question Results	6
Demographics and General Information (Questions 1–15)	6
Alcohol, Tobacco, and Other Drug (ATOD) Use	8
Lifetime Use (Questions 16-24)	8
30-Day Use (Questions 25-37)	9
Other Tobacco-Related Questions (Questions 38–42)	11
Other Alcohol-Related Questions (Questions 43–54)	12
Other Marijuana-Related Questions (Questions 55–60)	14
Other Alcohol- and Drug-Related Questions (Questions 61-65)	15
Other Health Concerns	16
Nutrition and Fitness (Questions 66–87)	16
Health Conditions and Health Care (Questions 88–99)	20
Sexual Behavior (Questions 100-104)	22
Behaviors Related to Unintentional Injuries (Questions 105–115)	23
Behaviors Related to Intentional Injuries (Questions 116–121)	25
Physical, Emotional and Sexual Abuse (Questions 122–126)	26
Mental Health (Questions 127–136)	27
Social and Emotional Learning (Questions 137–141)	28
Gambling (Question 142)	28
School Climate (Questions 143-156)	29
Quality of Life (Questions 157–163)	31
Risk and Protective Factors	33
Risk and Protective Factor: Reporting Schedule	33
Risk and Protective Factor: Domain Scale Results and Graphs	34
Risk and Protective Factor: Individual Question Results	38
Community Domain (Questions 164–180)	38
Family Domain (Questions 181–198)	40
School Domain (Questions 199–216)	43
Peer and Individual Domain (Questions 217–249)	46
Questions by Topic	52
Core Questions	53

For a detailed list of all of the survey questions by topic - please use the Questions by Topic section in the back of this report.

Statewide Results

Grades 6, 8, 10 and 12

Introduction and Overview

This report presents results from the fall 2014 Healthy Youth Survey in Washington State. This survey was sponsored by the Department of Health, the Office of Superintendent of Public Instruction, the Department of Social and Health Services and the Liquor Control Board in cooperation with schools throughout the state of Washington.

This Multiple Grade Report does not provide comparisons to the state. To compare your results to the state sample, see the Multiple Grade State Sample Report for grades 6, 8, 10 and 12. This report are available at: <http://www.askhys.net/Reports> under State Reports.

Survey questions covering the same topics are grouped together when possible but be sure to consult the index for related questions when you are searching for information on a specific topic. The numbering in this report is not the same as the question order on the survey itself.

Additional information may be found in the Interpretive Guide posted to the Healthy Youth Survey web site: www.AskHYS.net/Reports

Survey Participation in this Report

	<u>Grade 6</u>	<u>Grade 8</u>	<u>Grade 10</u>	<u>Grade 12</u>
Number of students surveyed:	9,301	11,451	9,201	6,907
Number of valid responses:	9,129	10,673	8,821	6,639
Number of enrolled students*:	11,511	13,556	13,269	13,239
Your survey participation rate**:	79%	79%	66%	50%

* The estimate of enrolled students is based on figures from the 2014-2015 school year, provided by OSPI.

** The survey participation rate is the number of valid responses divided by the number of enrolled students.

Caution about Participation Rates and Bias

Please use the following guidance when reviewing your results:

- 70% or greater participation—Results are probably representative of students in this grade.
- 40–69% participation—Results may be representative of students in this grade.
- Less than 40% participation—Results are likely not representative of students in this grade but do reflect students who completed the survey.

There may be limitations to your results even if you have a high participation rate. For instance, a particular group of students (say, the school orchestra) may have been away from school the day of the survey, and that could bias the results. It is important to acknowledge the potential limitations when using the results in this report. For reports summarizing results at the county or School District level, you should also consider whether the schools that participated represent all students in that area.

Small Numbers: Caution about Number of Students Participating

Results based on small numbers of students answering a question are unstable---that is, they could easily change with the absence from school of only a couple of students. This is especially the case when only a few students choose a particular answer option. Also, in this situation, the reported 95% confidence interval might be too narrow. Thus, use caution if fewer than 30 students answered a question and fewer than 5 students selected a given response option.

For example, if 20 students answered a question and of those 20 only 3 students answered “Yes”, the estimate is unreliable.

See the Understanding Your Report section later in this report for a discussion of “confidence intervals” to help guide your interpretation of the results.

Key to the Notes

Five versions of the survey were used in the administration of the 2014 HYS. Forms A and B were given to secondary students (grades 8, 10, and 12). Some questions on the Forms were available on enhanced versions of the survey and used at the discretion of the school. Form A-enhanced included one optional question on sexual orientation and Form B-enhanced included six optional questions on sexual orientation, behavior and abuse. Form C was given to elementary students (grade 6).

A list of the topics asked and their location in the report are provided in the Questions by Topic section at the end of this report. The Questions by Form Type section provides a list of which Form each question was on.

The following notes are used throughout this report to document the differences between the questions on different versions and indicate the optional questions:


A = wording on Form A; B = wording on Form B; C = wording on Form C

† = additional question on the enhanced survey forms


Highlights of the Local Results

Your students reported the following behaviors and attitudes:

Substance Use	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI)	% (± CI)	% (± CI)	% (± CI)
Smoked cigarettes in past 30 days	1.1% (±0.3)	4.0% (±0.6)	7.9% (±1.1)	13.0% (±1.6)
Drank alcohol in past 30 days	2.1% (±0.4)	8.1% (±1.0)	20.6% (±1.6)	32.9% (±2.6)
Used marijuana or hashish in past 30 days	1.3% (±0.4)	7.3% (±1.1)	18.1% (±1.7)	26.7% (±2.2)
Binge drinking in past 2 weeks	2.3% (±0.5)	4.5% (±0.7)	10.6% (±1.1)	19.2% (±2.0)



Bullying and School Climate	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI)	% (± CI)	% (± CI)	% (± CI)
Carried a weapon at school in the past 30 days	3.1% (±0.3)	3.8% (±0.6)	5.7% (±0.7)	6.4% (±1.1)
Bullied in the past 30 days	30.9% (±1.9)	28.0% (±1.8)	22.6% (±1.3)	16.4% (±1.3)
Enjoyed being at school over the past year	58.3% (±1.8)	46.4% (±1.5)	37.5% (±2.0)	36.5% (±2.5)
Felt safe at school	88.8% (±1.1)	86.0% (±1.7)	84.6% (±2.5)	87.0% (±2.3)




For more information on specific HYS topics, see the Fact Sheets at www.AskHYS.net, under HYS Results – Fact Sheets

Understanding Your Report

N's

In this report “n” is shorthand for “number”---the number of students who took this survey or who answered a survey question. N's are important! We also report the point estimates (%), and the 95% confidence intervals (CI).

Here is an example of how results are displayed in the report tables and charts.



How many students took this survey?

Look on page 2, under Survey Participation. The “number of students surveyed” is the number of surveys that we received. We have careful procedures for “cleaning” to remove surveys with many inconsistent or false answers – the “clean” surveys are the “number of valid surveys”. Only valid surveys are presented in this report.

How many students answered each survey question?

The n's for each question vary. Why? There are several reasons:

- There are four versions of the grades 8, 10, and 12 survey – A, A-enhanced, B, and B-enhanced. Core questions are on all versions, so the n's for those questions are about the same as the number of valid surveys. For non-Core questions the n's will be about half, or less.
- Some versions of the survey had additional questions about sexual orientation, behavior and abuse. Schools that chose to not ask those additional questions will have no responses for those questions. [District and county reports may include some schools that DID ask those questions and some that DID NOT.]
- Not all students finish the survey, so questions near the end have smaller n's. [NOTE: the order of the questions in the report is not the same as on the survey itself.] Also, some students may decide to skip certain questions.

Here are examples from a district with 400 valid surveys --- notice how the n's change with each question type.

17. Have you ever, even once in your life: Used marijuana?	(n=398)	(n=4,000)	N for a core question (almost all students)
a. Yes	49.0% (±3.0)	46.0% (±1.0)	
b. No	51.0% (±3.0)	54.0% (±1.0)	
18. Have you ever, even once in your life: Used inhalants (things you sniff to get high)?	(n=200)	(n=2,000)	N for a question on only one version (about half the students)
a. Yes	11.0% (±4.0)	12.0% (±2.0)	
b. No	89.0% (±4.0)	88.0% (±2.0)	
100. How old were you when you had sexual intercourse for the first time? †	(n=100)	(n=1,500)	N for an additional sexual behavior question (about half, or less if some schools requested surveys without the questions)
a. I have never had sexual intercourse.	70.0% (±8.0)	75.0% (±4.0)	
b. 11 years old or younger	2.0% (±2.0)	2.0% (±1.0)	
c. 12 years old	2.0% (±2.0)	1.0% (±1.0)	
d. 13 years old	5.0% (±2.0)	4.0% (±1.0)	
e. 14 years old	8.0% (±6.0)	7.0% (±2.0)	
f. 15 years old	9.0% (±5.0)	8.0% (±2.0)	
g. 16 years old	3.0% (±3.0)	2.0% (±1.0)	
h. 17 years old or older	1.0% (±1.0)	1.0% (±1.0)	
150. Does your school have a counselor	(n=375)	(n=3,750)	N for a core question near the end of the survey (almost all, except students who don't finish the survey)
a. Yes	90.0% (±3.0)	95.0% (±1.0)	
b. No	5.0% (±2.0)	2.0% (±1.0)	
c. Not sure	5.0% (±2.0)	3.0% (±1.0)	

How many students are we talking about?

N's are only provided for the total number of students who answered a question, not the individual answer options. Some people find it useful to translate their point estimates (%) into the number of students.

For example, if 200 students are enrolled in the district and 10% ($\pm 2\%$) reported "Yes" they used inhalants in their lifetime, then an estimated 20 (± 4) students in the school use inhalants ($200 \times .10 = 20$; and $200 \times .02 = 4$). Enrollment numbers are available on page 2 under Survey Participation.

Confidence Intervals

It is unlikely that the point estimate (%) reported for each question is *exactly* the same as the "true" value for all your students. This report includes 95% confidence intervals (CI) to describe this uncertainty. If there is no bias in the data, then there is a 95% chance that the CI will include the true value. So for example, if the point estimate and CI are 51.5% ($\pm 1.5\%$), then there is a 95% chance that the true percentage is between 50.0% and 53.0%.

The size of your CIs depend in part on the n for each question – as more students answer a question, your point estimate is likely to be closer to the "true" value. If you are a small school, district or county your CIs will be large. The confidence interval does not take into account the participation rate – in other words, a low participation rate can create bias which will not be reflected in the CI.

How can you "use" confidence intervals?

CIs help you to decide if the differences between your students at different grade levels are statistically significant.

For an example, let's say your 8th Grade students report 25% $\pm 5\%$. That means that the CI is between 20% and 30% Here is the math: $25\% - 5\% = 20\%$, and $25\% + 5\% = 30\%$

Using CIs, you can decide if the difference between your 8th grade and 10th grade students are statistically significant:

1. A significant difference:

- 8th Grade students report 25% $\pm 5\%$, so the point estimate is 25% and the CI is 20% to 30%
- 10th Grade students report 36% $\pm 3\%$, so the point estimate is 36% and the CI is 33% to 39%
- The CIs don't overlap, so the difference is significant



2. Not a significant difference:

- 8th Grade students report 25% $\pm 5\%$, so the point estimate is 25% and the CI is 20% to 30%
- 10th Grade students report 28% $\pm 3\%$, so the point estimate is 28% and the CI is 25% to 31%
- The CI for your 8th Grade students overlaps the point estimate for 10th Grade students, so the difference is not significant



3. Not sure if there is a difference:

- 8th Grade students report 25% $\pm 5\%$, so the point estimate is 25% and the CI is 20% to 30%
- 10th Grade students report 32% $\pm 3\%$, so the point estimate is 32% and the CI is 29% to 35%
- The CIs overlap, but don't include the other point estimates, so you don't know for sure if they are really different without using the significance testing tool described in the next paragraph.



More information about interpreting CIs is available on the HYS Training web page at www.AskHYS.net/Training. The Training web page includes past HYS trainings materials and a "Tool for Determining HYS Statistical Significance". Use the "tool" to find out if your results are significantly different.

More Information

Much of the information on the Healthy Youth Survey is available online at www.AskHYS.net.

- Under HYS Results, you can find these "Frequency Reports". Also included is the 2012 Interpretive Guide with a detailed overview of the reports, information on statistical issues and how to use your data. There are also topical fact sheets and a data query system where you can run your own simple analysis.
- Under Training, there are copies of past HYS training workshop materials and the tool for testing significant differences.

Technical information is also at www.doh.wa.gov/DataandStatisticalReports/HealthBehaviors/HealthyYouthSurvey

Frequency Results

Demographics and General Information

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,670)	% (± CI) (n=8,817)	% (± CI) (n=6,638)
1. How old are you?				
a. 12 or younger	**	1.1% (±0.2)	0.1% (±0.1)	0.1% (±0.1)
b. 13	**	74.8% (±1.4)	0.1% (±0.1)	0.0% (±0.0)
c. 14	**	23.4% (±1.4)	1.4% (±0.3)	0.0% (±0.0)
d. 15	**	0.7% (±0.2)	73.7% (±1.6)	0.0% (±0.0)
e. 16	**	0.0% (±0.0)	24.0% (±1.6)	1.4% (±0.3)
f. 17	**	0.0% (±0.0)	0.7% (±0.2)	72.7% (±1.7)
g. 18	**	0.0% (±0.0)	0.1% (±0.0)	24.1% (±1.5)
h. 19 or older	**	0.0% (±0.0)	0.0% (±0.0)	1.6% (±0.6)
2. How old are you?				
	(n=9,126)	(n=0)	(n=0)	(n=0)
a. 10 or younger	1.5% (±0.3)	**	**	**
b. 11	76.0% (±1.2)	**	**	**
c. 12	21.8% (±1.2)	**	**	**
d. 13	0.6% (±0.2)	**	**	**
e. 14	0.0% (±0.0)	**	**	**
f. 15 or older	0.1% (±0.1)	**	**	**
3. Are you:				
	(n=9,118)	(n=10,653)	(n=8,805)	(n=6,626)
a. Female	50.6% (±1.1)	50.3% (±1.0)	51.6% (±1.0)	49.8% (±1.4)
b. Male	49.4% (±1.1)	49.7% (±1.0)	48.4% (±1.0)	50.2% (±1.4)
4. How do you describe yourself? (Select one or more responses.)				
	(n=8,696)	(n=10,544)	(n=8,782)	(n=6,631)
a. American Indian or Alaskan Native	5.9% (±1.1)	3.3% (±0.7)	2.4% (±0.6)	2.1% (±0.8)
b. Asian or Asian American	9.7% (±3.4)	9.9% (±3.1)	9.3% (±2.8)	8.6% (±2.8)
c. Black or African-American	4.8% (±1.4)	4.1% (±1.0)	4.9% (±1.5)	4.9% (±1.6)
d. Hispanic or Latino/Latina	12.2% (±3.5)	15.1% (±5.1)	11.8% (±4.7)	12.4% (±5.0)
e. Native Hawaiian or other Pacific Islander	1.6% (±0.4)	1.7% (±0.4)	2.2% (±0.7)	1.9% (±0.8)
f. White or Caucasian	39.1% (±4.7)	49.3% (±5.5)	56.0% (±6.2)	58.8% (±6.5)
g. Other	17.4% (±1.8)	8.0% (±0.7)	5.6% (±0.6)	3.9% (±0.5)
More than one race/ethnicity marked	9.3% (±0.8)	8.7% (±1.1)	7.9% (±1.0)	7.4% (±0.9)
5. What language is usually spoken at home?				
	(n=0)	(n=10,425)	(n=8,598)	(n=6,492)
a. English	**	79.5% (±4.0)	82.0% (±4.1)	82.8% (±4.1)
b. Spanish	**	10.2% (±3.5)	7.4% (±3.2)	7.5% (±3.5)
c. Russian	**	1.0% (±0.3)	1.5% (±0.5)	1.2% (±0.5)
d. Ukrainian	**	0.8% (±0.3)	0.9% (±0.3)	1.0% (±0.3)
e. Vietnamese	**	1.4% (±0.9)	1.5% (±0.7)	1.4% (±0.7)
f. Chinese	**	1.6% (±1.0)	1.1% (±1.0)	1.2% (±0.7)
g. Korean	**	0.5% (±0.2)	0.6% (±0.4)	0.6% (±0.4)
h. Japanese	**	0.4% (±0.1)	0.3% (±0.1)	0.3% (±0.1)
i. Other	**	4.5% (±1.0)	4.6% (±1.1)	3.9% (±1.0)
6. What language is usually spoken at home?				
	(n=9,089)	(n=0)	(n=0)	(n=0)
a. English	80.7% (±4.4)	**	**	**
b. Spanish	10.0% (±2.9)	**	**	**
c. Other	9.3% (±3.1)	**	**	**

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,082)	% (± CI) (n=8,413)	% (± CI) (n=6,339)
7. How far did your mother get in school?				
a. Did not finish high school	**	9.3% (±1.9)	11.4% (±2.9)	11.6% (±3.2)
b. Graduated from high school or GED	**	15.1% (±1.5)	18.1% (±2.3)	20.2% (±2.3)
c. Had some college or technical training after high school	**	15.7% (±1.2)	21.4% (±2.0)	24.7% (±2.1)
d. Graduated from a 4-year college	**	22.3% (±2.6)	22.9% (±3.6)	23.1% (±3.8)
e. Earned an advanced graduate degree	**	13.8% (±2.4)	12.5% (±3.0)	11.5% (±3.0)
f. Don't know	**	21.9% (±1.7)	11.6% (±1.4)	6.9% (±1.0)
g. Does not apply	**	1.9% (±0.4)	2.1% (±0.4)	2.1% (±0.5)
8. How far did your father get in school?	(n=0)	(n=10,064)	(n=8,397)	(n=6,329)
a. Did not finish high school	**	9.2% (±1.9)	11.2% (±2.6)	12.3% (±3.2)
b. Graduated from high school or GED	**	15.4% (±1.4)	18.1% (±2.4)	19.8% (±2.5)
c. Had some college or technical training after high school	**	12.6% (±1.3)	16.5% (±1.7)	19.9% (±1.8)
d. Graduated from a 4-year college	**	19.6% (±2.6)	20.2% (±3.3)	21.5% (±3.8)
e. Earned an advanced graduate degree	**	15.1% (±3.0)	15.1% (±3.9)	13.5% (±3.3)
f. Don't know	**	25.4% (±2.4)	15.8% (±2.1)	9.7% (±1.5)
g. Does not apply	**	2.7% (±0.6)	3.2% (±0.6)	3.2% (±0.7)
9. Which of the following best describes you? †	(n=0)	(n=1,649)	(n=1,728)	(n=1,284)
a. Heterosexual (straight)	**	87.1% (±2.0)	87.6% (±1.6)	88.2% (±2.4)
b. Gay or lesbian	**	1.0% (±0.6)	1.5% (±0.7)	2.2% (±0.8)
c. Bisexual	**	4.6% (±1.0)	6.3% (±1.0)	6.2% (±1.7)
d. Not sure	**	7.3% (±1.1)	4.6% (±1.0)	3.4% (±1.0)
10. Who did you live with most of the time in the last 30 days?	(n=0)	(n=10,388)	(n=8,584)	(n=6,486)
a. Parent(s) and/or step-parent(s)	**	94.5% (±0.7)	93.5% (±0.8)	91.0% (±1.4)
b. Relatives—like a grandparent, an aunt, an older brother—but NOT your parents	**	3.0% (±0.5)	3.5% (±0.6)	4.2% (±0.7)
c. Foster care parent(s)	**	0.4% (±0.1)	0.6% (±0.2)	0.7% (±0.2)
d. An adult friend(s) of your family	**	0.2% (±0.1)	0.6% (±0.1)	1.0% (±0.3)
e. Friends of yours with no adults present	**	0.4% (±0.1)	0.5% (±0.2)	0.8% (±0.2)
f. On your own	**	0.5% (±0.1)	0.4% (±0.1)	0.9% (±0.2)
g. Other	**	1.0% (±0.2)	0.9% (±0.2)	1.4% (±0.4)
11. Where did you live most of the time in the last 30 days?	(n=0)	(n=10,422)	(n=8,606)	(n=6,504)
a. In a house, apartment, or mobile home	**	98.1% (±0.3)	97.5% (±0.4)	97.3% (±0.6)
b. In a motel or hotel	**	0.3% (±0.1)	0.4% (±0.2)	0.6% (±0.2)
c. In a group home	**	0.2% (±0.1)	0.4% (±0.1)	0.5% (±0.2)
d. In a shelter	**	0.2% (±0.1)	0.3% (±0.1)	0.4% (±0.2)
e. In a car, park, or campground	**	0.1% (±0.1)	0.3% (±0.1)	0.2% (±0.1)
f. On the street	**	0.3% (±0.1)	0.3% (±0.1)	0.3% (±0.2)
g. Moved from place to place	**	0.4% (±0.1)	0.4% (±0.2)	0.3% (±0.2)
h. Other	**	0.4% (±0.1)	0.5% (±0.1)	0.4% (±0.1)
12. Are your current living arrangements the result of losing your home because your family cannot afford housing?	(n=0)	(n=10,243)	(n=8,513)	(n=6,439)
a. No	**	87.2% (±1.2)	88.5% (±1.1)	90.7% (±1.1)
b. Yes	**	4.4% (±0.4)	5.2% (±0.5)	5.4% (±0.6)
c. Not sure	**	8.3% (±1.0)	6.2% (±0.9)	3.9% (±0.6)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=9,014)	% (± CI) (n=5,004)	% (± CI) (n=4,203)	% (± CI) (n=3,180)
13. Has your parent or guardian served in the military (Army, Navy, Air Force, Marines, Coast Guard, National Guard, and Reserves)?				
a. No	65.5% (±1.9)	72.7% (±2.4)	74.2% (±3.3)	76.6% (±3.5)
b. Yes	22.7% (±2.0)	21.3% (±2.4)	21.9% (±3.5)	20.6% (±3.3)
c. Not sure	11.8% (±1.1)	5.9% (±0.8)	3.8% (±0.7)	2.8% (±0.7)
14. Has your military parent or guardian been sent to Iraq, Afghanistan, or other combat zone?	(n=0)	(n=4,993)	(n=4,196)	(n=3,157)
a. I do not have a parent or guardian who has ever served in the military.	**	70.6% (±2.6)	70.1% (±3.2)	71.0% (±3.3)
b. No	**	13.1% (±1.0)	15.7% (±1.6)	16.9% (±1.5)
c. Yes	**	7.3% (±1.4)	8.2% (±2.0)	7.7% (±1.8)
d. Not sure	**	9.0% (±1.1)	6.0% (±0.8)	4.4% (±1.1)
15. How honest were you in filling out this survey?	(n=7,989)	(n=8,656)	(n=7,591)	(n=5,867)
a. I was very honest.	85.6% (±1.3)	82.0% (±1.2)	83.0% (±1.4)	84.0% (±1.8)
b. I was honest pretty much of the time.	13.0% (±1.2)	15.6% (±1.1)	14.5% (±1.1)	13.4% (±1.4)
c. I was honest some of the time	1.5% (±0.4)	2.4% (±0.5)	2.5% (±0.4)	2.6% (±0.6)
d. I was honest once in a while.		surveys pulled		
e. I was not honest at all.		surveys pulled		

Alcohol, Tobacco and Other Drug Use

Alcohol, tobacco, and other drug use has been a major concern in this country for many years. The consequences of ATOD use are well known. In the short term, ATOD use interferes with positive physical, emotional, and social development. In the long term, ATOD use is associated with delinquency and criminal activity, unintended injuries, and a variety of health complications including shorter life expectancy. Tobacco use is the world's leading cause of preventable death, disease, and disability. This section provides information about lifetime ATOD use (which in part reflects experimental use), use in the past 30 days (i.e., current use), and other tobacco-, alcohol-, and drug-related issues.

Lifetime Use

Have you ever, even once in your life:

16. Smoked a cigarette, even just a puff? (Computed from question 222)	(n=0)	(n=5,068)	(n=4,144)	(n=3,119)
a. No	**	88.2% (±1.7)	78.0% (±2.6)	68.5% (±3.1)
b. Yes	**	11.8% (±1.7)	22.0% (±2.6)	31.5% (±3.1)
17. Drank more than a sip or two of beer, wine, or hard liquor (for example: vodka, whiskey, or gin)? (Computed from question 17 or 223)	(n=8,650)	(n=10,348)	(n=8,579)	(n=6,423)
a. No	78.8% (±1.3)	71.0% (±1.9)	49.9% (±2.1)	33.8% (±2.2)
b. Yes	21.2% (±1.3)	29.0% (±1.9)	50.1% (±2.1)	66.2% (±2.2)
18. Used marijuana? (Computed from question 18 or 221)	(n=8,667)	(n=10,348)	(n=8,563)	(n=6,423)
a. No	96.9% (±0.5)	89.6% (±1.6)	70.6% (±2.5)	54.3% (±3.2)
b. Yes	3.1% (±0.5)	10.4% (±1.6)	29.4% (±2.5)	45.7% (±3.2)
19. Used inhalants (things you sniff to get high)?	(n=8,662)	(n=4,969)	(n=4,095)	(n=3,088)
a. No	97.7% (±0.4)	95.5% (±0.7)	92.4% (±0.9)	92.0% (±1.2)
b. Yes	2.3% (±0.4)	4.5% (±0.7)	7.6% (±0.9)	8.0% (±1.2)
20. Used heroin?	(n=0)	(n=4,991)	(n=4,098)	(n=3,091)
a. No	**	97.4% (±0.5)	96.6% (±0.7)	96.8% (±0.8)
b. Yes	**	2.6% (±0.5)	3.4% (±0.7)	3.2% (±0.8)

	Grade 6	Grade 8	Grade 10	Grade 12
21. Used methamphetamines (meth, crystal meth, ice, crank)? Do not include other types of amphetamines.	% (± CI) (n=0)	% (± CI) (n=4,987)	% (± CI) (n=4,094)	% (± CI) (n=3,090)
a. No	**	97.5% (±0.5)	95.9% (±0.8)	96.2% (±0.9)
b. Yes	**	2.5% (±0.5)	4.1% (±0.8)	3.8% (±0.9)
22. Used cocaine?	(n=0)	(n=4,979)	(n=4,096)	(n=3,088)
a. No	**	97.1% (±0.6)	95.8% (±0.8)	93.5% (±1.0)
b. Yes	**	2.9% (±0.6)	4.2% (±0.8)	6.5% (±1.0)
23. Used steroids (muscle builders) without a doctor's prescription?	(n=0)	(n=4,974)	(n=4,100)	(n=3,083)
a. No	**	97.7% (±0.5)	96.8% (±0.6)	96.8% (±0.8)
b. Yes	**	2.3% (±0.5)	3.2% (±0.6)	3.2% (±0.8)
24. Have you ever, even once in your lifetime: Used other illegal drugs?	(n=8,654)	(n=0)	(n=0)	(n=0)
a. Yes	2.4% (±0.4)	**	**	**
b. No	97.6% (±0.4)	**	**	**

30-Day Use (Use in the Past 30 Days)

During the past 30 days, on how many days did you:

25. Smoke cigarettes?	(n=8,790)	(n=10,437)	(n=8,619)	(n=6,491)
a. None	98.9% (±0.3)	96.0% (±0.6)	92.1% (±1.1)	87.0% (±1.6)
b. 1-2 days	0.5% (±0.2)	2.0% (±0.3)	2.8% (±0.4)	4.1% (±0.7)
c. 3-5 days	0.2% (±0.1)	0.5% (±0.1)	1.4% (±0.3)	2.3% (±0.4)
d. 6-9 days	0.2% (±0.1)	0.5% (±0.2)	1.0% (±0.2)	1.6% (±0.4)
e. 10-29 days	0.1% (±0.1)	0.5% (±0.1)	1.2% (±0.3)	2.5% (±0.5)
f. All 30 days	0.2% (±0.1)	0.5% (±0.2)	1.5% (±0.4)	2.5% (±0.7)
<i>Any use in past 30 days</i>	1.1% (±0.3)	4.0% (±0.6)	7.9% (±1.1)	13.0% (±1.6)
26. Use chewing tobacco, snuff, or dip?	(n=8,780)	(n=5,297)	(n=4,447)	(n=3,330)
a. None	98.8% (±0.3)	98.7% (±0.4)	96.3% (±0.6)	94.9% (±1.1)
b. 1-2 days	0.6% (±0.2)	0.5% (±0.2)	1.7% (±0.4)	2.0% (±0.6)
c. 3-5 days	0.1% (±0.1)	0.2% (±0.1)	0.4% (±0.2)	0.7% (±0.3)
d. 6-9 days	0.2% (±0.1)	0.1% (±0.1)	0.4% (±0.2)	0.5% (±0.3)
e. 10-29 days	0.2% (±0.1)	0.2% (±0.1)	0.5% (±0.2)	0.8% (±0.4)
f. All 30 days	0.1% (±0.1)	0.2% (±0.1)	0.6% (±0.3)	1.1% (±0.4)
<i>Any use in past 30 days</i>	1.2% (±0.3)	1.3% (±0.4)	3.7% (±0.6)	5.1% (±1.1)
27. Smoke cigars, cigarillos, or little cigars?	(n=0)	(n=5,293)	(n=4,445)	(n=3,325)
a. None	**	98.1% (±0.5)	94.9% (±0.7)	89.8% (±1.4)
b. 1-2 days	**	1.0% (±0.4)	2.8% (±0.5)	6.2% (±1.1)
c. 3-9 days	**	0.5% (±0.2)	1.2% (±0.3)	2.6% (±0.7)
d. 10-29 days	**	0.2% (±0.1)	0.7% (±0.3)	0.8% (±0.3)
e. All 30 days	**	0.2% (±0.1)	0.5% (±0.2)	0.5% (±0.3)
<i>Any use in past 30 days</i>	**	1.9% (±0.5)	5.1% (±0.7)	10.2% (±1.4)

	Grade 6	Grade 8	Grade 10	Grade 12
28. Use an electronic cigarette, also called e-cigs, or vape pens?	% (± CI) (n=0)	% (± CI) (n=5,288)	% (± CI) (n=4,440)	% (± CI) (n=3,320)
a. None	**	91.5% (±1.3)	82.0% (±1.6)	76.9% (±2.2)
b. 1-2 days	**	4.3% (±0.7)	8.1% (±1.0)	10.0% (±1.1)
c. 3-9 days	**	2.2% (±0.5)	4.9% (±0.7)	7.1% (±1.0)
d. 10-29 days	**	1.0% (±0.3)	2.9% (±0.6)	3.1% (±0.9)
e. All 30 days	**	1.0% (±0.3)	2.0% (±0.6)	2.9% (±0.7)
<i>Any use in past 30 days</i>	**	8.5% (±1.3)	18.0% (±1.6)	23.1% (±2.2)

	Grade 6	Grade 8	Grade 10	Grade 12
29. Smoke tobacco or flavored tobacco in a hookah, even just a puff?	(n=0)	(n=5,288)	(n=4,443)	(n=3,320)
a. None	**	95.3% (±0.8)	90.0% (±1.3)	85.2% (±1.7)
b. 1-2 days	**	2.5% (±0.5)	5.1% (±0.8)	8.5% (±1.1)
c. 3-9 days	**	1.0% (±0.3)	2.8% (±0.6)	3.8% (±0.8)
d. 10-29 days	**	0.5% (±0.2)	1.1% (±0.3)	1.6% (±0.5)
e. All 30 days	**	0.6% (±0.2)	0.9% (±0.3)	1.0% (±0.4)
<i>Any use in past 30 days</i>	**	4.7% (±0.8)	10.0% (±1.3)	14.8% (±1.7)

	Grade 6	Grade 8	Grade 10	Grade 12
30. Not including menthols – during the past 30 days on how many days did you use tobacco that tastes like candy, fruit or alcohol (tobacco includes: little cigars, bidis, cloves, chew, spit, snus, hookah)?	(n=0)	(n=5,282)	(n=4,444)	(n=3,326)
a. None	**	95.8% (±0.8)	90.9% (±1.0)	84.9% (±1.6)
b. 1-2 days	**	2.0% (±0.4)	4.3% (±0.5)	8.0% (±1.0)
c. 3-9 days	**	1.2% (±0.3)	2.8% (±0.5)	4.1% (±0.8)
d. 10-29 days	**	0.6% (±0.2)	1.5% (±0.3)	2.0% (±0.5)
e. All 30 days	**	0.4% (±0.2)	0.5% (±0.2)	1.0% (±0.4)
<i>Any use in past 30 days</i>	**	4.2% (±0.8)	9.1% (±1.0)	15.1% (±1.6)

	Grade 6	Grade 8	Grade 10	Grade 12
31. Are the cigarettes that you usually smoke menthol cigarettes?	(n=0)	(n=5,293)	(n=4,441)	(n=3,325)
a. I do not smoke cigarettes.	**	95.2% (±0.8)	91.8% (±1.2)	87.1% (±1.7)
b. Yes	**	1.3% (±0.3)	3.2% (±0.7)	4.5% (±0.7)
c. No	**	3.5% (±0.6)	5.0% (±0.7)	8.4% (±1.4)

During the past 30 days, on how many days did you:

	Grade 6	Grade 8	Grade 10	Grade 12
32. Drink a glass, can or bottle of alcohol (beer, wine, wine coolers, hard liquor)?	(n=8,754)	(n=10,409)	(n=8,604)	(n=6,468)
a. None	97.9% (±0.4)	91.9% (±1.0)	79.4% (±1.6)	67.1% (±2.6)
b. 1-2 days	1.5% (±0.3)	5.5% (±0.6)	12.9% (±1.1)	18.2% (±1.3)
c. 3-5 days	0.3% (±0.1)	1.3% (±0.3)	4.3% (±0.6)	8.5% (±1.2)
d. 6-9 days	0.1% (±0.1)	0.5% (±0.2)	1.7% (±0.3)	3.3% (±0.6)
e. 10 or more days	0.1% (±0.1)	0.7% (±0.2)	1.7% (±0.3)	2.8% (±0.6)
<i>Any use in past 30 days</i>	2.1% (±0.4)	8.1% (±1.0)	20.6% (±1.6)	32.9% (±2.6)

	Grade 6	Grade 8	Grade 10	Grade 12
33. Use marijuana or hashish (weed, hash, pot)?	(n=8,713)	(n=10,375)	(n=8,579)	(n=6,455)
a. None	98.7% (±0.4)	92.7% (±1.1)	81.9% (±1.7)	73.3% (±2.2)
b. 1-2 days	0.7% (±0.2)	2.8% (±0.5)	6.7% (±0.7)	8.5% (±1.0)
c. 3-5 days	0.2% (±0.1)	1.5% (±0.3)	3.6% (±0.5)	5.3% (±0.6)
d. 6-9 days	0.1% (±0.1)	0.7% (±0.2)	2.0% (±0.4)	2.6% (±0.5)
e. 10 or more days	0.3% (±0.1)	2.3% (±0.5)	5.9% (±0.9)	10.3% (±1.3)
<i>Any use in past 30 days</i>	1.3% (±0.4)	7.3% (±1.1)	18.1% (±1.7)	26.7% (±2.2)

	Grade 6	Grade 8	Grade 10	Grade 12
34. Not counting alcohol, tobacco, or marijuana, use another illegal drug?	% (± CI) (n=8,707)	% (± CI) (n=10,365)	% (± CI) (n=8,572)	% (± CI) (n=6,454)
a. None	99.4% (±0.2)	98.1% (±0.3)	95.6% (±0.5)	93.4% (±1.0)
b. 1-2 days	0.3% (±0.1)	1.0% (±0.2)	2.3% (±0.3)	3.5% (±0.6)
c. 3-5 days	0.1% (±0.1)	0.3% (±0.1)	1.0% (±0.2)	1.6% (±0.5)
d. 6-9 days	0.0% (±0.0)	0.3% (±0.1)	0.4% (±0.2)	0.7% (±0.2)
e. 10 or more days	0.2% (±0.1)	0.3% (±0.1)	0.6% (±0.2)	0.8% (±0.3)
<i>Any use in past 30 days</i>	0.6% (±0.2)	1.9% (±0.3)	4.4% (±0.5)	6.6% (±1.0)
35. Use any illegal drug, including marijuana? (Computed from questions 33 and 34)	(n=8,675)	(n=10,333)	(n=8,544)	(n=6,442)
None	98.6% (±0.4)	92.4% (±1.1)	81.1% (±1.7)	72.4% (±2.2)
Any use in past 30 days	1.4% (±0.4)	7.6% (±1.1)	18.9% (±1.7)	27.6% (±2.2)
36. Use a pain killer TO GET HIGH, like Vicodin, OxyContin (sometimes called Oxy or OC) or Percocet (sometimes called Percs)?	(n=0)	(n=10,345)	(n=8,565)	(n=6,449)
a. None	**	97.7% (±0.4)	95.4% (±0.6)	94.4% (±0.9)
b. 1-2 days	**	1.3% (±0.3)	2.4% (±0.4)	2.9% (±0.6)
c. 3-5 days	**	0.4% (±0.1)	1.3% (±0.3)	1.5% (±0.4)
d. 6-9 days	**	0.2% (±0.1)	0.5% (±0.2)	0.7% (±0.2)
e. 10 or more days	**	0.3% (±0.1)	0.4% (±0.1)	0.5% (±0.2)
<i>Any use in past 30 days</i>	**	2.3% (±0.4)	4.6% (±0.6)	5.6% (±0.9)
37. Use prescription drugs not prescribed to you?	(n=0)	(n=5,061)	(n=4,129)	(n=3,118)
a. None	**	95.8% (±0.6)	92.4% (±0.9)	91.0% (±1.3)
b. 1-2 days	**	2.8% (±0.4)	5.0% (±0.7)	5.0% (±0.9)
c. 3-5 days	**	0.7% (±0.3)	1.3% (±0.4)	2.5% (±0.6)
d. 6-9 days	**	0.4% (±0.2)	0.5% (±0.2)	1.0% (±0.3)
e. 10 or more days	**	0.3% (±0.1)	0.8% (±0.3)	0.6% (±0.3)
<i>Any use in past 30 days</i>	**	4.2% (±0.6)	7.6% (±0.9)	9.0% (±1.3)

Other Tobacco-Related Questions

38. During the past 7 days, on how many days were you in the same room with someone who was smoking cigarettes?	(n=8,239)	(n=4,803)	(n=4,100)	(n=3,110)
a. 0 days	81.3% (±1.5)	78.1% (±2.2)	72.0% (±2.8)	70.1% (±2.5)
b. 1-2 days	11.2% (±0.9)	12.0% (±1.3)	14.4% (±1.5)	15.8% (±1.3)
c. 3-4 days	2.8% (±0.4)	3.6% (±0.6)	5.3% (±0.9)	6.0% (±0.8)
d. 5-6 days	1.1% (±0.3)	2.0% (±0.5)	2.8% (±0.5)	3.0% (±0.8)
e. 7 days	3.6% (±0.6)	4.3% (±0.8)	5.6% (±1.1)	5.1% (±1.1)
39. During the past 7 days, on how many days did you ride in a car with someone who was smoking cigarettes?	(n=8,206)	(n=4,843)	(n=4,098)	(n=3,114)
a. 0 days	87.3% (±1.5)	85.1% (±2.0)	81.7% (±2.5)	79.0% (±2.6)
b. 1-2 days	6.7% (±0.9)	7.3% (±1.1)	8.7% (±1.3)	11.1% (±1.5)
c. 3-4 days	2.2% (±0.3)	2.5% (±0.6)	3.6% (±0.7)	3.8% (±0.9)
d. 5-6 days	1.0% (±0.3)	2.4% (±0.5)	2.8% (±0.6)	2.7% (±0.8)
e. 7 days	2.7% (±0.5)	2.6% (±0.6)	3.2% (±0.8)	3.3% (±0.9)

	Grade 6	Grade 8	Grade 10	Grade 12
40. Does anyone who lives with you now smoke cigarettes?	% (± CI) (n=0)	% (± CI) (n=4,822)	% (± CI) (n=4,092)	% (± CI) (n=3,104)
a. No	**	76.5% (±2.6)	74.0% (±3.0)	75.4% (±3.3)
b. Yes	**	23.5% (±2.6)	26.0% (±3.0)	24.6% (±3.3)
41. Which of these best describes the rules about smoking inside the house where you live? Smoking is...	(n=0)	(n=4,746)	(n=4,070)	(n=3,094)
a. Never allowed inside my house	**	87.7% (±1.4)	88.3% (±1.6)	88.6% (±2.0)
b. Allowed only at some times or in some places	**	9.0% (±1.1)	8.5% (±1.2)	7.7% (±1.4)
c. Always allowed inside my house	**	3.3% (±0.7)	3.3% (±0.8)	3.7% (±0.9)
42. How wrong do your friends feel it would be for you to: Use tobacco?	(n=0)	(n=4,240)	(n=3,674)	(n=2,862)
a. Very wrong	**	76.0% (±2.2)	59.3% (±2.5)	46.6% (±2.8)
b. Wrong	**	16.1% (±1.4)	23.3% (±1.7)	24.4% (±1.9)
c. A little bit wrong	**	4.7% (±0.8)	10.5% (±1.1)	14.0% (±1.3)
d. Not at all wrong	**	3.2% (±0.8)	6.9% (±1.3)	14.9% (±1.9)

Other Alcohol-Related Questions

43. Think back over the last 2 weeks. How many times have you had five or more drinks in a row? (A drink is a glass of wine, a bottle of beer, a shot glass of liquor, or a mixed drink.)	(n=8,642)	(n=10,326)	(n=8,562)	(n=6,429)
a. None	97.7% (±0.5)	95.5% (±0.7)	89.4% (±1.1)	80.8% (±2.0)
b. Once	1.4% (±0.3)	2.2% (±0.3)	5.0% (±0.6)	8.8% (±1.1)
c. Twice	0.5% (±0.1)	1.1% (±0.3)	2.6% (±0.4)	5.5% (±0.7)
d. 3-5 times	0.2% (±0.1)	0.6% (±0.2)	1.9% (±0.4)	3.2% (±0.6)
e. 6-9 times	0.1% (±0.1)	0.3% (±0.1)	0.6% (±0.1)	0.8% (±0.2)
f. 10 or more times	0.2% (±0.1)	0.4% (±0.1)	0.5% (±0.2)	1.0% (±0.3)
44. Drinking Categories Variable (computed from questions 32 and 43)	(n=8,615)	(n=10,309)	(n=8,555)	(n=6,431)
No drinking past 30 days and no bingeing past 2 weeks	96.4% (±0.6)	90.8% (±1.1)	78.1% (±1.7)	65.5% (±2.6)
1-2 days drinking past 30 days and no bingeing past 2 weeks	1.0% (±0.2)	3.9% (±0.5)	9.2% (±0.9)	11.7% (±0.9)
3-5 days drinking past 30 days and/or 1 binge past 2 weeks	1.5% (±0.4)	2.6% (±0.4)	5.8% (±0.7)	10.8% (±1.3)
6+ days drinking past 30 days and/or 2+ binge past 2 weeks	1.0% (±0.2)	2.8% (±0.5)	6.9% (±0.8)	12.0% (±1.4)
45. During the past 30 days, what type of alcohol did you usually drink?	(n=0)	(n=4,468)	(n=3,806)	(n=2,931)
a. I did not drink alcohol during the past 30 days.	**	91.7% (±1.1)	79.1% (±2.2)	67.0% (±3.1)
b. I do not have a usual type.	**	1.5% (±0.3)	2.8% (±0.5)	4.3% (±0.7)
c. Beer	**	1.4% (±0.4)	4.5% (±1.1)	7.7% (±1.3)
d. Flavored malt beverages, such as Smirnoff Ice, Bacardi Silver, or Hard Lemonade	**	1.2% (±0.4)	2.7% (±0.7)	3.7% (±0.7)
e. Wine coolers, such as Bartles & Jaymes or Seagrams	**	0.2% (±0.2)	0.5% (±0.2)	0.7% (±0.3)
f. Wine	**	0.9% (±0.3)	1.6% (±0.4)	1.7% (±0.6)
g. Liquor, such as vodka, rum, scotch, bourbon or whiskey	**	2.3% (±0.6)	8.1% (±1.1)	13.9% (±1.9)
h. Some other type	**	0.8% (±0.3)	0.8% (±0.3)	1.0% (±0.4)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,511)	% (± CI) (n=3,816)	% (± CI) (n=2,941)
46. During the past 30 days, how did you get alcohol (beer, wine or hard liquor)? Choose all that apply.				
a. I did not get alcohol in the past 30 days.	**	91.5% (±1.1)	80.2% (±1.9)	68.7% (±3.0)
b. I bought it from a store.	**	0.7% (±0.2)	1.5% (±0.4)	2.0% (±0.5)
c. I stole it from a store.	**	0.7% (±0.3)	1.5% (±0.4)	2.2% (±0.6)
d. I got it from friends.	**	2.2% (±0.5)	8.5% (±1.2)	14.4% (±2.1)
e. I got it at a party.	**	1.5% (±0.3)	6.1% (±1.3)	10.8% (±1.7)
f. I got it from an older brother or sister.	**	0.6% (±0.2)	2.0% (±0.4)	3.0% (±0.6)
g. I gave money to someone to get it for me.	**	0.8% (±0.3)	3.5% (±0.6)	9.2% (±1.7)
h. I took it from home without my parents' permission.	**	1.8% (±0.4)	4.8% (±0.9)	3.8% (±0.8)
i. I got it at home with my parents' permission.	**	1.6% (±0.4)	3.3% (±0.7)	5.4% (±0.8)
j. I got it some other way.	**	2.0% (±0.5)	2.5% (±0.5)	4.3% (±0.8)
47. NOT including talks on drinking and driving, in the past year have your parents or guardians talked to you about why you should not drink alcohol?	(n=0)	(n=5,194)	(n=4,245)	(n=3,188)
a. Yes, a number of times	**	42.8% (±2.1)	38.4% (±1.7)	34.0% (±1.6)
b. Yes, once	**	23.0% (±1.4)	21.3% (±1.1)	19.4% (±1.7)
c. No	**	20.9% (±1.6)	28.9% (±1.7)	37.6% (±1.8)
d. I don't remember.	**	13.3% (±0.9)	11.3% (±0.9)	8.9% (±0.9)
48. How wrong do your parents feel it would be for you to: Have one or two drinks of an alcoholic beverage nearly every day?	(n=0)	(n=5,325)	(n=4,340)	(n=3,281)
a. Very wrong	**	89.7% (±1.3)	84.0% (±1.6)	75.5% (±2.1)
b. Wrong	**	7.8% (±1.0)	11.6% (±1.4)	17.1% (±1.5)
c. A little bit wrong	**	1.7% (±0.4)	3.0% (±0.4)	4.9% (±0.9)
d. Not wrong at all	**	0.8% (±0.2)	1.4% (±0.5)	2.5% (±0.5)
49. During the past 30 days, on how many days did you have at least one drink of alcohol on school property?	(n=0)	(n=4,563)	(n=3,848)	(n=2,966)
a. 0 days	**	96.7% (±0.8)	93.9% (±1.0)	93.9% (±1.4)
b. 1-2 days	**	1.2% (±0.4)	3.3% (±0.6)	3.4% (±1.0)
c. 3-9 days	**	1.0% (±0.3)	1.4% (±0.4)	1.5% (±0.5)
d. 10-29 days	**	0.4% (±0.2)	0.6% (±0.3)	0.6% (±0.3)
e. All 30 days	**	0.7% (±0.3)	0.7% (±0.3)	0.6% (±0.3)
50. How wrong do your friends feel it would be for you to: Have one or two drinks of an alcoholic beverage nearly every day?	(n=0)	(n=4,236)	(n=3,670)	(n=2,862)
a. Very wrong	**	71.5% (±2.1)	51.3% (±2.6)	41.4% (±2.4)
b. Wrong	**	18.1% (±1.5)	25.6% (±1.4)	26.7% (±1.6)
c. A little bit wrong	**	6.9% (±1.0)	14.6% (±1.5)	17.8% (±1.5)
d. Not at all wrong	**	3.6% (±0.8)	8.6% (±1.3)	14.0% (±1.8)
51. Think about the students in your school. If you had to guess, how wrong do most students in your grade think it is to drink alcohol regularly?	(n=0)	(n=4,874)	(n=4,027)	(n=3,039)
a. Very wrong	**	30.7% (±2.6)	10.2% (±1.7)	7.0% (±1.2)
b. Wrong	**	41.1% (±1.7)	26.0% (±2.3)	18.1% (±2.0)
c. A little bit wrong	**	23.4% (±2.3)	44.9% (±1.9)	47.3% (±1.8)
d. Not wrong at all	**	4.9% (±0.9)	18.8% (±2.6)	27.6% (±2.7)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,183)	% (± CI) (n=4,216)	% (± CI) (n=3,177)
52. How much do you think people risk harming themselves if they: Have 5 or more drinks of an alcoholic beverage once or twice a week?				
a. No risk	**	4.3% (±0.8)	3.4% (±0.6)	4.4% (±0.9)
b. Slight risk	**	8.0% (±1.0)	8.8% (±1.3)	11.8% (±1.2)
c. Moderate risk	**	24.6% (±1.2)	28.7% (±1.8)	28.9% (±1.6)
d. Great risk	**	57.2% (±2.6)	56.2% (±2.5)	52.3% (±2.2)
e. Not sure	**	6.0% (±1.0)	3.0% (±0.6)	2.5% (±0.6)
53. How do you feel about someone your age having one or two drinks of an alcoholic beverage nearly every day?	(n=0)	(n=4,802)	(n=3,981)	(n=3,027)
a. Neither approve nor disapprove	**	17.0% (±1.7)	21.6% (±1.7)	22.1% (±2.1)
b. Somewhat disapprove	**	12.9% (±1.3)	21.0% (±1.3)	24.2% (±1.7)
c. Strongly disapprove	**	61.2% (±3.2)	50.3% (±2.4)	47.0% (±2.7)
d. Don't know or can't say	**	8.9% (±1.4)	7.2% (±1.2)	6.7% (±1.2)
54. In the past year, which of the following happened because you drank alcohol or used drugs? Choose all that apply.	(n=0)	(n=4,405)	(n=3,759)	(n=2,896)
a. I did not use alcohol or drugs in the past year.	**	88.2% (±1.4)	69.5% (±2.4)	53.1% (±3.0)
b. I did not have any problems from drinking alcohol or using drugs in the past year.	**	8.7% (±1.1)	20.9% (±1.8)	31.0% (±2.3)
c. I missed classes or school.	**	1.1% (±0.3)	1.7% (±0.4)	3.0% (±0.7)
d. I failed classes or dropped out of school.	**	0.6% (±0.3)	1.1% (±0.3)	1.6% (±0.5)
e. I got sick (vomited) or had a hangover.	**	1.6% (±0.5)	5.1% (±0.9)	9.4% (±1.3)
f. I felt depressed, anxious, scared, or had other emotional problems.	**	0.9% (±0.3)	2.7% (±0.7)	3.6% (±0.6)
g. I got hurt or injured.	**	0.4% (±0.2)	1.2% (±0.4)	1.9% (±0.5)
h. I hurt or injured someone else.	**	0.2% (±0.2)	0.7% (±0.4)	0.6% (±0.3)
i. I got in trouble with my parents or family.	**	0.8% (±0.3)	3.5% (±0.9)	3.8% (±0.9)
j. I did things I didn't want to do or regretted afterward.	**	1.5% (±0.7)	3.5% (±0.7)	4.2% (±0.7)

Other Marijuana-Related Questions

	(n=0)	(n=4,438)	(n=3,784)	(n=2,923)
55. During the past 30 days, if you used marijuana, how did you usually use it?				
a. I did not use marijuana during the past 30 days.	**	92.5% (±1.3)	81.8% (±2.0)	73.0% (±2.5)
b. Smoked it (in a joint, bong, pipe, blunt)	**	4.9% (±1.0)	13.5% (±1.6)	19.9% (±2.3)
c. Ate it (in brownies, cakes, cookies, candy)	**	1.1% (±0.3)	2.7% (±0.7)	3.3% (±0.8)
d. Drank it (tea, cola, alcohol)	**	0.5% (±0.2)	0.5% (±0.2)	0.9% (±0.4)
e. Vaporized it	**	0.5% (±0.2)	1.0% (±0.3)	2.0% (±0.5)
f. Used it some other way	**	0.5% (±0.2)	0.5% (±0.2)	1.0% (±0.3)
56. During the past 30 days, how did you get marijuana? Choose all that apply.	(n=0)	(n=4,467)	(n=3,790)	(n=2,923)
a. I did not get marijuana in the past 30 days.	**	92.7% (±1.2)	81.7% (±2.0)	73.7% (±2.4)
b. I bought it from a store.	**	1.7% (±0.4)	2.2% (±0.5)	2.3% (±0.6)
c. I stole it from a store.	**	0.5% (±0.2)	1.1% (±0.4)	1.1% (±0.5)
d. I got it from friends.	**	3.4% (±0.7)	11.3% (±1.4)	16.8% (±1.8)
e. I got it at a party.	**	0.9% (±0.3)	3.5% (±0.7)	4.7% (±0.8)
f. I got it from an older brother or sister.	**	0.9% (±0.3)	1.8% (±0.5)	1.8% (±0.6)
g. I gave money to someone to get it for me.	**	1.5% (±0.5)	3.6% (±0.6)	4.2% (±0.7)
h. I took it from home without my parents' permission.	**	0.5% (±0.2)	0.8% (±0.3)	0.5% (±0.2)
i. I got it at home with my parents' permission.	**	0.4% (±0.2)	1.3% (±0.4)	1.7% (±0.6)
j. I got it some other way.	**	1.6% (±0.6)	3.1% (±0.7)	4.9% (±1.1)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,193)	% (± CI) (n=4,235)	% (± CI) (n=3,187)
57. In the past year, have your parents or guardians talked to you about why you should not use marijuana?				
a. Yes, a number of times	**	42.7% (±2.1)	38.0% (±1.7)	32.3% (±1.7)
b. Yes, once	**	22.5% (±1.2)	22.9% (±1.1)	19.6% (±1.3)
c. No	**	25.0% (±1.7)	32.1% (±1.3)	42.4% (±1.7)
d. I don't remember.	**	9.8% (±1.0)	7.1% (±0.8)	5.7% (±0.8)
58. Does anyone who lives with you now use marijuana?	(n=0)	(n=4,439)	(n=3,766)	(n=2,921)
a. No	**	85.8% (±1.7)	79.9% (±2.0)	79.3% (±2.1)
b. Yes	**	14.2% (±1.7)	20.1% (±2.0)	20.7% (±2.1)
59. During the past 30 days, on how many days did you use marijuana on school property?	(n=0)	(n=4,539)	(n=3,838)	(n=2,953)
a. 0 days	**	96.7% (±0.8)	92.9% (±1.2)	92.2% (±1.2)
b. 1-2 days	**	1.2% (±0.4)	3.5% (±0.8)	3.5% (±0.7)
c. 3-9 days	**	0.8% (±0.3)	1.7% (±0.5)	2.0% (±0.6)
d. 10-29 days	**	0.6% (±0.2)	1.0% (±0.3)	1.1% (±0.3)
e. All 30 days	**	0.8% (±0.3)	0.9% (±0.3)	1.3% (±0.4)
60. How wrong do your friends feel it would be for you to: Use marijuana?	(n=0)	(n=4,231)	(n=3,673)	(n=2,862)
a. Very wrong	**	74.4% (±2.5)	46.4% (±2.7)	32.4% (±2.5)
b. Wrong	**	13.2% (±1.2)	18.7% (±1.3)	19.6% (±1.8)
c. A little bit wrong	**	6.7% (±0.9)	17.2% (±1.4)	19.8% (±1.7)
d. Not at all wrong	**	5.7% (±1.3)	17.6% (±1.7)	28.2% (±2.8)

Other Alcohol- and Drug-Related Questions

61. How many times in the past year (12 months) have you been drunk or high at school?	(n=0)	(n=5,042)	(n=4,126)	(n=3,109)
a. Never	**	94.3% (±1.0)	86.1% (±1.7)	82.5% (±2.0)
b. 1-2 times	**	2.9% (±0.6)	6.4% (±0.9)	6.8% (±1.2)
c. 3-5 times	**	1.1% (±0.3)	2.7% (±0.5)	3.3% (±0.8)
d. 6-9 times	**	0.5% (±0.2)	1.8% (±0.4)	2.3% (±0.5)
e. 10 or more times	**	1.1% (±0.4)	3.0% (±0.6)	5.1% (±1.0)
62. During the past year in school, how many times did you get information in classes about reasons not to use alcohol or drugs?	(n=8,225)	(n=5,172)	(n=4,230)	(n=3,179)
a. Never	24.7% (±2.3)	12.8% (±2.2)	17.9% (±2.7)	30.9% (±3.1)
b. Once	18.7% (±1.5)	20.8% (±2.1)	26.3% (±2.0)	31.4% (±1.8)
c. 2-3 times	24.8% (±1.4)	32.2% (±1.8)	32.4% (±2.6)	26.8% (±2.6)
d. 4 or more times	31.9% (±3.4)	34.2% (±4.0)	23.4% (±2.5)	10.9% (±1.6)
63. How wrong do your parents feel it would be for you to: Use prescription drugs not prescribed to you?	(n=0)	(n=5,312)	(n=4,331)	(n=3,279)
a. Very wrong	**	86.8% (±1.2)	83.0% (±1.6)	81.6% (±1.7)
b. Wrong	**	9.9% (±0.9)	11.7% (±1.3)	12.3% (±1.2)
c. A little bit wrong	**	2.1% (±0.4)	3.5% (±0.6)	3.8% (±0.7)
d. Not at all wrong	**	1.3% (±0.3)	1.8% (±0.4)	2.3% (±0.6)

64. How wrong do your friends feel it would be for you to: Use prescription drugs not prescribed to you?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,213)	% (± CI) (n=3,662)	% (± CI) (n=2,854)
a. Very wrong	**	77.9% (±1.5)	65.0% (±1.9)	58.4% (±2.4)
b. Wrong	**	14.4% (±1.0)	20.8% (±1.3)	22.6% (±1.6)
c. A little bit wrong	**	5.1% (±0.8)	9.7% (±1.1)	11.8% (±1.4)
d. Not at all wrong	**	2.6% (±0.6)	4.6% (±0.8)	7.1% (±1.3)

65. How much do you think people risk harming themselves if they: Use prescription drugs that are not prescribed to them?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=0)	(n=5,165)	(n=4,212)	(n=3,173)
a. No risk	**	3.8% (±0.8)	2.8% (±0.7)	2.4% (±0.7)
b. Slight risk	**	5.1% (±0.6)	5.9% (±0.9)	6.0% (±1.0)
c. Moderate risk	**	16.5% (±1.2)	18.9% (±1.3)	19.5% (±1.7)
d. Great risk	**	67.1% (±2.3)	67.6% (±2.2)	68.9% (±2.1)
e. Not sure	**	7.6% (±1.1)	4.8% (±0.8)	3.3% (±0.6)

Other Health Concerns

Nutrition and Fitness

This section provides results regarding other health concerns including nutrition and fitness, health conditions and health care, safety, behaviors related to intentional injury, depression, and gambling behavior. Proper nutrition and exercise are critical components of a healthy life, as is access to preventive health care. Safety-related behaviors, such as wearing a seat belt when in a moving vehicle, can profoundly influence the outcome of an accident. Injury is the leading cause of death for adolescents aged 10 to 19, and violence contributes to injury-related deaths. People who are depressed experience a range of symptoms, and depression is associated with suicidal behavior.

Obese: "Obese" includes students who are in the top 5% for body mass index by age and gender based on growth charts developed by the Centers for Disease Control and Prevention (2000). "Overweight" includes students who are in the top 15% but not the top 5%.

66. Obese or overweight? (Computed from numeric responses to "How tall are you without your shoes on?" and "How much do you weigh without your shoes on?")

	(n=0)	(n=4,620)	(n=4,126)	(n=3,134)
Obese	**	9.3% (±1.2)	11.2% (±1.6)	11.1% (±1.7)
Overweight	**	13.6% (±1.4)	13.8% (±1.6)	13.3% (±1.5)
Normal weight	**	73.3% (±2.1)	71.3% (±2.6)	71.6% (±2.5)
Underweight	**	3.8% (±0.5)	3.7% (±0.6)	4.0% (±0.6)

67. Which of the following are you trying to do about your weight?

	(n=0)	(n=5,193)	(n=4,361)	(n=3,260)
a. I am not trying to do anything about my weight.	**	36.9% (±2.5)	30.3% (±2.2)	31.5% (±2.1)
b. Lose weight	**	38.1% (±2.7)	40.1% (±2.4)	40.4% (±2.7)
c. Gain weight	**	8.7% (±0.8)	13.4% (±1.1)	14.7% (±1.3)
d. Stay the same weight	**	16.3% (±1.0)	16.2% (±1.2)	13.4% (±1.4)

68. How often do you eat dinner with your family?

	(n=8,041)	(n=5,229)	(n=4,361)	(n=3,266)
a. Never	3.0% (±0.4)	5.2% (±0.8)	6.3% (±0.9)	8.7% (±1.2)
b. Rarely	9.2% (±1.0)	10.3% (±1.1)	13.6% (±1.4)	14.8% (±1.6)
c. Sometimes	12.5% (±0.8)	16.7% (±1.2)	19.0% (±1.4)	23.2% (±1.7)
d. Most of the time	29.6% (±1.1)	33.3% (±2.0)	34.3% (±2.2)	33.1% (±2.0)
e. Always	45.7% (±1.6)	34.5% (±1.6)	26.8% (±1.6)	20.2% (±1.3)

	Grade 6	Grade 8	Grade 10	Grade 12
69. How often in the past 12 months did you or your family have to cut meal size or skip meals because there wasn't enough money for food?	% (± CI) (n=0)	% (± CI) (n=10,155)	% (± CI) (n=8,427)	% (± CI) (n=6,390)
a. Almost every month	**	3.8% (±0.6)	3.9% (±0.7)	4.8% (±0.8)
b. Some months but not every month	**	3.5% (±0.6)	4.7% (±0.6)	5.6% (±1.2)
c. Only 1-2 months	**	3.9% (±0.6)	4.7% (±0.7)	3.8% (±0.7)
d. Did not have to skip or cut the size of meals.	**	88.9% (±1.3)	86.8% (±1.5)	85.8% (±2.1)
70. Number of servings of fruits and vegetables eaten per day (Computed from FV1 – FV6)	(n=0)	(n=5,212)	(n=4,382)	(n=3,274)
a. Less than 1	**	11.2% (±1.2)	10.2% (±1.4)	9.8% (±1.4)
b. 1 to less than 3	**	41.1% (±1.6)	45.1% (±1.8)	44.2% (±1.8)
c. 3 to less than 5	**	23.2% (±1.5)	23.2% (±1.8)	24.3% (±2.1)
d. 5 or more	**	24.5% (±1.5)	21.5% (±1.5)	21.7% (±1.7)
71. Did you eat breakfast today?	(n=8,362)	(n=5,223)	(n=4,365)	(n=3,265)
a. Yes	79.7% (±2.0)	69.4% (±2.7)	65.0% (±3.1)	58.6% (±2.6)
b. No	20.3% (±2.0)	30.6% (±2.7)	35.0% (±3.1)	41.4% (±2.6)
72. During the past 7 days, how many times did you drink regular soda, sports drinks (such as Gatorade) and other flavored sweetened drinks (such as Snapple or SoBe)? Do not include diet drinks.	(n=0)	(n=5,236)	(n=4,381)	(n=3,272)
a. 0 times	**	20.8% (±2.2)	20.2% (±1.9)	22.0% (±2.4)
b. 1-3 times	**	44.7% (±1.5)	40.5% (±1.2)	37.9% (±1.5)
c. 4-6 times	**	16.4% (±1.1)	19.8% (±1.5)	18.9% (±1.5)
d. 1 time per day	**	8.1% (±0.8)	8.5% (±1.1)	9.3% (±1.1)
e. 2 times per day	**	4.7% (±0.6)	5.8% (±0.9)	6.3% (±1.0)
f. 3 times per day	**	2.0% (±0.5)	2.5% (±0.5)	2.5% (±0.6)
g. 4 or more times per day	**	3.3% (±0.7)	2.7% (±0.6)	3.1% (±0.8)
73. During the past 7 days, how many times did you drink regular soda, sports drinks (such as Gatorade) and other flavored sweetened drinks (such as Snapple or SoBe) at school (including any after-school and weekend activities)? Do not include diet drinks.	(n=0)	(n=5,251)	(n=4,390)	(n=3,276)
a. 0 times	**	59.7% (±2.8)	51.4% (±2.8)	56.3% (±2.9)
b. 1-3 times	**	28.9% (±2.1)	35.4% (±2.1)	31.1% (±2.0)
c. 4-6 times	**	6.9% (±0.8)	9.4% (±1.0)	8.3% (±1.1)
d. 7-9 times	**	2.2% (±0.5)	2.1% (±0.5)	2.4% (±0.5)
e. 10 times or more	**	2.4% (±0.6)	1.6% (±0.3)	1.9% (±0.5)
74. During the past 7 days, where did you usually get the soda or other sweetened drinks that you drank at school? (Choose only one answer.)	(n=0)	(n=5,246)	(n=4,373)	(n=3,272)
a. I did not drink sodas, sports drinks, or other flavored drinks at school.	**	60.1% (±2.6)	48.0% (±2.6)	51.2% (±2.9)
b. I brought them from home.	**	21.9% (±1.9)	23.0% (±1.9)	24.1% (±2.3)
c. I got them from friends.	**	3.6% (±0.5)	5.3% (±0.7)	3.2% (±0.7)
d. I bought them at school.	**	6.6% (±1.6)	13.1% (±2.7)	8.3% (±1.9)
e. Other	**	7.7% (±1.2)	10.7% (±2.5)	13.1% (±2.1)

	Grade 6	Grade 8	Grade 10	Grade 12
75. How many sodas, sports drinks (such as Gatorade) and other sweetened drinks (such as Snapple or SoBe) did you drink yesterday?	% (± CI) (n=8,345)	% (± CI) (n=0)	% (± CI) (n=0)	% (± CI) (n=0)
a. None	53.2% (±1.7)	**	**	**
b. 1	33.6% (±1.3)	**	**	**
c. 2	8.7% (±0.7)	**	**	**
d. 3	2.1% (±0.3)	**	**	**
e. 4 or more	2.4% (±0.5)	**	**	**
76. Did you buy sodas, sports drinks, or other flavored sweetened drinks at school yesterday?	(n=8,274)	(n=0)	(n=0)	(n=0)
a. I did not drink sodas, sports drinks or other flavored sweetened drinks yesterday.	42.0% (±1.6)	**	**	**
b. Yes	11.3% (±1.3)	**	**	**
c. No	46.7% (±1.6)	**	**	**
77. During the past 7 days, how many times did you eat any potato chips or similar snack foods such as corn chips or cheese puffs at school (including any after-school and weekend activities)? Do not include reduced fat or fat-free items.	(n=0)	(n=5,239)	(n=4,368)	(n=3,266)
a. 0 times	**	41.7% (±2.5)	45.1% (±2.4)	49.6% (±2.8)
b. 1-3 times	**	40.8% (±1.8)	38.8% (±1.8)	35.9% (±2.1)
c. 4-6 times	**	12.0% (±1.2)	11.7% (±1.1)	10.0% (±1.2)
d. 7-9 times	**	3.0% (±0.5)	2.9% (±0.6)	2.5% (±0.6)
e. 10 times or more	**	2.4% (±0.5)	1.5% (±0.3)	1.9% (±0.4)
78. During the past 7 days, where did you usually get the chips and similar snack items you ate at school? (Choose only one answer)	(n=0)	(n=5,223)	(n=4,363)	(n=3,259)
a. I did not eat potato chips or similar snack foods at school.	**	42.9% (±2.6)	43.7% (±2.5)	47.3% (±2.8)
b. I brought them from home	**	29.0% (±1.7)	28.9% (±2.2)	28.2% (±2.5)
c. I got them from friends	**	8.6% (±1.1)	6.9% (±0.8)	5.3% (±0.9)
d. I bought them at school	**	10.4% (±2.3)	11.3% (±2.2)	10.6% (±2.0)
e. Other	**	9.1% (±1.1)	9.1% (±1.4)	8.6% (±1.0)
79. In the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increases your heart rate or makes you breathe hard some of the time.)	(n=8,898)	(n=5,206)	(n=4,356)	(n=3,258)
a. 0 days	5.4% (±0.7)	7.6% (±1.2)	12.0% (±1.7)	15.6% (±1.6)
b. 1 day	5.1% (±0.6)	5.6% (±0.8)	6.9% (±0.9)	7.9% (±1.0)
c. 2 days	7.6% (±0.7)	7.5% (±0.9)	8.2% (±0.9)	8.7% (±1.1)
d. 3 days	11.0% (±0.9)	9.8% (±1.0)	11.2% (±1.5)	10.8% (±1.1)
e. 4 days	12.7% (±0.7)	11.2% (±1.3)	9.8% (±1.1)	10.6% (±1.3)
f. 5 days	17.8% (±1.0)	16.8% (±1.2)	17.6% (±1.4)	15.2% (±1.3)
g. 6 days	11.2% (±1.1)	10.5% (±0.9)	10.7% (±1.4)	10.2% (±1.2)
h. 7 days	29.2% (±1.5)	31.0% (±1.8)	23.5% (±2.0)	21.1% (±1.9)

80. On how many of the past 7 days did you do exercises to strengthen or tone your muscles, such as push-ups, sit-ups, or weight lifting?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,208)	% (± CI) (n=4,348)	% (± CI) (n=3,254)
a. 0 days	**	20.3% (±2.0)	24.7% (±2.5)	29.4% (±2.1)
b. 1 day	**	10.9% (±0.8)	10.6% (±1.1)	9.8% (±1.2)
c. 2 days	**	13.1% (±1.3)	12.6% (±1.1)	10.4% (±1.1)
d. 3 days	**	13.0% (±1.2)	12.8% (±1.7)	11.9% (±1.3)
e. 4 days	**	9.9% (±0.9)	9.4% (±1.1)	9.6% (±1.3)
f. 5 days	**	14.2% (±1.6)	13.2% (±1.7)	13.1% (±1.8)
g. 6 days	**	4.1% (±0.6)	4.6% (±0.6)	5.1% (±1.0)
h. 7 days	**	14.6% (±1.5)	12.0% (±1.5)	10.5% (±1.0)

81. On average how many days a week do you walk to or from school?	(n=9,090)	(n=5,175)	(n=4,316)	(n=3,242)
a. Never	65.7% (±4.2)	65.8% (±5.2)	66.3% (±5.0)	78.4% (±4.0)
b. 1-2	10.5% (±1.2)	11.1% (±1.5)	11.5% (±1.6)	7.4% (±1.3)
c. 3-4	6.8% (±1.0)	5.7% (±1.0)	6.0% (±1.2)	3.6% (±1.0)
d. I walk every day.	17.0% (±2.6)	17.4% (±3.8)	16.1% (±3.0)	10.6% (±2.5)

82. On average how many days a week do you ride a bicycle to or from school?	(n=9,031)	(n=5,151)	(n=4,302)	(n=3,230)
a. Never	91.5% (±1.1)	94.1% (±0.8)	94.6% (±0.7)	95.7% (±0.8)
b. 1-2	4.7% (±0.7)	3.3% (±0.6)	2.6% (±0.5)	2.2% (±0.5)
c. 3-4	1.7% (±0.3)	1.2% (±0.3)	1.2% (±0.3)	1.1% (±0.4)
d. I bike every day.	2.1% (±0.4)	1.5% (±0.3)	1.5% (±0.4)	1.0% (±0.4)

83. On an average school day, how many hours do you watch TV?	(n=9,044)	(n=5,189)	(n=4,324)	(n=3,239)
a. I do not watch TV on an average school day.	15.9% (±1.5)	18.5% (±1.5)	22.5% (±1.6)	24.5% (±2.2)
b. Less than 1 hour per day	25.4% (±1.4)	22.9% (±1.3)	22.5% (±1.2)	22.4% (±1.3)
c. 1 hour per day	19.2% (±0.9)	19.0% (±1.3)	17.0% (±1.3)	15.5% (±1.7)
d. 2 hours per day	20.3% (±1.1)	19.1% (±1.0)	18.5% (±1.4)	17.7% (±1.4)
e. 3 hours per day	10.2% (±1.0)	10.9% (±1.2)	10.4% (±1.0)	10.5% (±1.2)
f. 4 hours per day	4.1% (±0.4)	4.3% (±0.8)	4.0% (±0.7)	4.8% (±1.0)
g. 5 or more hours per day	4.9% (±0.7)	5.2% (±0.9)	5.0% (±1.1)	4.6% (±0.9)

84. On an average school day, how many hours do you play video or computer games or use a computer for something that is not school work? (Count time spent on such things as Xbox, PlayStation, an iPod, an iPad or other tablet, a smartphone, YouTube, Facebook or other social networking tools, and the Internet.)	(n=8,971)	(n=5,193)	(n=4,329)	(n=3,241)
a. I do not play video games or use a computer for fun on an average school day.	14.5% (±0.9)	13.1% (±1.3)	16.4% (±1.3)	21.5% (±1.7)
b. Less than 1 hour per day	26.0% (±1.4)	17.4% (±1.5)	15.1% (±1.0)	14.6% (±1.3)
c. 1 hour per day	17.8% (±0.8)	15.0% (±1.1)	12.9% (±1.4)	13.5% (±1.5)
d. 2 hours per day	17.0% (±0.9)	16.5% (±1.2)	16.4% (±1.5)	15.2% (±1.4)
e. 3 hours per day	9.9% (±0.7)	14.2% (±1.3)	13.4% (±1.1)	11.7% (±1.0)
f. 4 hours per day	6.0% (±0.5)	8.6% (±0.7)	8.2% (±0.7)	9.2% (±1.2)
g. 5 or more hours per day	8.8% (±0.9)	15.1% (±1.8)	17.5% (±2.2)	14.2% (±1.6)

85. In an average week when you are in school, on how many days do you go to physical education (PE) classes?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,133)	% (± CI) (n=4,302)	% (± CI) (n=3,229)
a. 0 days	**	34.0% (±6.1)	56.1% (±8.5)	64.2% (±4.4)
b. 1 day	**	1.6% (±0.4)	1.6% (±0.6)	1.6% (±0.5)
c. 2 days	**	5.0% (±2.3)	2.8% (±1.6)	2.2% (±0.7)
d. 3 days	**	14.1% (±8.1)	10.7% (±8.6)	5.6% (±2.6)
e. 4 days	**	4.3% (±4.2)	3.3% (±2.7)	3.9% (±2.4)
f. 5 days	**	41.0% (±8.4)	25.6% (±7.0)	22.5% (±5.4)

86. During an average PE class, how many minutes do you spend actually exercising or playing sports?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=0)	(n=5,162)	(n=4,315)	(n=3,239)
a. I do not take PE	**	31.4% (±5.9)	51.9% (±7.9)	60.1% (±4.3)
b. Less than 10 minutes	**	2.0% (±0.6)	1.6% (±0.4)	1.8% (±0.6)
c. 10-20 minutes	**	4.4% (±1.0)	2.5% (±0.5)	2.1% (±0.6)
d. 21-30 minutes	**	9.0% (±1.4)	5.3% (±1.0)	4.9% (±1.0)
e. 31-40 minutes	**	14.8% (±1.9)	9.8% (±1.5)	9.3% (±1.9)
f. 41-50 minutes	**	18.1% (±2.3)	12.3% (±2.4)	10.4% (±2.0)
g. 51-60 minutes	**	12.0% (±1.8)	9.0% (±2.0)	6.2% (±0.9)
h. More than 60 minutes	**	8.3% (±4.1)	7.5% (±4.7)	5.2% (±2.2)

87. During the average week, on how many days do you participate in supervised after-school activities either at school or away from school? Include activities such as sports, art, music, dance, drama, or community service, religious, or club activities.	Grade 6	Grade 8	Grade 10	Grade 12
	(n=0)	(n=10,184)	(n=8,452)	(n=6,387)
a. 0 days	**	33.8% (±2.5)	32.2% (±2.6)	33.9% (±3.2)
b. 1-2 days	**	20.6% (±1.7)	23.3% (±1.4)	23.6% (±1.5)
c. 3 or more days	**	45.5% (±2.4)	44.5% (±2.6)	42.4% (±2.9)

Health Conditions and Health Care

88. Has a doctor or nurse ever told you that you have asthma?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=9,071)	(n=5,138)	(n=4,315)	(n=3,231)
a. Yes	14.1% (±0.9)	18.4% (±1.2)	21.5% (±1.4)	20.9% (±1.4)
b. No	74.9% (±1.4)	73.1% (±1.7)	72.9% (±1.6)	75.0% (±1.5)
c. Not sure	11.0% (±0.9)	8.5% (±0.9)	5.6% (±0.8)	4.1% (±0.8)

89. Do you still have asthma?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=9,052)	(n=5,127)	(n=4,308)	(n=3,228)
a. I have never had asthma.	67.3% (±1.4)	67.1% (±1.8)	65.1% (±1.9)	64.1% (±2.0)
b. Yes	10.4% (±0.8)	9.9% (±0.9)	12.7% (±1.1)	10.7% (±1.0)
c. No	11.7% (±0.8)	12.1% (±1.1)	13.4% (±1.0)	16.9% (±1.9)
d. Not sure	10.6% (±0.8)	10.9% (±1.1)	8.8% (±1.1)	8.2% (±0.9)

90. Symptoms of asthma include coughing, wheezing, shortness of breath, and chest tightness when you don't have a cold or the flu. During the past 4 weeks, about how many days per week on average did you have any symptoms of asthma?

	Grade 6 % (± CI) (n=0)	Grade 8 % (± CI) (n=4,701)	Grade 10 % (± CI) (n=4,005)	Grade 12 % (± CI) (n=3,072)
a. Never	**	78.3% (±1.8)	77.7% (±1.7)	80.7% (±1.4)
b. 1 or 2 days per week	**	11.3% (±1.0)	12.1% (±1.1)	10.5% (±1.0)
c. 3 to 6 days per week	**	3.9% (±0.6)	4.3% (±0.8)	3.5% (±0.5)
d. Every day, but not throughout the day	**	2.7% (±0.5)	2.2% (±0.5)	1.9% (±0.5)
e. Every day, throughout the day	**	0.9% (±0.3)	1.1% (±0.3)	0.9% (±0.4)
f. Not sure	**	3.0% (±0.7)	2.6% (±0.5)	2.5% (±0.6)

91. RESCUE inhalers are asthma medicine that you breath in through your mouth that gives you QUICK relief from asthma symptoms. They are PRESCRIBED by a doctor. During the past 4 weeks, about how many days per week on average did you use a rescue inhaler?

	(n=0)	(n=4,816)	(n=4,072)	(n=3,109)
a. Never	**	87.4% (±2.1)	88.1% (±1.8)	89.3% (±2.1)
b. 1 or 2 days per week	**	2.8% (±0.5)	4.5% (±0.7)	4.3% (±0.7)
c. 3 to 6 days per week	**	1.5% (±0.3)	1.8% (±0.5)	1.7% (±0.5)
d. Every day, once per day	**	0.6% (±0.3)	0.9% (±0.3)	0.7% (±0.3)
e. Every day, twice or more per day	**	6.5% (±1.8)	3.7% (±1.5)	3.0% (±1.8)
f. Not sure	**	1.2% (±0.4)	1.0% (±0.3)	1.0% (±0.4)

92. During the past 12 months, how many times did you visit an emergency room or urgent care center because of your asthma?

	(n=0)	(n=4,699)	(n=4,001)	(n=3,050)
a. I do not have asthma.	**	77.4% (±2.3)	75.0% (±2.1)	76.3% (±1.7)
b. None	**	15.9% (±1.5)	19.4% (±1.7)	19.9% (±1.3)
c. 1 to 3 times	**	2.8% (±0.5)	3.4% (±0.6)	2.1% (±0.5)
d. 4 to 9 times	**	0.5% (±0.2)	0.7% (±0.2)	0.7% (±0.4)
e. 10 to 12 times	**	1.1% (±0.5)	0.3% (±0.2)	0.3% (±0.2)
f. More than 12 times	**	0.1% (±0.1)	0.2% (±0.2)	0.1% (±0.1)
g. Not sure	**	2.2% (±0.6)	1.0% (±0.4)	0.7% (±0.3)

93. During the past 12 months, how many days did you stay out of school or stay away from your usual activities because of your asthma?

	(n=0)	(n=4,654)	(n=3,976)	(n=3,042)
a. I do not have asthma.	**	79.8% (±2.0)	77.1% (±2.0)	78.6% (±1.7)
b. None	**	13.8% (±1.3)	17.7% (±1.7)	17.4% (±1.3)
c. 1 to 2 days	**	3.8% (±1.3)	2.3% (±0.5)	2.2% (±0.7)
d. 3 to 4 days	**	0.6% (±0.2)	1.0% (±0.3)	0.8% (±0.4)
e. 5 to 10 days	**	0.4% (±0.2)	0.9% (±0.3)	0.4% (±0.2)
f. More than 10 days	**	0.5% (±0.2)	0.4% (±0.2)	0.4% (±0.2)
g. Not sure	**	1.1% (±0.3)	0.6% (±0.3)	0.3% (±0.2)

94. When was the last time you saw a doctor or health care provider for a check-up or physical exam when you were not sick or injured?

	(n=0)	(n=5,118)	(n=4,295)	(n=3,224)
a. During the past 12 months	**	65.2% (±2.7)	66.1% (±2.1)	60.6% (±2.4)
b. Between 12 and 24 months ago	**	13.1% (±1.2)	15.5% (±1.4)	19.0% (±1.4)
c. More than 24 months ago	**	4.2% (±0.6)	6.2% (±0.7)	8.5% (±1.3)
d. Never	**	2.0% (±0.5)	2.1% (±0.5)	3.1% (±0.7)
e. Not sure	**	15.6% (±2.0)	10.1% (±1.5)	8.8% (±1.2)

	Grade 6	Grade 8	Grade 10	Grade 12
95. When was the last time you saw a dentist for a check-up, exam, teeth cleaning, or other dental work?	% (± CI) (n=0)	% (± CI) (n=5,118)	% (± CI) (n=4,291)	% (± CI) (n=3,225)
a. During the past 12 months	**	76.9% (±2.3)	79.0% (±2.5)	75.5% (±2.6)
b. Between 12 and 24 months ago	**	8.3% (±0.9)	8.5% (±1.1)	11.3% (±1.1)
c. More than 24 months ago	**	4.0% (±0.7)	5.1% (±0.9)	7.0% (±1.2)
d. Never	**	1.3% (±0.4)	1.5% (±0.4)	1.4% (±0.4)
e. Not sure	**	9.4% (±1.4)	5.8% (±1.2)	4.8% (±0.9)
96. During the past 12 months, how many days did you miss some school because of toothache? (Do not include toothache due to braces or injury.)	(n=0)	(n=5,116)	(n=4,285)	(n=3,220)
a. None	**	93.8% (±1.1)	94.0% (±0.9)	93.2% (±1.3)
b. 1-4 days	**	3.4% (±0.7)	3.4% (±0.6)	3.8% (±0.8)
c. 5 days or more	**	0.9% (±0.3)	1.1% (±0.3)	1.3% (±0.5)
d. Not sure	**	1.9% (±0.5)	1.5% (±0.4)	1.7% (±0.5)
97. During the past year, did you miss any time from school because of toothache? (Do not include toothache due to braces or an injury.)	(n=8,988)	(n=0)	(n=0)	(n=0)
a. Yes	6.1% (±0.6)	**	**	**
b. No	85.5% (±1.1)	**	**	**
c. Not sure	8.4% (±0.8)	**	**	**
98. Have you ever been told by a doctor or other health professional that you have diabetes?	(n=0)	(n=5,108)	(n=4,278)	(n=3,224)
a. No	**	94.7% (±0.7)	94.7% (±0.8)	94.4% (±1.0)
b. Yes	**	2.9% (±0.5)	3.2% (±0.6)	3.2% (±0.7)
c. Not sure	**	2.4% (±0.5)	2.2% (±0.5)	2.4% (±0.5)
99. On an average school night how many hours do you sleep?	(n=0)	(n=4,916)	(n=4,141)	(n=3,127)
a. 5 hours or less	**	8.9% (±1.1)	14.5% (±1.4)	16.3% (±1.8)
b. About 6 hours	**	12.3% (±1.2)	23.9% (±1.5)	28.8% (±1.9)
c. About 7 hours	**	23.8% (±1.7)	30.5% (±1.6)	31.8% (±1.9)
d. About 8 hours	**	39.2% (±1.7)	25.5% (±1.7)	18.5% (±1.9)
e. 9 hours or more	**	15.8% (±1.7)	5.6% (±0.8)	4.5% (±1.0)

Sexual Behavior

100. Have you ever had sexual intercourse? †	(n=0)	(n=1,091)	(n=1,510)	(n=1,124)
a. Yes	**	9.3% (±4.0)	26.6% (±4.6)	51.6% (±3.8)
b. No	**	90.7% (±4.0)	73.4% (±4.6)	48.4% (±3.8)
101. How old were you when you had sexual intercourse for the first time? †	(n=0)	(n=1,091)	(n=1,510)	(n=1,124)
a. I have never had sexual intercourse.	**	90.7% (±4.0)	73.4% (±4.6)	48.4% (±3.8)
b. 11 years old or younger	**	2.2% (±1.4)	2.1% (±0.7)	2.6% (±1.1)
c. 12 years old	**	2.0% (±1.0)	1.5% (±0.5)	1.7% (±0.5)
d. 13 years old	**	3.6% (±1.9)	3.2% (±0.9)	3.6% (±1.5)
e. 14 years old	**	1.1% (±0.7)	8.1% (±2.8)	6.6% (±1.4)
f. 15 years old	**	0.1% (±0.2)	9.9% (±1.5)	10.9% (±1.9)
g. 16 years old	**	0.2% (±0.2)	1.4% (±0.6)	14.9% (±2.1)
h. 17 years old or older	**	0.2% (±0.3)	0.5% (±0.4)	11.3% (±1.7)

	Grade 6	Grade 8	Grade 10	Grade 12
102. During your life, with whom have you had sexual contact? †	% (± CI) (n=0)	% (± CI) (n=1,091)	% (± CI) (n=1,517)	% (± CI) (n=1,124)
a. I have never had sexual contact.	**	85.0% (±3.6)	57.5% (±4.6)	36.6% (±3.5)
b. Females	**	8.2% (±2.1)	20.8% (±2.3)	30.2% (±2.9)
c. Males	**	4.6% (±2.2)	16.2% (±2.4)	26.5% (±2.9)
d. Females and males	**	2.3% (±1.2)	5.6% (±1.3)	6.7% (±1.7)
103. With how many people have you ever had sexual intercourse? †	(n=0)	(n=1,089)	(n=1,508)	(n=1,126)
a. I have never had sexual intercourse.	**	91.1% (±3.8)	74.1% (±4.6)	49.6% (±3.3)
b. 1 person	**	3.9% (±1.5)	12.6% (±2.2)	19.3% (±2.6)
c. 2 people	**	1.9% (±1.1)	5.4% (±1.9)	10.0% (±1.7)
d. 3 people	**	1.2% (±1.1)	2.7% (±1.0)	6.8% (±2.0)
e. 4 people	**	0.6% (±0.8)	1.8% (±0.5)	4.8% (±1.3)
f. 5 people	**	0.1% (±0.2)	0.8% (±0.4)	2.8% (±1.3)
g. 6 or more people	**	1.3% (±0.7)	2.7% (±0.7)	6.7% (±2.3)
104. The last time you had sexual intercourse, did you or your partner use a condom? †	(n=0)	(n=1,088)	(n=1,509)	(n=1,127)
a. I have never had sexual intercourse.	**	90.7% (±4.2)	74.4% (±4.4)	48.8% (±3.6)
b. Yes	**	6.1% (±2.9)	16.0% (±2.3)	27.8% (±2.5)
c. No	**	3.2% (±1.7)	9.6% (±2.9)	23.4% (±2.7)

Behaviors Related to Unintentional Injury

105. How often do you wear a life vest when you're in a small boat like a canoe, raft, or small motorboat?	(n=0)	(n=5,294)	(n=4,455)	(n=3,338)
a. Never go boating	**	22.1% (±3.0)	19.1% (±2.9)	18.2% (±2.9)
b. Never	**	7.3% (±0.8)	11.2% (±1.2)	14.4% (±1.3)
c. Less than half the time	**	7.9% (±0.7)	12.6% (±1.3)	14.8% (±2.0)
d. About half the time	**	11.4% (±1.0)	14.1% (±1.5)	13.8% (±1.7)
e. More than half the time	**	18.6% (±1.9)	17.4% (±1.9)	16.4% (±1.4)
f. Always	**	32.6% (±1.4)	25.6% (±1.9)	22.5% (±2.0)
106. Have you ever taken formal swimming lessons?	(n=9,090)	(n=5,315)	(n=4,462)	(n=3,343)
a. Yes	55.8% (±3.7)	59.5% (±4.9)	59.3% (±5.8)	63.5% (±5.8)
b. No	37.2% (±3.8)	32.9% (±4.7)	34.8% (±5.6)	32.5% (±5.7)
c. Not sure	7.0% (±0.7)	7.6% (±1.0)	5.9% (±0.9)	4.0% (±0.9)
107. I am comfortable playing and swimming in water over my head.	(n=8,996)	(n=5,295)	(n=4,458)	(n=3,342)
a. Strongly agree	44.2% (±2.6)	53.8% (±3.5)	55.5% (±3.8)	58.1% (±3.8)
b. Agree	41.1% (±1.4)	33.5% (±2.2)	30.7% (±2.2)	28.3% (±2.3)
c. Disagree	10.5% (±1.3)	9.0% (±1.4)	8.6% (±1.5)	9.0% (±1.6)
d. Strongly disagree	4.1% (±0.6)	3.7% (±0.7)	5.2% (±1.1)	4.6% (±1.2)

	Grade 6	Grade 8	Grade 10	Grade 12
108. During the past 30 days, how many days did you ride in a car or other vehicle driven by someone who was texting or emailing?	% (± CI) (n=0)	% (± CI) (n=5,284)	% (± CI) (n=4,453)	% (± CI) (n=3,338)
a. 0 days	**	52.8% (±2.1)	43.2% (±2.3)	41.0% (±2.7)
b. 1 or 2 days	**	19.7% (±1.2)	20.9% (±1.4)	20.6% (±1.5)
c. 3 to 5 days	**	8.1% (±0.7)	11.8% (±1.0)	12.4% (±1.1)
d. 6 to 9 days	**	5.7% (±0.6)	7.1% (±0.9)	9.2% (±1.0)
e. 10 to 19 days	**	5.5% (±0.6)	7.6% (±0.8)	8.1% (±1.1)
f. 20 to 29 days	**	3.4% (±0.5)	3.5% (±0.6)	3.1% (±0.5)
g. All 30 days	**	4.7% (±0.6)	6.0% (±0.8)	5.6% (±1.0)
109. During the past 30 days, did you ride in a car or other vehicle driven by someone who was texting or emailing?	(n=8,944)	(n=0)	(n=0)	(n=0)
a. Yes	22.2% (±1.4)	**	**	**
b. No	60.3% (±2.0)	**	**	**
c. Not sure	17.5% (±1.1)	**	**	**
110. During the past 30 days, how many days did you text or email while driving a car or other vehicle?	(n=0)	(n=10,012)	(n=8,406)	(n=6,350)
a. I did not drive a car or other vehicle during the past 30 days.	**	84.1% (±1.1)	58.2% (±2.5)	27.2% (±3.2)
b. 0 days	**	11.5% (±1.0)	33.8% (±2.3)	29.9% (±1.4)
c. 1 or 2 days	**	1.0% (±0.2)	3.7% (±0.4)	12.4% (±1.2)
d. 3 to 5 days	**	0.8% (±0.2)	1.4% (±0.3)	6.9% (±0.8)
e. 6 to 9 days	**	0.4% (±0.1)	0.9% (±0.2)	5.0% (±0.8)
f. 10 to 19 days	**	0.6% (±0.2)	0.6% (±0.2)	5.9% (±0.9)
g. 20 to 29 days	**	0.4% (±0.1)	0.3% (±0.1)	4.8% (±0.8)
h. All 30 days	**	1.3% (±0.3)	1.1% (±0.2)	7.8% (±1.1)
111. During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been drinking alcohol?	(n=0)	(n=4,803)	(n=3,986)	(n=3,024)
a. 0 times	**	83.2% (±1.9)	82.3% (±1.3)	83.2% (±1.7)
b. 1 time	**	8.3% (±1.0)	8.3% (±0.7)	8.7% (±1.0)
c. 2-3 times	**	4.9% (±0.7)	6.0% (±0.8)	5.5% (±0.9)
d. 4-5 times	**	1.4% (±0.4)	1.7% (±0.5)	1.3% (±0.4)
e. 6 or more times	**	2.2% (±0.6)	1.7% (±0.5)	1.3% (±0.4)
112. In the last 30 days, have you ridden in a car driven by someone who had been drinking alcohol?	(n=8,343)	(n=0)	(n=0)	(n=0)
a. Yes	5.6% (±0.5)	**	**	**
b. No	86.6% (±0.8)	**	**	**
c. Not sure	7.8% (±0.6)	**	**	**
113. During the past 30 days, how many times did you drive a car or other vehicle when you had been drinking alcohol?	(n=0)	(n=4,793)	(n=3,979)	(n=3,019)
a. 0 times	**	97.0% (±0.6)	95.1% (±0.9)	91.2% (±1.4)
b. 1 time	**	1.3% (±0.4)	2.3% (±0.6)	5.0% (±0.8)
c. 2-3 times	**	0.8% (±0.3)	1.3% (±0.4)	2.3% (±0.7)
d. 4-5 times	**	0.4% (±0.2)	0.7% (±0.3)	0.9% (±0.3)
e. 6 or more times	**	0.6% (±0.2)	0.7% (±0.3)	0.6% (±0.2)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,786)	% (± CI) (n=3,976)	% (± CI) (n=3,015)
114. During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been using marijuana?				
a. 0 times	**	90.4% (±1.4)	81.0% (±1.7)	74.1% (±2.5)
b. 1 time	**	3.0% (±0.7)	6.7% (±0.9)	7.9% (±1.1)
c. 2-3 times	**	2.6% (±0.5)	5.5% (±0.9)	7.1% (±1.0)
d. 4-5 times	**	1.2% (±0.3)	2.0% (±0.5)	3.4% (±0.6)
e. 6 or more times	**	2.8% (±0.7)	4.8% (±1.0)	7.5% (±1.3)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,768)	% (± CI) (n=3,961)	% (± CI) (n=3,012)
115. During the past 30 days, how many times did you drive a car or other vehicle within three hours after using marijuana?				
a. 0 times	**	95.5% (±0.8)	90.9% (±1.4)	83.2% (±1.6)
b. 1 time	**	1.5% (±0.4)	3.5% (±0.6)	5.6% (±0.8)
c. 2-3 times	**	1.3% (±0.4)	2.5% (±0.7)	4.0% (±0.7)
d. 4-5 times	**	0.6% (±0.2)	1.1% (±0.4)	2.1% (±0.4)
e. 6 or more times	**	1.1% (±0.4)	2.1% (±0.6)	5.1% (±1.0)

Behaviors Related to Intentional Injury

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,253)	% (± CI) (n=8,537)	% (± CI) (n=6,402)
116. During the past 30 days, on how many days did you: Carry a weapon such as a gun, knife, or club on school property?				
a. 0 days	**	96.2% (±0.6)	94.3% (±0.7)	93.6% (±1.1)
b. 1-5 days	**	2.5% (±0.4)	3.2% (±0.5)	3.4% (±0.7)
c. 6 or more days	**	1.3% (±0.3)	2.5% (±0.4)	3.1% (±0.6)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,697)	% (± CI) (n=0)	% (± CI) (n=0)	% (± CI) (n=0)
117. During the past 30 days, did you carry a weapon such as a gun, knife, or club on school property?				
a. Yes	3.1% (±0.3)	**	**	**
b. No	96.9% (±0.3)	**	**	**

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,280)	% (± CI) (n=4,448)	% (± CI) (n=3,328)
118. During the past 30 days, on how many days did you carry a gun? (Do not include carrying a gun while hunting.)				
a. 0 times	**	95.7% (±0.6)	95.6% (±0.6)	95.7% (±0.9)
b. 1 time	**	2.3% (±0.4)	2.1% (±0.4)	2.0% (±0.5)
c. 2-3 times	**	0.9% (±0.2)	1.2% (±0.3)	0.8% (±0.3)
d. 4-5 times	**	0.3% (±0.1)	0.3% (±0.2)	0.5% (±0.2)
e. 6 or more times	**	0.9% (±0.3)	0.8% (±0.3)	1.0% (±0.4)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,615)	% (± CI) (n=10,480)	% (± CI) (n=8,657)	% (± CI) (n=6,513)
119. During the past 12 months, how many times were you in a physical fight?				
a. 0 times	76.6% (±1.4)	73.2% (±1.6)	77.6% (±1.2)	83.6% (±1.8)
b. 1 time	11.6% (±0.9)	13.3% (±0.8)	11.8% (±0.8)	8.0% (±0.9)
c. 2-3 times	6.5% (±0.6)	8.1% (±0.8)	6.5% (±0.5)	4.8% (±0.6)
d. 4-5 times	1.9% (±0.3)	2.3% (±0.3)	1.5% (±0.3)	1.4% (±0.4)
e. 6 or more times	3.4% (±0.4)	3.2% (±0.4)	2.6% (±0.4)	2.1% (±0.5)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,295)	% (± CI) (n=8,563)	% (± CI) (n=6,423)
120. A gang is a group of people with a leader who act together often for violent or illegal activities. During the past 12 months, have you been a member of a gang?				
a. No	**	94.5% (±0.6)	94.1% (±0.5)	95.0% (±0.6)
b. Yes	**	5.5% (±0.6)	5.9% (±0.5)	5.0% (±0.6)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,263)	% (± CI) (n=8,540)	% (± CI) (n=6,410)
121. Are there gangs at your school?				
a. No	**	31.6% (±3.6)	20.0% (±3.1)	31.0% (±4.7)
b. Yes	**	11.1% (±1.5)	19.4% (±3.3)	17.2% (±3.7)
c. Don't know	**	57.2% (±2.9)	60.7% (±3.1)	51.8% (±3.1)

Physical, Emotional and Sexual Abuse

122. Has an adult ever physically hurt you on purpose (like pushed, slapped, hit, kicked or punched you), leaving a mark, bruise or injury?

	(n=0)	(n=4,673)	(n=4,012)	(n=3,064)
a. No	**	77.8% (±1.5)	73.6% (±1.8)	75.1% (±2.2)
b. Yes	**	22.2% (±1.5)	26.4% (±1.8)	24.9% (±2.2)

123. During the past 12 months, did someone you were dating or going out with ever limit your activities, threaten you, or make you feel unsafe in any other way?

	(n=0)	(n=4,652)	(n=4,020)	(n=3,056)
a. I did not date or go out with anyone during the past 12 months.	**	57.5% (±3.0)	47.0% (±2.9)	35.3% (±2.0)
b. No	**	38.2% (±2.8)	44.7% (±2.3)	56.3% (±1.8)
c. Yes	**	4.3% (±0.7)	8.3% (±1.3)	8.4% (±1.0)

124. In the past 12 months, how many times did someone you were dating or going out with physically hurt you on purpose? (Count such things as being hit, slammed into something, or injured with an object or weapon.)

	(n=0)	(n=4,701)	(n=4,021)	(n=3,064)
a. I did not date or go out with anyone during the past 12 months.	**	59.2% (±3.0)	48.8% (±2.8)	38.0% (±2.3)
b. 0 times	**	36.4% (±2.6)	45.4% (±2.4)	55.8% (±2.1)
c. 1 time	**	1.5% (±0.4)	2.2% (±0.5)	2.2% (±0.5)
d. 2 or 3 times	**	1.1% (±0.3)	1.9% (±0.4)	2.0% (±0.7)
e. 4 or 5 times	**	0.4% (±0.2)	0.6% (±0.2)	0.7% (±0.3)
f. 6 or more times	**	1.3% (±0.5)	1.1% (±0.4)	1.3% (±0.4)

125. Have you ever been in a situation where someone made you engage in kissing, sexual touch or intercourse when you did not want to? †

	(n=0)	(n=1,159)	(n=1,572)	(n=1,153)
a. No	**	91.5% (±2.1)	84.9% (±2.5)	82.9% (±2.5)
b. Yes	**	8.5% (±2.1)	15.1% (±2.5)	17.1% (±2.5)

126. In the past 12 months, have you been in a situation where someone made you engage in kissing, sexual touch or intercourse when you did not want to? †

	(n=0)	(n=1,148)	(n=1,562)	(n=1,150)
a. No	**	93.9% (±1.8)	91.8% (±1.6)	90.5% (±2.1)
b. Yes	**	6.1% (±1.8)	8.2% (±1.6)	9.5% (±2.1)

Mental Health

	Grade 6	Grade 8	Grade 10	Grade 12
127. During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?	% (± CI) (n=0)	% (± CI) (n=10,440)	% (± CI) (n=8,625)	% (± CI) (n=6,493)
a. Yes	**	27.2% (±1.9)	34.9% (±2.0)	33.7% (±2.1)
b. No	**	72.8% (±1.9)	65.1% (±2.0)	66.3% (±2.1)
128. During the past 12 months, did you ever seriously consider attempting suicide?	(n=0)	(n=10,401)	(n=8,601)	(n=6,483)
a. Yes	**	16.1% (±1.5)	20.5% (±1.4)	18.1% (±1.3)
b. No	**	83.9% (±1.5)	79.5% (±1.4)	81.9% (±1.3)
129. During the past 12 months, did you make a plan about how you would attempt suicide?	(n=0)	(n=5,269)	(n=4,426)	(n=3,333)
a. Yes	**	13.9% (±1.5)	16.4% (±1.5)	14.3% (±1.5)
b. No	**	86.1% (±1.5)	83.6% (±1.5)	85.7% (±1.5)
130. During the past 12 months, how many times did you actually attempt suicide?	(n=0)	(n=5,286)	(n=4,440)	(n=3,332)
a. 0 times	**	91.1% (±1.1)	89.8% (±1.2)	92.4% (±1.1)
b. 1 time	**	5.2% (±0.7)	5.3% (±0.7)	4.6% (±0.8)
c. 2-3 times	**	2.4% (±0.5)	2.9% (±0.6)	2.2% (±0.6)
d. 4-5 times	**	0.4% (±0.2)	0.7% (±0.2)	0.3% (±0.2)
e. 6 or more times	**	0.9% (±0.4)	1.3% (±0.3)	0.5% (±0.2)
131. Have you ever seriously thought about killing yourself?	(n=8,525)	(n=0)	(n=0)	(n=0)
a. Yes	15.0% (±1.1)	**	**	**
b. No	85.0% (±1.1)	**	**	**
132. Have you ever tried to kill yourself?	(n=8,527)	(n=0)	(n=0)	(n=0)
a. Yes	4.5% (±0.5)	**	**	**
b. No	95.5% (±0.5)	**	**	**
133. How often over the last 2 weeks, were you bothered by: Feeling nervous, anxious or on edge?	(n=0)	(n=9,537)	(n=8,060)	(n=6,143)
a. Not at all	**	50.2% (±1.6)	35.5% (±1.6)	36.9% (±1.7)
b. Several days	**	32.8% (±1.2)	38.0% (±1.4)	38.3% (±1.4)
c. More than half the days	**	7.5% (±0.7)	11.6% (±0.7)	11.4% (±1.0)
d. Nearly every day	**	9.6% (±0.7)	14.9% (±1.2)	13.3% (±0.9)
134. How often over the last 2 weeks, were you bothered by: Not being able to stop or control worrying?	(n=0)	(n=9,506)	(n=8,058)	(n=6,128)
a. Not at all	**	62.7% (±1.6)	49.0% (±1.7)	48.7% (±1.5)
b. Several days	**	22.4% (±1.0)	28.1% (±1.0)	29.8% (±1.4)
c. More than half the days	**	6.7% (±0.6)	9.8% (±0.8)	9.8% (±1.0)
d. Nearly every day	**	8.2% (±0.8)	13.1% (±1.1)	11.7% (±0.7)
135. When you feel sad or hopeless, are there adults that you can turn to for help?	(n=8,581)	(n=5,275)	(n=4,428)	(n=3,333)
a. I never feel sad or hopeless.	20.6% (±1.2)	25.9% (±1.8)	20.6% (±1.2)	19.1% (±1.6)
b. Yes	62.4% (±1.7)	49.4% (±2.1)	50.3% (±2.0)	56.6% (±2.0)
c. No	8.1% (±0.9)	12.6% (±1.2)	15.0% (±1.5)	12.6% (±1.0)
d. Not sure	9.0% (±0.9)	12.2% (±1.2)	14.1% (±1.2)	11.7% (±1.4)

136. Last year, did you hear or see information at your school about the warning signs of suicide and how to get help for yourself or a friend?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,584)	% (± CI) (n=3,966)	% (± CI) (n=3,025)
a. Yes	**	38.6% (±5.5)	53.7% (±5.1)	47.5% (±4.3)
b. No	**	39.3% (±4.2)	30.5% (±4.2)	35.5% (±4.0)
c. Not sure	**	22.1% (±1.9)	15.8% (±1.6)	17.0% (±1.3)

Social and Emotional Learning

137. I know how to disagree without starting a fight or argument.	(n=7,903)	(n=4,485)	(n=3,920)	(n=3,003)
a. Strongly agree	38.7% (±1.4)	44.3% (±2.6)	47.0% (±2.1)	52.4% (±2.0)
b. Agree	48.3% (±1.3)	43.3% (±2.0)	43.5% (±1.8)	39.6% (±1.9)
c. Disagree	9.0% (±0.7)	9.3% (±1.2)	7.3% (±1.1)	5.9% (±0.9)
d. Strongly disagree	4.0% (±0.6)	3.1% (±0.6)	2.2% (±0.5)	2.1% (±0.5)

138. When I have problems at school, I am good at finding ways to solve them.	(n=7,913)	(n=4,559)	(n=3,915)	(n=3,003)
a. Strongly agree	32.0% (±1.4)	34.4% (±2.4)	34.3% (±2.0)	38.6% (±2.2)
b. Agree	54.2% (±1.2)	49.6% (±2.0)	51.7% (±1.9)	51.5% (±2.1)
c. Disagree	11.1% (±1.0)	12.6% (±1.5)	11.5% (±1.4)	8.2% (±1.3)
d. Strongly disagree	2.7% (±0.4)	3.4% (±0.7)	2.5% (±0.5)	1.6% (±0.5)

139. When I make a decision, I think about what might happen afterward.	(n=7,901)	(n=4,522)	(n=3,913)	(n=2,997)
a. Strongly agree	36.0% (±1.3)	35.1% (±1.9)	37.8% (±2.0)	42.3% (±1.8)
b. Agree	50.4% (±1.3)	47.2% (±1.6)	47.0% (±1.9)	45.4% (±1.9)
c. Disagree	10.7% (±0.7)	14.1% (±1.5)	12.3% (±1.1)	9.9% (±1.1)
d. Strongly disagree	2.8% (±0.5)	3.6% (±0.8)	2.9% (±0.7)	2.4% (±0.7)

140. I get along well with students who are different from me.	(n=7,946)	(n=4,541)	(n=3,915)	(n=3,003)
a. Strongly agree	45.2% (±1.4)	42.6% (±2.0)	40.1% (±1.9)	43.0% (±2.2)
b. Agree	46.6% (±1.2)	45.6% (±1.8)	48.9% (±1.8)	47.9% (±2.0)
c. Disagree	6.1% (±0.7)	8.9% (±1.3)	8.5% (±1.4)	7.2% (±0.8)
d. Strongly disagree	2.2% (±0.4)	3.0% (±0.6)	2.5% (±0.6)	2.0% (±0.5)

141. I try to understand how other people feel and think.	(n=7,928)	(n=4,494)	(n=3,910)	(n=2,998)
a. Strongly agree	42.6% (±1.5)	42.8% (±2.1)	45.4% (±2.0)	50.2% (±2.4)
b. Agree	49.0% (±1.2)	45.6% (±1.8)	45.3% (±1.7)	41.4% (±2.5)
c. Disagree	6.7% (±0.6)	9.4% (±1.3)	6.8% (±0.9)	6.0% (±1.1)
d. Strongly disagree	1.7% (±0.4)	2.2% (±0.5)	2.4% (±0.6)	2.4% (±0.6)

Gambling

142. In the past 12 months, how often have you gambled (bet) for money or possessions?	(n=0)	(n=4,728)	(n=3,937)	(n=3,005)
a. Never in the past year	**	77.9% (±1.5)	74.3% (±1.5)	76.1% (±1.8)
b. Once or twice in the past year	**	12.6% (±1.0)	13.5% (±1.0)	12.8% (±1.3)
c. A few times in the past year	**	5.5% (±0.7)	8.3% (±0.8)	6.4% (±1.0)
d. Once or twice a month	**	2.3% (±0.4)	2.4% (±0.5)	2.3% (±0.6)
e. At least once a week	**	1.7% (±0.4)	1.5% (±0.4)	2.3% (±0.5)

School Climate

In the past few years Washington State has given increased attention to supportive learning environments. Students need a safe, nurturing, healthy, and civil learning environment if they are to be successful in school. This section provides information about student perceptions of school climate.

143. A student is being bullied when another student, or group of students, say or do nasty or unpleasant things to him or her. It is also bullying when a student is teased repeatedly in a way he or she doesn't like. It is NOT bullying when two students of about the same strength argue or fight. In the last 30 days, how often have you been bullied?

	Grade 6 % (± CI) (n=8,668)	Grade 8 % (± CI) (n=10,602)	Grade 10 % (± CI) (n=8,739)	Grade 12 % (± CI) (n=6,573)
a. I have not been bullied.	69.1% (±1.9)	72.0% (±1.8)	77.4% (±1.3)	83.6% (±1.3)
b. Once	15.2% (±1.0)	12.7% (±0.9)	10.6% (±0.8)	7.6% (±0.6)
c. 2-3 times	8.4% (±0.6)	7.6% (±0.6)	6.4% (±0.6)	5.0% (±0.7)
d. About once a week	2.8% (±0.4)	3.6% (±0.4)	2.5% (±0.4)	1.8% (±0.3)
e. Several times a week	4.4% (±0.5)	4.1% (±0.6)	3.2% (±0.4)	2.1% (±0.3)

144. When a student is being bullied at school, how often do teachers or other adults at school try to put a stop to it?

	(n=0)	(n=5,007)	(n=4,184)	(n=3,186)
a. Almost always	**	38.9% (±2.6)	28.8% (±2.0)	34.1% (±1.8)
b. Often	**	21.6% (±1.5)	22.7% (±1.7)	25.4% (±1.8)
c. Sometimes	**	16.4% (±1.4)	21.5% (±1.4)	20.2% (±1.8)
d. Once in a while	**	11.5% (±1.0)	14.8% (±1.1)	12.1% (±1.3)
e. Almost never	**	11.5% (±1.2)	12.2% (±1.4)	8.3% (±1.1)

145. If you see bullying or have been bullied at school do you know how to report it?

	(n=0)	(n=5,063)	(n=4,230)	(n=3,206)
a. Yes	**	79.6% (±2.1)	68.5% (±2.1)	72.9% (±1.9)
b. No	**	8.7% (±1.3)	15.1% (±1.2)	13.4% (±1.4)
c. Not sure	**	11.6% (±1.3)	16.4% (±1.7)	13.7% (±1.4)

146. In the past 30 days, how often were you bullied, harassed, or intimidated at school or on your way to or from school because of your race, ethnicity, or national origin or what someone thought it was?

	(n=0)	(n=5,084)	(n=4,259)	(n=3,204)
a. 0 times	**	87.8% (±1.1)	89.0% (±1.1)	91.6% (±0.9)
b. 1 time	**	6.0% (±0.7)	5.3% (±0.6)	4.1% (±0.5)
c. 2-3 times	**	3.1% (±0.5)	3.5% (±0.7)	2.3% (±0.6)
d. About once a week	**	1.0% (±0.3)	0.9% (±0.3)	0.7% (±0.2)
e. Several times a week or more	**	2.1% (±0.4)	1.3% (±0.4)	1.3% (±0.3)

147. In the past 30 days, how often were you bullied, harassed, or intimidated at school or on your way to or from school: Because someone thought you were gay, lesbian, or bisexual (whether you are or are not)?

	(n=0)	(n=5,066)	(n=4,252)	(n=3,203)
a. 0 times	**	88.6% (±1.3)	90.7% (±1.0)	93.2% (±0.8)
b. 1 time	**	5.7% (±0.7)	4.7% (±0.6)	3.3% (±0.5)
c. 2-3 times	**	2.7% (±0.5)	2.4% (±0.5)	1.8% (±0.6)
d. About once a week	**	1.1% (±0.3)	1.1% (±0.3)	0.9% (±0.3)
e. Several times a week or more	**	1.8% (±0.5)	1.2% (±0.4)	0.8% (±0.3)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,953)	% (± CI) (n=4,167)	% (± CI) (n=3,137)
148. During the past 30 days, on how many days did you not go to school because you felt you would be unsafe at school or on your way to and from school?				
a. 0 days	**	91.5% (±1.2)	91.6% (±1.2)	93.1% (±1.3)
b. 1 day	**	3.4% (±0.6)	3.6% (±0.7)	3.2% (±0.8)
c. 2 or 3 days	**	2.5% (±0.5)	2.6% (±0.7)	2.0% (±0.6)
d. 4 or 5 days	**	1.1% (±0.3)	0.9% (±0.3)	0.7% (±0.3)
e. 6 or more days	**	1.6% (±0.4)	1.4% (±0.4)	1.0% (±0.3)
149. In the past 30 days, has someone used the computer or a cell phone to bully, harass or intimidate you?	(n=0)	(n=5,070)	(n=4,245)	(n=3,202)
a. Yes	**	13.2% (±1.0)	12.4% (±1.0)	11.4% (±1.2)
b. No	**	81.4% (±1.3)	83.5% (±1.4)	85.1% (±1.4)
c. Not sure	**	5.4% (±0.7)	4.1% (±0.7)	3.6% (±0.6)
150. Does your school provide a counselor, intervention specialist, or other school staff member for students to discuss problems with alcohol, tobacco, or other drugs?	(n=0)	(n=5,065)	(n=4,135)	(n=3,119)
a. No	**	7.3% (±1.1)	6.6% (±1.0)	8.4% (±1.8)
b. Yes	**	65.2% (±3.7)	56.2% (±5.2)	55.9% (±6.9)
c. Not sure	**	27.5% (±3.3)	37.3% (±4.7)	35.7% (±5.7)
151. Does your school have a counselor?	(n=0)	(n=4,561)	(n=3,964)	(n=3,019)
a. Yes	**	92.7% (±1.5)	94.4% (±1.0)	94.4% (±1.6)
b. No	**	2.1% (±0.5)	2.4% (±0.8)	2.7% (±1.0)
c. Not sure	**	5.2% (±1.3)	3.1% (±0.6)	2.9% (±0.8)
152. In the last year, did you have any contact with the counselor?	(n=0)	(n=4,659)	(n=4,007)	(n=3,053)
a. Yes	**	42.5% (±3.5)	55.0% (±3.1)	74.0% (±2.5)
b. No	**	57.5% (±3.5)	45.0% (±3.1)	26.0% (±2.5)
153. There are people in this school who will help me if I need it?	(n=0)	(n=4,586)	(n=3,969)	(n=3,026)
a. Yes	**	78.8% (±2.0)	77.3% (±2.0)	81.8% (±1.4)
b. No	**	6.6% (±1.0)	7.0% (±0.9)	6.1% (±0.9)
c. Not sure	**	14.6% (±1.7)	15.7% (±1.5)	12.1% (±1.2)
154. Last year in school, were you taught about AIDS or HIV infection?	(n=0)	(n=4,968)	(n=4,170)	(n=3,144)
a. Yes	**	77.8% (±5.9)	72.8% (±6.7)	43.1% (±6.8)
b. No	**	15.1% (±4.5)	19.6% (±5.9)	45.6% (±6.3)
c. Not sure	**	7.1% (±1.8)	7.6% (±1.4)	11.3% (±1.8)
155. Last year in school, were you taught about abstinence (not having sex) to prevent sexually transmitted diseases (STDs) and pregnancy?	(n=0)	(n=4,572)	(n=3,936)	(n=3,022)
a. Yes	**	67.5% (±6.4)	72.8% (±7.3)	41.9% (±4.6)
b. No	**	19.3% (±4.7)	20.3% (±6.2)	47.1% (±4.8)
c. Not sure	**	13.2% (±2.2)	6.9% (±1.6)	11.0% (±1.7)

156. Last year in school, were you taught about ways other than abstinence to prevent sexually transmitted diseases (STDs) and pregnancy?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,516)	% (± CI) (n=3,930)	% (± CI) (n=3,010)
a. Yes	**	66.3% (±6.4)	73.2% (±7.5)	44.5% (±4.8)
b. No	**	18.9% (±4.6)	19.2% (±6.2)	44.3% (±5.1)
c. Not sure	**	14.9% (±2.2)	7.7% (±1.8)	11.2% (±1.8)

Quality of Life

Health-related quality of life is an individual's or group's perceived physical and mental health over time. At the individual level it involves a person's health and health-related conditions; at the community level it involves conditions that influence people's health. This section contains the results of the survey questions related to individual quality of life.

157. Youth Quality of Life (Computed from questions 158-162)	(n=0)	(n=4,867)	(n=4,131)	(n=3,119)
a. Low	**	23.1% (±2.0)	26.5% (±1.5)	27.4% (±1.8)
b. Medium low	**	21.3% (±1.4)	27.9% (±1.4)	24.4% (±2.0)
c. Medium high	**	21.4% (±1.1)	24.7% (±1.7)	22.4% (±1.8)
d. High	**	34.1% (±2.3)	20.9% (±1.7)	25.7% (±2.1)

158. I feel I am getting along with my parents or guardians.	(n=0)	(n=4,851)	(n=4,099)	(n=3,102)
a. 0 not at all true	**	4.1% (±0.7)	3.6% (±0.6)	3.8% (±0.6)
b. 1	**	1.7% (±0.4)	1.9% (±0.4)	2.2% (±0.6)
c. 2	**	1.3% (±0.3)	2.7% (±0.5)	2.6% (±0.6)
d. 3	**	2.4% (±0.4)	3.4% (±0.6)	3.4% (±0.7)
e. 4	**	2.7% (±0.5)	4.0% (±0.7)	4.1% (±0.7)
f. 5	**	5.6% (±0.8)	7.1% (±0.9)	6.4% (±1.0)
g. 6	**	4.4% (±0.5)	5.7% (±0.7)	6.2% (±0.9)
h. 7	**	9.2% (±0.7)	11.5% (±1.0)	12.5% (±1.2)
i. 8	**	12.8% (±1.1)	15.0% (±1.1)	15.4% (±1.5)
j. 9	**	16.3% (±1.4)	16.3% (±1.2)	14.9% (±1.4)
k. 10 completely true	**	39.5% (±2.1)	28.7% (±1.4)	28.4% (±1.9)

159. I look forward to the future.	(n=0)	(n=4,928)	(n=4,154)	(n=3,132)
a. 0 not at all true	**	3.0% (±0.6)	2.9% (±0.7)	2.9% (±0.7)
b. 1	**	1.0% (±0.3)	1.7% (±0.3)	1.2% (±0.5)
c. 2	**	1.3% (±0.4)	1.9% (±0.5)	1.9% (±0.4)
d. 3	**	2.8% (±0.7)	2.1% (±0.5)	2.1% (±0.7)
e. 4	**	2.5% (±0.5)	2.3% (±0.4)	2.4% (±0.5)
f. 5	**	6.3% (±0.9)	6.6% (±0.9)	6.2% (±0.7)
g. 6	**	3.4% (±0.6)	4.9% (±0.8)	4.3% (±0.7)
h. 7	**	7.9% (±0.7)	10.5% (±1.2)	10.0% (±0.9)
i. 8	**	12.0% (±0.8)	13.5% (±1.2)	13.2% (±1.4)
j. 9	**	14.0% (±1.3)	12.3% (±1.1)	12.6% (±1.3)
k. 10 completely true	**	45.9% (±1.7)	41.4% (±1.9)	43.2% (±1.6)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,922)	% (± CI) (n=4,142)	% (± CI) (n=3,133)
160. I feel good about myself.				
a. 0 not at all true	**	4.6% (±0.9)	4.9% (±0.8)	3.2% (±0.6)
b. 1	**	2.1% (±0.5)	2.6% (±0.5)	2.5% (±0.5)
c. 2	**	2.7% (±0.5)	3.2% (±0.5)	3.0% (±0.6)
d. 3	**	4.5% (±0.7)	4.5% (±0.6)	3.7% (±0.7)
e. 4	**	4.2% (±0.6)	4.6% (±0.6)	4.9% (±0.8)
f. 5	**	6.6% (±0.8)	8.7% (±1.0)	8.7% (±1.1)
g. 6	**	5.3% (±0.7)	8.1% (±1.0)	6.9% (±0.9)
h. 7	**	9.0% (±0.9)	12.9% (±1.1)	13.1% (±1.0)
i. 8	**	12.4% (±1.0)	15.4% (±1.1)	16.7% (±1.3)
j. 9	**	15.4% (±1.2)	13.1% (±1.6)	13.6% (±1.2)
k. 10 completely true	**	33.1% (±1.8)	22.0% (±1.8)	23.8% (±1.8)
161. I am satisfied with the way my life is now.	(n=0)	(n=4,911)	(n=4,148)	(n=3,131)
a. 0 not at all true	**	5.5% (±0.8)	6.2% (±0.7)	5.6% (±0.8)
b. 1	**	2.2% (±0.4)	2.7% (±0.6)	2.2% (±0.5)
c. 2	**	2.2% (±0.5)	3.7% (±0.6)	3.8% (±0.7)
d. 3	**	4.9% (±0.7)	4.7% (±0.7)	5.7% (±0.9)
e. 4	**	4.1% (±0.5)	5.5% (±0.8)	5.1% (±0.7)
f. 5	**	6.7% (±0.8)	8.8% (±0.9)	8.0% (±1.1)
g. 6	**	5.6% (±0.6)	8.5% (±0.9)	7.7% (±1.3)
h. 7	**	8.7% (±0.9)	12.2% (±1.1)	12.5% (±1.1)
i. 8	**	11.3% (±1.0)	13.7% (±0.9)	16.3% (±1.6)
j. 9	**	16.0% (±1.2)	13.6% (±1.1)	12.6% (±1.1)
k. 10 completely true	**	32.9% (±1.8)	20.3% (±1.6)	20.5% (±1.7)
162. I feel alone in my life.	(n=0)	(n=4,902)	(n=4,135)	(n=3,130)
a. 0 not at all true	**	48.3% (±2.1)	35.0% (±1.6)	34.4% (±2.2)
b. 1	**	9.5% (±1.1)	10.4% (±1.2)	11.9% (±1.5)
c. 2	**	6.0% (±0.6)	9.1% (±1.0)	10.6% (±1.5)
d. 3	**	5.2% (±0.8)	5.9% (±0.6)	6.0% (±1.0)
e. 4	**	3.9% (±0.6)	4.5% (±0.9)	4.7% (±0.8)
f. 5	**	5.2% (±0.8)	7.5% (±0.8)	7.5% (±1.0)
g. 6	**	3.1% (±0.6)	4.9% (±0.7)	4.6% (±0.8)
h. 7	**	4.7% (±0.6)	6.7% (±0.7)	6.4% (±0.9)
i. 8	**	5.1% (±0.7)	6.5% (±0.9)	6.2% (±1.1)
j. 9	**	3.7% (±0.5)	4.3% (±0.8)	3.7% (±0.7)
k. 10 completely true	**	5.3% (±0.8)	5.3% (±0.7)	4.1% (±0.8)
163. Do you have goals and plans for the future?	(n=8,935)	(n=0)	(n=0)	(n=0)
a. No	12.4% (±0.9)	**	**	**
b. Yes	87.6% (±0.9)	**	**	**

Risk and Protective Factors

Decades of research have shown that certain risk factors are associated with increased likelihood of health risk behaviors including ATOD use, violence, and delinquent behaviors. Similarly, research from Drs. Hawkins and Catalano and associates at the University of Washington, has shown that protective factors exert a positive influence against the negative influence of risk factors. The premise of the risk reduction and protective factor enhancement approach to prevention is that preventing a problem before it occurs requires addressing the factors that predict the problem. Ideally, this strategy entails discovering the causes of the problem behavior and influencing those causes. Many of the survey questions were used to assess students' status on risk and protective factors in the community, school, and peer-individual domains. Composite scales were computed for use in local program planning.

Risk and Protective Factor Framework and Reporting Schedule

This table provides a list of risk and protective factors included in the Healthy Youth Survey by year.

	2002	2004	2006	2008	2010	2012	2014
Community Protective Factors							
Opportunities for Prosocial Involvement	X	X ^S	X ^S				
Rewards for Prosocial Involvement	X	X	X	X	X ^E	X ^E	X ^E
Community Risk Factors							
Transitions and Mobility		X ^S					
Perceived Availability of Handguns	X ^S						
Laws And Norms Favorable to Drug Use	X	X	X	X	X	X	X
Low Neighborhood Attachment	X ^S		X ^S	X ^S	X ^S		
Perceived Availability of Drugs	X	X	X	X	X	X	X
Family Protective Factors							
Opportunities for Prosocial Involvement	X [†]	X ^{E, †}	X [†]	X [†]	X [†]	X [†]	X
Rewards for Prosocial Involvement	X [†]	X ^{E, †}	X [†]	X [†]	X [†]	X [†]	X ^E
Family Risk Factors							
Poor Family Management	X ^{S, †}	X ^S					
Parental Attitudes Favorable towards Drug Use		X ^{S, †}		X ^{S, †}	X ^S	X ^S	X ^S
Parental Attitudes Favorable to Antisocial Behavior		X ^{S, †}					
Antisocial Behavior Among Familiar Adults	X ^S						
Peer-Individual Protective Factors							
Social Skills	X ^S	X ^S					
Interaction With Prosocial Peers		X	X	X	X ^S	X ^S	X ^S
Belief in the Moral Order	X ^S	X ^S					
Prosocial Involvement		X	X	X ^E	X ^E	X ^E	X ^E
Peer-Individual Risk Factors							
Perceived Risk of Drug Use	X	X	X	X	X	X	X
Early Initiation of Drug Use	X ^S	X ^S					
Early Initiation of Antisocial Behavior	X ^S						
Favorable Attitudes Towards Drug Use	X	X	X	X	X	X	X
Favorable Attitudes Towards Antisocial Behavior	X ^S						
Rewards for Antisocial Involvement	X ^S	X ^S	X ^S	X ^S			
Friends' Use of Drugs	X ^S	X ^S					
Interaction With Antisocial Peers		X ^S					
Intentions to Use	X ^S						
School Protective Factors							
Opportunities for Prosocial Involvement	X ^S	X ^S					
Rewards for Prosocial Involvement	X	X	X	X	X	X	X
School Risk Factors							
Low Commitment to School	X	X	X	X	X	X	X
Academic Failure	X	X	X	X	X	X	X


Note: X^S = Included only on the secondary version; X^E = Included only on the elementary version.

Risk and Protective Factor Scale Results and Graphs

For each risk factor scale, the percentage of students who are at risk (i.e., who agreed with statements that predict ATOD use or other problem behaviors) is reported; higher percentages indicate that more students are likely to engage in problem behaviors. For each protective factor scale, the percentage of students who are resilient (i.e., who agreed with statements that predict the ability to resist ATOD use or other problem behaviors) is reported; higher percentages indicate that fewer students are likely to engage in problem behaviors. These percentages are based on computational methods provided by the University of Washington's Social Development Research Group.

Community Domain

Risk Factors	Grade 6 % (±CI) (n=8,312)	Grade 8 % (±CI) (n=5,249)	Grade 10 % (±CI) (n=4,294)	Grade 12 % (±CI) (n=3,255)
Perceived Availability of Drugs (Questions 148-151)	18.7% (±1.3)	17.1% (±1.8)	26.5% (±2.2)	31.7% (±2.1)
Laws And Norms Favorable to Drug Use (Questions 153-158)	33.9% (±1.8)	23.1% (±2.1)	31.7% (±3.1)	31.2% (±2.5)
Protective Factors				
Opportunities for Prosocial Involvement (Questions 159-162)	NA ^S	(n=5,150) 75.3% (±3.2)	(n=4,229) 75.4% (±3.2)	(n=3,225) 77.7% (±3.7)
Rewards for Prosocial Involvement (Questions 163-165)	(n=8,794) 36.9% (±1.8)	NA ^E	NA ^E	NA ^E




Note: NA^S = Included only on the secondary version; NA^E = Included only on the elementary version.

For more information, see the History of Risk and Protective Factor Report on www.AskHYS.net, under HYS Results – Frequency Reports - Additional Reports

Family Domain
Risk Factors

	Grade 6 % (±CI)	Grade 8 % (±CI)	Grade 10 % (±CI)	Grade 12 % (±CI)
Poor Family Management (Questions 166-173)	NA ^S	(n=4,599) 30.6% (±2.4)	(n=3,875) 32.8% (±2.1)	(n=2,969) 34.4% (±2.5)
Parental Attitudes Favorable Towards Drug Use (Questions 181-183)	NA ^S	(n=5,330) 22.2% (±2.1)	(n=4,342) 40.5% (±2.3)	(n=3,281) 41.3% (±2.8)
Protective Factors				
Opportunities for Prosocial Involvement (Questions 174-176)	(n=8,223) 52.6% (±2.1)	(n=4,656) 68.4% (±2.1)	(n=3,905) 57.8% (±2.6)	(n=2,980) 57.8% (±2.6)
Rewards for Prosocial Involvement (Questions 177-180)	(n=8,141) 53.6% (±2.3)	NA ^E	NA ^E	NA ^E



School Domain

Risk Factors

Academic Failure (Questions 184–185)

Grade 6	Grade 8	Grade 10	Grade 12
% (±CI)	% (±CI)	% (±CI)	% (±CI)
(n=8,275)	(n=5,090)	(n=4,220)	(n=3,190)
39.5% (±1.9)	43.9% (±2.3)	46.1% (±2.6)	49.2% (±2.9)

Low Commitment to School (Questions 186–192)

(n=9,001)	(n=5,312)	(n=4,306)	(n=3,252)
38.1% (±1.9)	31.9% (±1.9)	38.3% (±2.8)	40.4% (±2.7)


Protective Factors

Opportunities for Prosocial Involvement (Questions 193–197)

NA ^s	(n=5,259)	(n=4,273)	(n=3,223)
	70.0% (±2.4)	65.2% (±2.7)	68.3% (±3.2)

Rewards for Prosocial Involvement (Questions 198–201)


(n=8,962)	(n=5,246)	(n=4,267)	(n=3,216)
44.9% (±2.0)	52.8% (±2.9)	57.5% (±2.5)	43.3% (±2.7)




Peer-Individual Domain

Risk Factors

	Grade 6 % (±CI) (n=7,244)	Grade 8 % (±CI) (n=4,912)	Grade 10 % (±CI) (n=4,109)	Grade 12 % (±CI) (n=3,116)
Perceived Risk of Drug Use (Questions 202–205)	40.7% (±2.5)	37.9% (±3.5)	41.2% (±2.9)	52.8% (±2.5)
Early Initiation of Drug Use (Questions 206-209)	NA ^S	13.7% (±2.0) (n=5,048)	20.5% (±2.4) (n=4,120)	22.8% (±2.8) (n=3,099)
Favorable Attitudes Towards Drug Use (Questions 216-219)	19.6% (±1.3) (n=8,387)	23.8% (±2.4) (n=4,870)	41.0% (±2.3) (n=4,022)	39.9% (±2.9) (n=3,037)
Friends' Use of Drugs (Questions 221-224)	NA ^S	15.3% (±2.4) (n=4,248)	23.0% (±2.2) (n=3,689)	22.5% (±2.3) (n=2,872)
Protective Factors				
Interaction With Prosocial Peers (Questions 234-238)	NA ^S	61.5% (±2.8) (n=4,249)	57.7% (±3.3) (n=3,705)	50.9% (±3.5) (n=2,865)
Belief in the Moral Order (Questions 239-242)	NA ^S	71.8% (±2.5) (n=5,164)	72.9% (±2.1) (n=4,191)	55.4% (±2.4) (n=3,157)
Opportunities for Prosocial Involvement (Questions 243-245)	40.9% (±2.5) (n=8,858)	NA ^E	NA ^E	NA ^E
Social Skills (Questions 246-248)	NA ^S	73.8% (±2.3) (n=4,212)	61.1% (±2.1) (n=3,677)	54.9% (±2.4) (n=2,851)



Peer-Individual Risk Factors



Peer-Individual Protective Factors

Risk and Protective Factors: Individual Question Results

The remainder of the report provides results for the individual survey questions used to compute the risk and protective factors.

Community Domain

Perceived Availability of Drugs (Questions 164-167)

	Grade 6	Grade 8	Grade 10	Grade 12
164. If you wanted to get some beer, wine, or hard liquor (for example vodka, whiskey, or gin), how easy would it be for you to get some?	% (± CI) (n=8,359)	% (± CI) (n=5,249)	% (± CI) (n=4,305)	% (± CI) (n=3,258)
a. Very hard	73.0% (±1.1)	48.2% (±1.8)	22.0% (±1.7)	13.8% (±1.6)
b. Sort of hard	14.4% (±0.9)	23.7% (±1.3)	26.5% (±1.4)	22.8% (±1.7)
c. Sort of easy	7.5% (±0.5)	16.8% (±1.0)	28.3% (±1.5)	31.7% (±1.3)
d. Very easy	5.1% (±0.6)	11.3% (±1.1)	23.2% (±1.5)	31.6% (±2.0)
165. If you wanted to get some cigarettes, how easy would it be for you to get some?	(n=8,339)	(n=5,243)	(n=4,290)	(n=3,261)
a. Very hard	76.7% (±1.4)	60.4% (±2.2)	33.6% (±2.0)	18.0% (±1.6)
b. Sort of hard	12.2% (±0.8)	18.5% (±1.2)	24.9% (±1.7)	19.3% (±1.9)
c. Sort of easy	6.0% (±0.6)	10.9% (±1.0)	20.5% (±1.4)	21.8% (±1.5)
d. Very easy	5.1% (±0.7)	10.1% (±1.2)	21.0% (±2.4)	40.9% (±2.7)
166. If you wanted to get some marijuana, how easy would it be for you to get some?	(n=8,277)	(n=5,240)	(n=4,290)	(n=3,254)
a. Very hard	86.0% (±1.4)	65.8% (±2.9)	29.1% (±2.4)	17.1% (±1.5)
b. Sort of hard	7.2% (±0.7)	13.6% (±1.1)	18.0% (±1.3)	17.1% (±1.8)
c. Sort of easy	3.4% (±0.5)	9.7% (±1.2)	20.5% (±1.3)	23.7% (±1.6)
d. Very easy	3.5% (±0.6)	10.9% (±1.6)	32.4% (±2.5)	42.1% (±2.6)
167. If you wanted to get a drug like cocaine, LSD, or amphetamines, how easy would it be for you to get some?	(n=8,255)	(n=5,246)	(n=4,283)	(n=3,249)
a. Very hard	90.3% (±1.1)	84.4% (±1.6)	61.8% (±2.5)	50.8% (±2.3)
b. Sort of hard	5.8% (±0.6)	9.4% (±1.1)	22.3% (±1.7)	25.6% (±1.7)
c. Sort of easy	1.8% (±0.4)	4.1% (±0.6)	11.3% (±1.0)	16.0% (±1.8)
d. Very easy	2.1% (±0.4)	2.1% (±0.4)	4.6% (±0.8)	7.6% (±1.1)

Laws and Norms Favorable to Drug Use (Questions 168-173)

	Grade 6	Grade 8	Grade 10	Grade 12
168. How wrong would most adults in your neighborhood^C/ neighborhood or community^A think it was for kids your age: To use marijuana?	(n=8,620)	(n=5,237)	(n=4,290)	(n=3,248)
a. Very wrong	82.9% (±1.5)	65.5% (±2.6)	42.0% (±2.8)	32.5% (±2.7)
b. Wrong	10.7% (±0.9)	23.0% (±1.6)	37.6% (±1.7)	40.2% (±2.2)
c. A little bit wrong	3.9% (±0.6)	8.2% (±1.2)	15.6% (±1.9)	21.0% (±2.1)
d. Not wrong at all	2.5% (±0.4)	3.4% (±0.6)	4.8% (±0.9)	6.3% (±1.2)

169. How wrong would most adults in your neighborhood ^C / neighborhood or community ^A think it was for kids your age: To drink alcohol?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,715)	% (± CI) (n=5,289)	% (± CI) (n=4,327)	% (± CI) (n=3,276)
a. Very wrong	79.6% (±1.1)	60.3% (±2.4)	38.2% (±2.2)	28.9% (±1.8)
b. Wrong	13.7% (±0.8)	28.9% (±1.7)	41.7% (±1.7)	42.3% (±1.8)
c. A little bit wrong	4.7% (±0.5)	8.2% (±0.9)	16.2% (±1.6)	22.3% (±1.8)
d. Not wrong at all	2.0% (±0.4)	2.6% (±0.5)	4.0% (±0.8)	6.5% (±1.0)

170. How wrong would most adults in your neighborhood ^C / neighborhood or community ^A think it was for kids your age: To smoke cigarettes?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=8,689)	(n=5,285)	(n=4,323)	(n=3,275)
a. Very wrong	81.0% (±1.3)	67.6% (±2.2)	51.8% (±2.6)	43.2% (±2.8)
b. Wrong	12.1% (±0.8)	23.4% (±1.6)	32.4% (±1.6)	32.5% (±2.3)
c. A little bit wrong	4.4% (±0.6)	6.5% (±0.9)	11.1% (±1.4)	15.9% (±1.5)
d. Not wrong at all	2.5% (±0.4)	2.5% (±0.5)	4.7% (±0.9)	8.4% (±1.6)

171. If a kid drank some beer, wine, or hard liquor (for example vodka, whiskey, or gin) in your neighborhood ^C / community ^A , would he or she be caught by the police?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=8,519)	(n=5,227)	(n=4,293)	(n=3,266)
a. NO!	9.1% (±0.9)	9.1% (±1.0)	15.1% (±1.7)	19.1% (±1.8)
b. No	27.5% (±1.3)	40.1% (±1.7)	58.4% (±2.0)	61.8% (±2.3)
c. yes	34.2% (±1.2)	35.3% (±1.3)	21.5% (±1.8)	14.9% (±1.5)
d. YES!	29.2% (±1.3)	15.5% (±1.3)	5.0% (±0.6)	4.2% (±0.9)

172. If a kid carried a handgun in your neighborhood ^C / community ^A , would he or she be caught by the police?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=8,595)	(n=5,235)	(n=4,274)	(n=3,256)
a. NO!	6.5% (±0.7)	6.6% (±0.7)	6.3% (±0.9)	7.5% (±1.4)
b. No	11.8% (±1.1)	13.9% (±1.5)	23.4% (±1.7)	27.2% (±1.9)
c. yes	28.2% (±1.2)	33.3% (±1.5)	40.8% (±1.7)	41.8% (±2.2)
d. YES!	53.5% (±1.8)	46.2% (±2.3)	29.6% (±1.9)	23.5% (±1.5)

173. If a kid used marijuana in your neighborhood ^C / community ^A , would he or she be caught by the police?	Grade 6	Grade 8	Grade 10	Grade 12
	(n=8,450)	(n=5,231)	(n=4,292)	(n=3,265)
a. NO!	7.6% (±0.9)	9.9% (±1.1)	17.7% (±1.9)	21.9% (±1.9)
b. No	19.6% (±1.1)	31.4% (±2.0)	52.4% (±1.8)	57.9% (±2.3)
c. yes	35.5% (±1.0)	34.0% (±1.4)	21.9% (±1.6)	15.4% (±1.3)
d. YES!	37.3% (±1.4)	24.6% (±2.1)	7.9% (±0.7)	4.8% (±1.0)

Opportunities for Prosocial Involvement (Questions 174-177)

174. There are adults in my neighborhood or community I could talk to about something important.	Grade 6	Grade 8	Grade 10	Grade 12
	(n=0)	(n=5,290)	(n=4,302)	(n=3,264)
a. NO!	**	9.1% (±1.2)	9.2% (±1.5)	8.6% (±1.7)
b. No	**	13.4% (±1.3)	16.4% (±1.3)	15.7% (±2.0)
c. yes	**	37.5% (±1.5)	42.4% (±1.4)	43.5% (±1.6)
d. YES!	**	40.0% (±2.1)	32.0% (±2.2)	32.2% (±2.8)

Which of the following activities for people your age are available in your community?

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,262)	% (± CI) (n=4,297)	% (± CI) (n=3,264)
175. Sports teams and recreation				
a. Yes	**	88.0% (±2.0)	89.4% (±1.8)	90.6% (±2.1)
b. No	**	12.0% (±2.0)	10.6% (±1.8)	9.4% (±2.1)
176. Scouts, Camp Fire, 4-H Clubs, or other service clubs	(n=0)	(n=5,221)	(n=4,265)	(n=3,243)
a. Yes	**	69.6% (±4.0)	71.2% (±4.6)	75.6% (±4.7)
b. No	**	30.4% (±4.0)	28.8% (±4.6)	24.4% (±4.7)
177. Boys and Girls Club, YMCA, or other activity clubs	(n=0)	(n=5,227)	(n=4,277)	(n=3,250)
a. Yes	**	76.4% (±3.7)	79.2% (±3.3)	80.4% (±4.4)
b. No	**	23.6% (±3.7)	20.8% (±3.3)	19.6% (±4.4)

Rewards for Prosocial Involvement (Questions 178-180)

178. My neighbors notice when I am doing a good job and let me know.

	(n=8,759)	(n=0)	(n=0)	(n=0)
a. NO!	35.3% (±1.6)	**	**	**
b. No	38.9% (±1.1)	**	**	**
c. yes	19.1% (±1.1)	**	**	**
d. YES!	6.6% (±0.6)	**	**	**

179. There are people in my neighborhood who encourage me to do my best.

	(n=8,758)	(n=0)	(n=0)	(n=0)
a. NO!	25.4% (±1.6)	**	**	**
b. No	29.9% (±1.1)	**	**	**
c. yes	31.8% (±1.3)	**	**	**
d. YES!	12.9% (±0.9)	**	**	**

180. There are people in my neighborhood who are proud of me when I do something well.

	(n=8,798)	(n=0)	(n=0)	(n=0)
a. NO!	24.9% (±1.6)	**	**	**
b. No	31.6% (±1.1)	**	**	**
c. yes	32.4% (±1.4)	**	**	**
d. YES!	11.1% (±0.8)	**	**	**

Family Domain

Poor Family Management (Questions 181-188)

181. My parents ask if I've gotten my homework done.

	(n=0)	(n=4,746)	(n=3,943)	(n=2,996)
a. NO!	**	3.7% (±0.6)	4.9% (±0.8)	7.3% (±1.0)
b. No	**	6.2% (±0.8)	10.2% (±1.0)	18.2% (±1.8)
c. yes	**	31.9% (±1.8)	40.9% (±2.2)	41.9% (±2.0)
d. YES!	**	58.2% (±2.5)	44.0% (±2.4)	32.6% (±1.9)

	Grade 6	Grade 8	Grade 10	Grade 12
182. Would your parents know if you did not come home on time?	% (± CI) (n=0)	% (± CI) (n=4,606)	% (± CI) (n=3,880)	% (± CI) (n=2,970)
a. NO!	**	3.1% (±0.6)	4.6% (±0.8)	5.8% (±0.9)
b. No	**	10.8% (±1.2)	14.4% (±1.1)	17.2% (±1.5)
c. yes	**	34.3% (±1.6)	43.1% (±1.8)	45.7% (±1.7)
d. YES!	**	51.8% (±2.3)	37.8% (±2.0)	31.3% (±1.8)
183. When I am not at home, one of my parents knows where I am and who I am with.	(n=0)	(n=4,740)	(n=3,937)	(n=2,993)
a. NO!	**	3.2% (±0.7)	3.3% (±0.6)	5.0% (±1.0)
b. No	**	5.1% (±0.8)	8.1% (±1.1)	11.0% (±1.2)
c. yes	**	32.3% (±1.3)	45.9% (±2.1)	50.7% (±2.0)
d. YES!	**	59.3% (±2.1)	42.8% (±2.2)	33.4% (±1.5)
184. The rules in my family are clear.	(n=0)	(n=4,687)	(n=3,920)	(n=2,986)
a. NO!	**	2.9% (±0.5)	3.5% (±0.6)	3.7% (±0.9)
b. No	**	8.2% (±0.9)	10.4% (±1.1)	11.5% (±1.3)
c. yes	**	35.6% (±1.7)	47.2% (±1.7)	48.8% (±2.1)
d. YES!	**	53.3% (±2.2)	38.8% (±2.2)	36.0% (±1.6)
185. My family has clear rules about alcohol and drug use.	(n=0)	(n=4,656)	(n=3,907)	(n=2,983)
a. NO!	**	3.0% (±0.5)	3.2% (±0.6)	4.1% (±1.0)
b. No	**	7.7% (±0.9)	12.6% (±1.2)	15.4% (±1.7)
c. yes	**	24.3% (±1.7)	35.1% (±2.1)	40.7% (±1.4)
d. YES!	**	64.9% (±2.3)	49.1% (±2.9)	39.8% (±1.9)
186. If you drank some beer, wine, or liquor (for example vodka, whiskey, or gin) without your parent's permission, would you be caught by them?	(n=0)	(n=4,576)	(n=3,871)	(n=2,969)
a. NO!	**	6.1% (±1.0)	10.3% (±1.4)	14.4% (±1.4)
b. No	**	17.3% (±1.7)	36.0% (±2.4)	44.8% (±2.4)
c. yes	**	27.6% (±1.5)	29.7% (±1.5)	25.8% (±1.8)
d. YES!	**	49.0% (±2.3)	24.0% (±1.8)	15.0% (±1.5)
187. If you carried a handgun without your parent's permission, would you be caught by them?	(n=0)	(n=4,566)	(n=3,853)	(n=2,950)
a. NO!	**	3.9% (±0.7)	4.9% (±0.8)	6.8% (±1.2)
b. No	**	6.1% (±0.7)	13.3% (±1.4)	17.7% (±1.8)
c. yes	**	23.9% (±1.6)	32.0% (±1.6)	33.7% (±1.8)
d. YES!	**	66.1% (±2.0)	49.8% (±1.8)	41.8% (±1.9)
188. If you skipped school, would you be caught by your parents?	(n=0)	(n=4,576)	(n=3,871)	(n=2,966)
a. NO!	**	4.7% (±0.7)	5.4% (±0.9)	7.9% (±1.1)
b. No	**	5.3% (±1.0)	11.1% (±1.8)	14.8% (±1.9)
c. yes	**	22.9% (±1.6)	32.8% (±1.6)	39.8% (±1.6)
d. YES!	**	67.0% (±2.5)	50.7% (±2.7)	37.6% (±2.8)

Opportunities for Prosocial Involvement (Questions 189-191)

	Grade 6	Grade 8	Grade 10	Grade 12
189. If I had a personal problem, I could ask my mom or dad for help.	% (± CI) (n=8,199)	% (± CI) (n=4,692)	% (± CI) (n=3,919)	% (± CI) (n=2,984)
a. NO!	5.6% (±0.7)	6.5% (±1.0)	8.2% (±1.2)	7.4% (±1.3)
b. No	7.0% (±0.8)	9.6% (±0.9)	13.5% (±1.2)	12.7% (±1.6)
c. yes	30.9% (±1.2)	32.4% (±1.4)	42.2% (±2.2)	43.8% (±1.7)
d. YES!	56.4% (±1.9)	51.5% (±2.1)	36.1% (±2.1)	36.1% (±2.2)
190. My parents give me lots of chances to do fun things with them.	(n=8,254)	(n=4,644)	(n=3,901)	(n=2,978)
a. NO!	5.3% (±0.8)	6.1% (±0.8)	7.5% (±0.9)	7.5% (±1.2)
b. No	12.7% (±0.9)	15.5% (±1.3)	22.0% (±1.6)	22.1% (±1.4)
c. yes	41.9% (±1.1)	37.9% (±1.5)	44.7% (±1.4)	46.1% (±1.7)
d. YES!	40.1% (±1.7)	40.5% (±2.0)	25.8% (±1.8)	24.3% (±1.4)
191. My parents ask me what I think before most family decisions affecting me are made.	(n=8,102)	(n=4,598)	(n=3,882)	(n=2,973)
a. NO!	10.3% (±1.0)	10.9% (±1.2)	11.9% (±1.4)	12.2% (±1.7)
b. No	21.4% (±1.4)	20.6% (±1.4)	26.1% (±1.7)	26.4% (±1.7)
c. yes	40.0% (±1.1)	36.9% (±1.4)	41.2% (±1.9)	41.5% (±2.2)
d. YES!	28.3% (±1.7)	31.6% (±1.9)	20.8% (±1.5)	19.9% (±1.6)

Rewards for Prosocial Involvement (Questions 192-195)

192. My parents notice when I am doing a good job and let me know about it.	(n=8,189)	(n=0)	(n=0)	(n=0)
a. Never or almost never	7.8% (±1.0)	**	**	**
b. Sometimes	21.9% (±1.3)	**	**	**
c. Often	30.1% (±1.0)	**	**	**
d. All the time	40.2% (±1.9)	**	**	**
193. How often do your parents tell you they're proud of you for something you've done?	(n=8,115)	(n=0)	(n=0)	(n=0)
a. Never or almost never	6.3% (±0.9)	**	**	**
b. Sometimes	20.0% (±1.2)	**	**	**
c. Often	32.9% (±1.2)	**	**	**
d. All the time	40.8% (±1.9)	**	**	**
194. Do you enjoy spending time with your mom?	(n=8,049)	(n=0)	(n=0)	(n=0)
a. NO!	5.7% (±0.9)	**	**	**
b. No	4.6% (±0.5)	**	**	**
c. yes	23.7% (±1.3)	**	**	**
d. YES!	66.1% (±2.1)	**	**	**
195. Do you enjoy spending time with your dad?	(n=8,143)	(n=0)	(n=0)	(n=0)
a. NO!	2.8% (±0.7)	**	**	**
b. No	3.2% (±0.4)	**	**	**
c. yes	23.8% (±1.4)	**	**	**
d. YES!	70.2% (±1.7)	**	**	**

Parental Attitudes Favorable Towards Drug Use (Questions 196-198)

196. How wrong do your parents feel it would be for you to: Drink beer, wine, or hard liquor (for example vodka, whiskey or gin) regularly (at least once or twice a month)?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,331)	% (± CI) (n=4,339)	% (± CI) (n=3,283)
a. Very wrong	**	84.5% (±1.4)	69.0% (±2.1)	54.1% (±2.2)
b. Wrong	**	10.7% (±1.0)	18.9% (±1.4)	22.1% (±1.6)
c. A little bit wrong	**	3.4% (±0.6)	8.9% (±1.0)	15.8% (±1.5)
d. Not wrong at all	**	1.3% (±0.3)	3.2% (±0.8)	8.0% (±1.1)

197. How wrong do your parents feel it would be for you to: Smoke cigarettes?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,328)	% (± CI) (n=4,342)	% (± CI) (n=3,280)
a. Very wrong	**	91.2% (±1.1)	85.6% (±1.7)	78.9% (±2.2)
b. Wrong	**	6.6% (±0.9)	10.3% (±1.2)	13.0% (±1.3)
c. A little bit wrong	**	1.6% (±0.3)	2.5% (±0.6)	4.7% (±0.9)
d. Not wrong at all	**	0.6% (±0.2)	1.6% (±0.5)	3.4% (±1.0)

198. How wrong do your parents feel it would be for you to: Use marijuana?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,322)	% (± CI) (n=4,337)	% (± CI) (n=3,279)
a. Very wrong	**	87.6% (±1.7)	73.0% (±2.3)	64.7% (±2.7)
b. Wrong	**	7.8% (±1.1)	15.8% (±1.2)	18.2% (±1.6)
c. A little bit wrong	**	2.9% (±0.6)	6.9% (±0.9)	10.5% (±1.2)
d. Not wrong at all	**	1.7% (±0.5)	4.2% (±0.9)	6.6% (±1.1)

School Domain

Academic Failure Questions (Questions 199-200)

199. Putting them all together, what were your grades like last year?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,578)	% (± CI) (n=10,182)	% (± CI) (n=8,433)	% (± CI) (n=6,370)
a. Mostly As	43.0% (±2.8)	47.6% (±3.7)	41.8% (±3.9)	35.3% (±3.9)
b. Mostly Bs	42.1% (±2.0)	32.7% (±2.3)	31.7% (±1.5)	34.9% (±1.4)
c. Mostly Cs	11.4% (±1.2)	13.5% (±1.5)	17.7% (±1.9)	22.1% (±2.5)
d. Mostly Ds	2.2% (±0.5)	3.8% (±0.5)	5.4% (±0.9)	5.1% (±0.9)
e. Mostly Fs	1.2% (±0.3)	2.4% (±0.4)	3.3% (±0.7)	2.6% (±0.6)

200. Are your school grades better than the grades of most students in your class?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,625)	% (± CI) (n=5,128)	% (± CI) (n=4,234)	% (± CI) (n=3,195)
a. NO!	5.4% (±0.8)	7.2% (±0.9)	7.4% (±0.9)	7.3% (±0.9)
b. No	30.7% (±1.4)	31.3% (±1.5)	31.6% (±1.8)	35.1% (±2.2)
c. yes	53.1% (±1.7)	48.5% (±1.5)	47.5% (±1.8)	45.2% (±2.0)
d. YES!	10.8% (±1.0)	12.9% (±1.2)	13.4% (±1.1)	12.4% (±1.2)

Low Commitment to School (Questions 201-207)

201. How often do you feel the schoolwork you are assigned is meaningful and important?	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,960)	% (± CI) (n=5,319)	% (± CI) (n=4,318)	% (± CI) (n=3,264)
a. Almost always	31.1% (±1.3)	24.9% (±1.5)	13.2% (±1.4)	10.8% (±1.5)
b. Often	30.1% (±1.0)	30.0% (±1.5)	27.3% (±1.6)	26.2% (±2.3)
c. Sometimes	27.3% (±1.3)	30.5% (±1.4)	36.1% (±1.5)	35.5% (±2.0)
d. Seldom	7.5% (±0.7)	9.4% (±0.8)	16.9% (±1.6)	20.6% (±1.7)
e. Never	4.1% (±0.6)	5.2% (±0.8)	6.5% (±1.0)	6.8% (±1.1)

	Grade 6	Grade 8	Grade 10	Grade 12
202. How interesting are most of your courses to you?	% (± CI) (n=8,726)	% (± CI) (n=5,290)	% (± CI) (n=4,314)	% (± CI) (n=3,257)
a. Very interesting and stimulating	14.7% (±1.0)	10.1% (±1.0)	6.5% (±1.2)	9.9% (±1.3)
b. Quite interesting	39.4% (±1.5)	32.0% (±1.5)	26.0% (±1.8)	28.9% (±1.9)
c. Fairly interesting	32.7% (±1.4)	35.7% (±1.5)	39.0% (±1.7)	37.7% (±1.4)
d. Slightly dull	9.4% (±0.8)	15.3% (±1.1)	20.2% (±1.5)	16.3% (±1.4)
e. Very dull	3.7% (±0.5)	7.0% (±0.9)	8.2% (±1.1)	7.2% (±1.1)
203. How important do you think the things you are learning in school are going to be for you later in life?	(n=8,959)	(n=5,317)	(n=4,307)	(n=3,256)
a. Very important	52.9% (±2.1)	36.1% (±1.9)	19.4% (±2.0)	16.9% (±1.6)
b. Quite important	26.6% (±1.2)	28.5% (±1.6)	25.6% (±1.6)	24.3% (±2.0)
c. Fairly important	13.5% (±1.1)	21.5% (±1.3)	28.3% (±1.4)	29.3% (±1.5)
d. Slightly important	5.5% (±0.5)	10.9% (±0.9)	20.6% (±1.3)	22.7% (±1.8)
e. Not at all important	1.5% (±0.3)	2.9% (±0.5)	6.1% (±0.8)	6.8% (±1.2)
Think back over the past year in school. How often did you:				
204. Enjoy being in school?	(n=9,066)	(n=10,284)	(n=8,480)	(n=6,398)
a. Never	4.7% (±0.6)	8.7% (±0.8)	10.0% (±0.8)	10.0% (±1.2)
b. Seldom	5.7% (±0.5)	12.1% (±0.7)	17.2% (±1.0)	17.6% (±1.1)
c. Sometimes	31.3% (±1.4)	32.7% (±1.1)	35.3% (±1.1)	35.9% (±1.6)
d. Often	27.7% (±1.3)	28.9% (±1.3)	26.7% (±1.4)	26.5% (±1.9)
e. Almost always	30.6% (±1.2)	17.6% (±1.1)	10.8% (±1.1)	10.0% (±0.9)
205. Hate being in school?	(n=9,041)	(n=5,303)	(n=4,305)	(n=3,252)
a. Never	23.0% (±1.2)	15.0% (±1.2)	8.0% (±0.9)	8.4% (±1.1)
b. Seldom	27.6% (±1.4)	27.5% (±1.7)	26.0% (±1.8)	25.2% (±2.0)
c. Sometimes	36.2% (±1.2)	36.4% (±1.7)	36.9% (±1.5)	37.8% (±2.0)
d. Often	7.6% (±0.7)	13.1% (±1.0)	18.6% (±1.4)	18.7% (±1.6)
e. Almost always	5.6% (±0.8)	8.0% (±1.0)	10.5% (±1.4)	9.9% (±1.6)
206. Try to do your best work in school?	(n=8,984)	(n=5,302)	(n=4,295)	(n=3,246)
a. Never	1.7% (±0.3)	2.7% (±0.4)	1.8% (±0.4)	1.9% (±0.5)
b. Seldom	1.3% (±0.2)	4.3% (±0.6)	5.6% (±0.9)	5.8% (±0.8)
c. Sometimes	6.0% (±0.6)	10.0% (±1.1)	14.8% (±1.1)	18.1% (±1.3)
d. Often	18.9% (±1.1)	27.6% (±1.4)	35.0% (±1.2)	34.3% (±1.8)
e. Almost always	72.1% (±1.4)	55.5% (±2.2)	42.8% (±2.0)	39.9% (±1.9)
207. During the LAST 4 WEEKS, how many whole days of school have you missed because you skipped or "cut"?	(n=9,020)	(n=5,300)	(n=4,297)	(n=3,248)
a. None	83.2% (±1.4)	84.7% (±1.5)	82.0% (±1.8)	71.6% (±3.2)
b. 1	8.1% (±0.8)	6.6% (±0.7)	7.7% (±1.0)	10.6% (±1.6)
c. 2	3.8% (±0.6)	3.5% (±0.6)	3.2% (±0.7)	6.0% (±1.0)
d. 3	2.3% (±0.3)	2.0% (±0.4)	2.8% (±0.6)	4.2% (±0.8)
e. 4-5	1.7% (±0.3)	1.5% (±0.4)	2.4% (±0.7)	4.3% (±0.9)
f. 6-10	0.6% (±0.1)	0.9% (±0.3)	0.9% (±0.3)	2.3% (±0.7)
g. 11 or more	0.3% (±0.1)	0.8% (±0.3)	1.0% (±0.3)	1.0% (±0.3)

Opportunities for Prosocial Involvement (Questions 208-212)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=5,245)	% (± CI) (n=4,263)	% (± CI) (n=3,223)
208. In my school, students have lots of chances to help decide things like class activities and rules.				
a. NO!	**	14.1% (±1.7)	14.7% (±1.6)	14.5% (±1.5)
b. No	**	29.2% (±1.7)	35.4% (±1.7)	34.3% (±2.6)
c. yes	**	44.8% (±1.7)	41.6% (±2.1)	42.7% (±2.3)
d. YES!	**	11.9% (±1.6)	8.3% (±1.2)	8.5% (±1.5)
209. There are lots of chances for students in my school to talk with a teacher one-on-one.	(n=0)	(n=5,268)	(n=4,283)	(n=3,236)
a. NO!	**	5.3% (±0.7)	4.4% (±0.7)	3.8% (±0.9)
b. No	**	13.8% (±1.4)	15.7% (±1.5)	12.5% (±1.7)
c. yes	**	48.2% (±1.6)	55.8% (±1.5)	55.3% (±2.4)
d. YES!	**	32.6% (±2.4)	24.1% (±2.4)	28.4% (±2.9)
210. Teachers ask me to work on special classroom projects.	(n=0)	(n=5,202)	(n=4,248)	(n=3,210)
a. NO!	**	14.8% (±1.3)	15.4% (±1.5)	13.4% (±1.3)
b. No	**	41.5% (±1.8)	51.0% (±1.7)	49.9% (±1.7)
c. yes	**	34.3% (±1.6)	28.7% (±1.9)	31.7% (±2.0)
d. YES!	**	9.5% (±0.8)	4.8% (±0.8)	5.0% (±0.8)
211. There are lots of chances for students in my school to get involved in sports, clubs, and other school activities outside of class.	(n=0)	(n=5,279)	(n=4,283)	(n=3,234)
a. NO!	**	2.6% (±0.5)	2.2% (±0.5)	2.7% (±0.7)
b. No	**	4.4% (±0.8)	4.6% (±0.9)	5.6% (±1.5)
c. yes	**	35.6% (±2.5)	36.0% (±2.4)	37.0% (±2.9)
d. YES!	**	57.4% (±3.1)	57.2% (±3.0)	54.7% (±4.0)
212. I have lots of chances to be part of class discussions or activities.	(n=0)	(n=5,267)	(n=4,281)	(n=3,226)
a. NO!	**	3.6% (±0.6)	2.7% (±0.5)	2.7% (±0.6)
b. No	**	10.4% (±1.2)	11.0% (±1.2)	9.2% (±1.6)
c. yes	**	50.2% (±2.0)	58.3% (±1.7)	58.5% (±2.0)
d. YES!	**	35.8% (±2.6)	28.0% (±2.2)	29.7% (±3.0)

Rewards for Prosocial Involvement (Questions 213-216)

213. My teacher(s) notices when I am doing a good job and lets me know about it.	(n=9,036)	(n=5,253)	(n=4,264)	(n=3,225)
a. NO!	7.0% (±0.8)	8.5% (±1.1)	9.8% (±1.1)	6.9% (±1.1)
b. No	20.6% (±1.3)	24.7% (±1.7)	32.3% (±2.2)	29.9% (±1.9)
c. yes	54.2% (±1.4)	49.3% (±1.6)	47.3% (±1.7)	51.9% (±1.7)
d. YES!	18.2% (±1.3)	17.4% (±1.5)	10.6% (±1.6)	11.3% (±1.5)
214. The school lets my parents know when I have done something well.	(n=8,865)	(n=5,243)	(n=4,266)	(n=3,214)
a. NO!	15.2% (±1.1)	22.4% (±1.9)	27.9% (±1.7)	27.0% (±2.2)
b. No	40.2% (±1.6)	41.3% (±1.7)	47.2% (±1.7)	49.8% (±2.1)
c. yes	33.2% (±1.3)	27.1% (±2.1)	20.7% (±1.6)	19.0% (±2.0)
d. YES!	11.4% (±0.9)	9.2% (±1.0)	4.3% (±0.7)	4.2% (±0.6)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=9,030)	% (± CI) (n=10,566)	% (± CI) (n=8,729)	% (± CI) (n=6,561)
215. I feel safe at my school.				
a. NO! ^{AC} /Definitely NOT true ^B	3.5% (±0.5)	4.4% (±0.7)	4.4% (±0.8)	4.0% (±1.0)
b. no ^{AC} /Mostly not true ^B	7.7% (±0.7)	9.5% (±1.1)	11.0% (±1.8)	9.0% (±1.5)
c. yes ^{AC} /Mostly true ^B	46.4% (±1.6)	51.8% (±2.3)	56.8% (±1.8)	54.0% (±2.7)
d. YES! ^{AC} /Definitely true ^B	42.4% (±1.9)	34.3% (±3.2)	27.8% (±3.6)	33.1% (±4.3)
216. My teachers praise me when I work hard in school.	(n=8,801)	(n=5,201)	(n=4,248)	(n=3,213)
a. NO!	10.8% (±0.9)	14.7% (±1.7)	16.7% (±1.4)	13.6% (±1.6)
b. No	31.6% (±1.6)	35.6% (±2.1)	43.5% (±2.1)	42.4% (±2.1)
c. yes	45.2% (±1.6)	38.5% (±1.8)	32.3% (±2.0)	36.1% (±2.6)
d. YES!	12.3% (±1.0)	11.3% (±1.3)	7.4% (±1.0)	7.9% (±1.1)

Peer and Individual Domain

Perceived Risk of Drug Use (Questions 217-220)

How much do you think people risk harming themselves if they:

217. Smoke one or more packs of cigarettes per day?	(n=8,400)	(n=5,200)	(n=4,234)	(n=3,184)
a. No risk	6.7% (±1.0)	2.1% (±0.5)	2.4% (±0.4)	2.7% (±0.7)
b. Slight risk	5.4% (±0.6)	3.6% (±0.6)	3.6% (±0.6)	3.3% (±0.7)
c. Moderate risk	14.8% (±0.8)	13.2% (±1.2)	12.7% (±1.2)	13.2% (±1.3)
d. Great risk	61.0% (±2.3)	75.2% (±2.6)	78.5% (±1.8)	78.9% (±2.1)
e. Not sure	12.1% (±1.2)	5.9% (±1.2)	2.7% (±0.5)	1.9% (±0.5)
218. Try marijuana once or twice?	(n=8,313)	(n=5,193)	(n=4,225)	(n=3,182)
a. No risk	12.4% (±1.2)	15.6% (±1.9)	31.9% (±2.5)	43.1% (±2.5)
b. Slight risk	20.1% (±1.1)	24.1% (±1.4)	28.9% (±1.7)	28.7% (±1.6)
c. Moderate risk	24.1% (±1.2)	27.6% (±1.9)	20.4% (±1.5)	14.0% (±1.4)
d. Great risk	30.2% (±1.2)	27.1% (±2.2)	15.8% (±1.6)	12.1% (±1.3)
e. Not sure	13.2% (±1.0)	5.7% (±0.9)	3.0% (±0.6)	2.1% (±0.6)
219. Use marijuana regularly (at least once or twice a week)?	(n=8,297)	(n=5,188)	(n=4,225)	(n=3,176)
a. No risk	10.9% (±1.3)	8.7% (±1.6)	15.4% (±2.1)	21.9% (±2.5)
b. Slight risk	8.1% (±0.8)	10.9% (±1.5)	18.6% (±1.0)	23.1% (±1.7)
c. Moderate risk	18.7% (±1.0)	21.5% (±1.4)	27.4% (±1.9)	26.6% (±1.9)
d. Great risk	49.6% (±2.5)	53.2% (±3.8)	35.6% (±2.8)	25.7% (±2.1)
e. Not sure	12.6% (±1.0)	5.8% (±0.9)	3.0% (±0.6)	2.6% (±0.8)
220. Take one or two drinks of an alcoholic beverage (wine, beer, a shot, liquor) nearly every day?	(n=8,310)	(n=5,180)	(n=4,219)	(n=3,178)
a. No risk	13.4% (±1.2)	6.6% (±0.7)	5.3% (±1.0)	7.0% (±1.2)
b. Slight risk	19.1% (±1.4)	14.8% (±1.3)	15.4% (±1.6)	18.2% (±1.7)
c. Moderate risk	23.9% (±1.0)	30.3% (±1.9)	32.7% (±1.7)	34.5% (±1.7)
d. Great risk	32.3% (±1.7)	42.9% (±2.7)	43.8% (±2.8)	38.4% (±2.7)
e. Not sure	11.3% (±1.0)	5.4% (±0.9)	2.8% (±0.5)	2.0% (±0.5)

Early Initiation of Drug Use (Questions 221-224)

How old were you the first time you:

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=10,348)	% (± CI) (n=8,563)	% (± CI) (n=6,423)
221. Used marijuana?				
a. Never have	**	89.6% (±1.6)	70.6% (±2.5)	54.3% (±3.2)
b. 10 or younger	**	1.9% (±0.4)	2.2% (±0.4)	1.9% (±0.5)
c. 11	**	1.4% (±0.4)	1.8% (±0.4)	1.3% (±0.3)
d. 12	**	2.8% (±0.6)	3.4% (±0.6)	3.0% (±0.7)
e. 13	**	3.4% (±0.5)	6.7% (±0.8)	6.0% (±0.7)
f. 14	**	0.6% (±0.2)	8.1% (±0.7)	8.3% (±1.2)
g. 15	**	0.1% (±0.1)	6.4% (±0.8)	9.6% (±1.0)
h. 16	**	0.1% (±0.0)	0.6% (±0.2)	9.7% (±1.0)
i. 17 or older	**	0.1% (±0.1)	0.2% (±0.1)	6.0% (±0.6)
222. Smoked a cigarette, even just a puff?	(n=0)	(n=5,068)	(n=4,144)	(n=3,119)
a. Never have	**	88.2% (±1.7)	78.0% (±2.6)	68.5% (±3.1)
b. 10 or younger	**	4.3% (±1.0)	4.8% (±1.0)	4.2% (±1.0)
c. 11	**	1.7% (±0.4)	2.1% (±0.5)	1.9% (±0.5)
d. 12	**	2.6% (±0.5)	2.7% (±0.5)	2.8% (±0.8)
e. 13	**	2.3% (±0.5)	3.6% (±0.8)	4.5% (±1.0)
f. 14	**	0.4% (±0.2)	4.4% (±0.8)	3.7% (±0.7)
g. 15	**	0.1% (±0.1)	3.5% (±0.7)	4.1% (±0.8)
h. 16	**	0.2% (±0.1)	0.6% (±0.3)	5.5% (±0.9)
i. 17 or older	**	0.1% (±0.1)	0.3% (±0.2)	4.6% (±0.8)
223. Had more than a sip or two of beer, wine, or hard liquor (for example vodka, whiskey, or gin)?	(n=0)	(n=10,348)	(n=8,579)	(n=6,423)
a. Never have	**	71.0% (±1.9)	49.9% (±2.1)	33.8% (±2.2)
b. 10 or younger	**	12.3% (±1.1)	10.6% (±0.9)	8.0% (±0.9)
c. 11	**	4.3% (±0.4)	3.4% (±0.4)	2.5% (±0.5)
d. 12	**	5.4% (±0.5)	5.5% (±0.5)	4.4% (±0.6)
e. 13	**	5.9% (±0.5)	8.0% (±0.7)	7.2% (±0.8)
f. 14	**	0.9% (±0.2)	11.2% (±0.7)	9.8% (±0.9)
g. 15	**	0.1% (±0.1)	10.1% (±1.0)	12.5% (±1.0)
h. 16	**	0.0% (±0.0)	1.0% (±0.2)	12.8% (±1.1)
i. 17 or older	**	0.1% (±0.1)	0.2% (±0.1)	8.8% (±0.8)
224. Began drinking alcoholic beverages regularly, that is, at least once or twice a month?	(n=0)	(n=5,051)	(n=4,114)	(n=3,086)
a. Never have	**	93.4% (±1.1)	82.9% (±1.8)	69.6% (±2.5)
b. 10 or younger	**	1.4% (±0.4)	1.2% (±0.4)	1.3% (±0.4)
c. 11	**	0.6% (±0.2)	0.8% (±0.2)	0.9% (±0.4)
d. 12	**	1.8% (±0.4)	1.5% (±0.5)	1.3% (±0.5)
e. 13	**	2.0% (±0.4)	2.1% (±0.5)	1.5% (±0.5)
f. 14	**	0.4% (±0.2)	4.3% (±0.7)	2.4% (±0.6)
g. 15	**	0.1% (±0.1)	5.6% (±1.0)	4.8% (±1.0)
h. 16	**	0.2% (±0.1)	1.2% (±0.4)	9.5% (±1.2)
i. 17 or older	**	0.1% (±0.1)	0.3% (±0.2)	8.8% (±1.3)

Favorable Attitudes Toward Drug Use (Questions 225-228)

How wrong do YOU think it is for someone your age to:

	Grade 6	Grade 8	Grade 10	Grade 12
225. Drink beer, wine, or hard liquor (for example vodka, whiskey, or gin) regularly?	% (± CI) (n=8,448)	% (± CI) (n=4,910)	% (± CI) (n=4,045)	% (± CI) (n=3,053)
a. Very wrong	86.2% (±1.1)	69.3% (±2.3)	41.5% (±2.8)	28.8% (±2.6)
b. Wrong	10.1% (±0.8)	20.2% (±1.6)	30.8% (±1.8)	30.2% (±1.9)
c. A little bit wrong	2.8% (±0.4)	8.0% (±1.1)	20.8% (±1.5)	29.6% (±2.6)
d. Not wrong at all	0.9% (±0.2)	2.5% (±0.5)	6.9% (±1.0)	11.4% (±2.0)
226. Smoke cigarettes?	(n=8,414)	(n=4,911)	(n=4,042)	(n=3,048)
a. Very wrong	88.5% (±1.0)	75.6% (±2.1)	59.4% (±2.4)	50.6% (±2.9)
b. Wrong	9.3% (±0.8)	17.5% (±1.5)	25.4% (±1.5)	24.6% (±1.8)
c. A little bit wrong	1.6% (±0.3)	4.9% (±0.8)	10.7% (±1.1)	15.2% (±1.2)
d. Not wrong at all	0.6% (±0.2)	2.0% (±0.5)	4.5% (±1.0)	9.6% (±1.8)
227. Use marijuana?	(n=8,380)	(n=4,874)	(n=4,023)	(n=3,042)
a. Very wrong	90.9% (±1.0)	72.4% (±2.6)	41.1% (±2.7)	28.8% (±2.8)
b. Wrong	6.3% (±0.7)	13.5% (±1.2)	20.0% (±1.2)	20.7% (±1.6)
c. A little bit wrong	1.8% (±0.4)	8.7% (±1.3)	21.3% (±1.5)	24.9% (±1.9)
d. Not wrong at all	1.0% (±0.3)	5.4% (±1.2)	17.6% (±1.8)	25.5% (±2.6)
228. Use LSD, cocaine, amphetamines, or another illegal drug?	(n=8,369)	(n=4,864)	(n=4,015)	(n=3,038)
a. Very wrong	94.6% (±0.9)	86.5% (±1.4)	76.5% (±1.8)	71.0% (±2.3)
b. Wrong	4.2% (±0.8)	9.8% (±1.0)	15.8% (±1.3)	18.6% (±1.5)
c. A little bit wrong	0.7% (±0.2)	2.2% (±0.4)	5.4% (±0.8)	6.8% (±1.2)
d. Not wrong at all	0.6% (±0.2)	1.5% (±0.4)	2.3% (±0.5)	3.6% (±0.7)

Friends Use Drugs (Questions 229-232)

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have...

	(n=0)	(n=4,258)	(n=3,694)	(n=2,873)
229. Smoked cigarettes?				
a. None of my friends	**	86.8% (±2.0)	73.8% (±3.0)	66.4% (±2.5)
b. 1 of my friends	**	7.1% (±1.1)	12.3% (±1.7)	15.2% (±1.3)
c. 2 of my friends	**	2.7% (±0.6)	6.3% (±1.0)	7.7% (±1.2)
d. 3 of my friends	**	1.3% (±0.4)	3.4% (±0.8)	4.5% (±0.8)
e. 4 of my friends	**	2.0% (±0.5)	4.2% (±0.8)	6.2% (±1.2)
230. Tried beer, wine, or hard liquor (for example vodka, whiskey, or gin) when their parents didn't know about it?	(n=0)	(n=4,254)	(n=3,685)	(n=2,866)
a. None of my friends	**	80.0% (±2.3)	51.0% (±2.7)	35.2% (±2.7)
b. 1 of my friends	**	9.9% (±1.1)	16.7% (±1.3)	17.4% (±1.4)
c. 2 of my friends	**	4.6% (±0.8)	11.3% (±1.2)	13.7% (±1.3)
d. 3 of my friends	**	2.4% (±0.6)	8.1% (±1.1)	11.2% (±1.3)
e. 4 of my friends	**	3.1% (±0.7)	12.8% (±1.7)	22.4% (±2.7)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,252)	% (± CI) (n=3,691)	% (± CI) (n=2,871)
231. Used marijuana?				
a. None of my friends	**	82.8% (±2.6)	54.7% (±3.1)	42.7% (±3.1)
b. 1 of my friends	**	7.4% (±1.2)	15.8% (±1.6)	17.1% (±1.4)
c. 2 of my friends	**	3.9% (±0.9)	9.8% (±1.1)	12.2% (±1.3)
d. 3 of my friends	**	2.0% (±0.6)	7.0% (±0.8)	9.6% (±1.3)
e. 4 of my friends	**	4.0% (±0.9)	12.7% (±1.6)	18.5% (±2.2)
232. Used LSD, cocaine, amphetamines, or other illegal drugs?	(n=0)	(n=4,243)	(n=3,684)	(n=2,868)
a. None of my friends	**	93.9% (±1.1)	87.4% (±1.5)	82.4% (±2.1)
b. 1 of my friends	**	3.1% (±0.6)	6.9% (±1.0)	8.5% (±1.1)
c. 2 of my friends	**	1.0% (±0.3)	2.5% (±0.6)	4.0% (±0.9)
d. 3 of my friends	**	0.8% (±0.3)	1.2% (±0.3)	2.0% (±0.6)
e. 4 of my friends	**	1.2% (±0.4)	2.0% (±0.5)	3.0% (±0.8)
Interaction with Prosocial Peers (Questions 233-237)				
<i>Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have...</i>				
233. Participated in clubs, organizations or activities at school?	(n=0)	(n=4,371)	(n=3,744)	(n=2,889)
a. None of my friends	**	13.9% (±1.5)	14.2% (±1.8)	15.7% (±2.2)
b. 1 of my friends	**	12.0% (±1.2)	12.6% (±1.6)	13.1% (±1.6)
c. 2 of my friends	**	14.8% (±1.2)	14.7% (±1.3)	14.4% (±1.3)
d. 3 of my friends	**	13.0% (±1.1)	13.4% (±1.2)	12.6% (±1.2)
e. 4 of my friends	**	46.4% (±2.5)	45.2% (±3.4)	44.1% (±3.9)
234. Made a commitment to stay drug-free?	(n=0)	(n=4,291)	(n=3,716)	(n=2,879)
a. None of my friends	**	24.1% (±2.3)	28.5% (±2.4)	37.5% (±2.8)
b. 1 of my friends	**	7.4% (±0.9)	13.7% (±1.4)	15.7% (±1.9)
c. 2 of my friends	**	5.8% (±0.8)	10.3% (±1.1)	10.5% (±1.1)
d. 3 of my friends	**	6.5% (±0.8)	10.6% (±0.8)	9.0% (±1.2)
e. 4 of my friends	**	56.2% (±3.2)	36.9% (±3.0)	27.2% (±2.8)
235. Liked school?	(n=0)	(n=4,294)	(n=3,718)	(n=2,876)
a. None of my friends	**	24.2% (±2.0)	28.7% (±2.5)	30.4% (±2.8)
b. 1 of my friends	**	12.3% (±1.1)	14.6% (±1.3)	14.3% (±1.0)
c. 2 of my friends	**	15.5% (±1.1)	17.4% (±1.0)	16.9% (±1.6)
d. 3 of my friends	**	15.1% (±1.1)	14.5% (±1.2)	12.6% (±1.5)
e. 4 of my friends	**	32.9% (±2.4)	24.7% (±2.3)	25.9% (±2.4)
236. Regularly attended religious services?	(n=0)	(n=4,220)	(n=3,698)	(n=2,867)
a. None of my friends	**	30.4% (±2.6)	33.8% (±2.4)	37.4% (±2.8)
b. 1 of my friends	**	22.7% (±1.3)	26.3% (±1.8)	25.7% (±1.7)
c. 2 of my friends	**	18.8% (±1.2)	17.1% (±1.3)	16.8% (±1.6)
d. 3 of my friends	**	11.2% (±1.2)	8.5% (±1.3)	8.4% (±1.4)
e. 4 of my friends	**	17.0% (±1.6)	14.2% (±2.1)	11.8% (±1.9)

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=0)	% (± CI) (n=4,260)	% (± CI) (n=3,707)	% (± CI) (n=2,862)
237. Tried to do well in school?				
a. None of my friends	**	7.0% (±1.1)	6.2% (±1.2)	6.5% (±1.1)
b. 1 of my friends	**	5.8% (±0.9)	6.7% (±0.9)	7.3% (±0.9)
c. 2 of my friends	**	7.4% (±0.9)	10.4% (±1.4)	11.8% (±1.4)
d. 3 of my friends	**	12.8% (±1.1)	16.5% (±1.2)	15.9% (±1.4)
e. 4 of my friends	**	66.9% (±2.6)	60.2% (±3.3)	58.5% (±2.8)

Belief in the Moral Order (Questions 238-241)

238. I think it is okay to take something without asking as long as you get away with it.	(n=0)	(n=5,172)	(n=4,198)	(n=3,158)
a. NO!	**	53.8% (±2.3)	44.6% (±1.8)	48.5% (±1.9)
b. No	**	36.1% (±1.5)	44.0% (±1.9)	42.5% (±1.7)
c. yes	**	7.5% (±1.0)	9.0% (±0.8)	6.7% (±1.0)
d. YES!	**	2.6% (±0.6)	2.4% (±0.6)	2.2% (±0.5)
239. I think sometimes it's okay to cheat at school.	(n=0)	(n=5,170)	(n=4,196)	(n=3,159)
a. NO!	**	52.2% (±2.5)	30.2% (±2.2)	25.4% (±1.9)
b. No	**	33.2% (±1.4)	38.4% (±1.8)	38.8% (±1.7)
c. yes	**	12.5% (±1.6)	27.6% (±2.3)	30.5% (±1.9)
d. YES!	**	2.0% (±0.5)	3.8% (±0.7)	5.4% (±1.0)
240. It is all right to beat up people if they start the fight.	(n=0)	(n=5,154)	(n=4,195)	(n=3,159)
a. NO!	**	38.0% (±2.6)	25.5% (±1.9)	24.5% (±2.2)
b. No	**	26.5% (±1.1)	28.4% (±1.7)	28.2% (±1.6)
c. yes	**	22.9% (±1.9)	29.0% (±1.5)	30.9% (±1.8)
d. YES!	**	12.7% (±1.5)	17.1% (±1.9)	16.4% (±1.7)
241. It is important to be honest with your parents, even if they become upset or you get punished.	(n=0)	(n=5,152)	(n=4,181)	(n=3,149)
a. NO!	**	6.9% (±0.8)	5.7% (±0.8)	5.8% (±0.9)
b. No	**	10.4% (±1.0)	13.6% (±1.3)	15.0% (±1.4)
c. yes	**	38.1% (±1.8)	49.4% (±1.6)	49.4% (±1.7)
d. YES!	**	44.7% (±2.1)	31.3% (±1.8)	29.8% (±1.9)

Prosocial Involvement (Questions 242-244)

How many times in the past year (12 months) have you...

242. Participated in clubs, organizations or activities at school?	(n=8,926)	(n=0)	(n=0)	(n=0)
a. Never	25.4% (±2.1)	**	**	**
b. 1 or 2 times	31.0% (±1.1)	**	**	**
c. 3 to 5 times	17.4% (±0.9)	**	**	**
d. 6 to 9 times	8.2% (±0.7)	**	**	**
e. 10 to 19 times	5.9% (±0.6)	**	**	**
f. 20 to 29 times	3.2% (±0.4)	**	**	**
g. 30 to 39 times	1.9% (±0.5)	**	**	**
h. 40+ times	6.9% (±0.8)	**	**	**

	Grade 6	Grade 8	Grade 10	Grade 12
	% (± CI) (n=8,777)	% (± CI) (n=0)	% (± CI) (n=0)	% (± CI) (n=0)
243. Done extra work on your own for school?				
a. Never	23.1% (±1.7)	**	**	**
b. 1 or 2 times	32.7% (±1.2)	**	**	**
c. 3 to 5 times	16.8% (±0.8)	**	**	**
d. 6 to 9 times	9.7% (±0.9)	**	**	**
e. 10 to 19 times	6.8% (±0.7)	**	**	**
f. 20 to 29 times	3.9% (±0.5)	**	**	**
g. 30 to 39 times	1.8% (±0.3)	**	**	**
h. 40+ times	5.1% (±0.6)	**	**	**
244. Volunteered to do community service?	(n=8,816)	(n=0)	(n=0)	(n=0)
a. Never	54.9% (±2.4)	**	**	**
b. 1 or 2 times	23.3% (±1.3)	**	**	**
c. 3 to 5 times	9.4% (±0.7)	**	**	**
d. 6 to 9 times	5.0% (±0.6)	**	**	**
e. 10 to 19 times	2.9% (±0.4)	**	**	**
f. 20 to 29 times	1.5% (±0.3)	**	**	**
g. 30 to 39 times	0.7% (±0.2)	**	**	**
h. 40+ times	2.4% (±0.4)	**	**	**

Social Skills (Questions 245-247)

245. You're looking at CDs in a music store with a friend. You look up and see her slip a CD under her coat. She smiles and says, "Which one do you want? Go ahead, take it while nobody's around." There is nobody in sight, no employees, and no other customers.

	(n=0)	(n=4,186)	(n=3,654)	(n=2,844)
a. Ignore her	**	15.2% (±1.7)	19.7% (±2.1)	23.1% (±1.8)
b. Grab a CD and leave the store	**	6.3% (±1.1)	9.3% (±1.0)	7.9% (±1.4)
c. Tell her to put the CD back	**	47.3% (±2.1)	39.0% (±2.1)	37.9% (±2.1)
d. Act like it's a joke and ask her to put the CD back	**	31.2% (±1.3)	32.0% (±1.6)	31.0% (±1.7)

246. You are visiting another part of town and you don't know any of the people your age there. You are walking down the street and some teenager you don't know is walking toward you. He is about your size. As he is about to pass you, he deliberately bumps int

	(n=0)	(n=4,206)	(n=3,676)	(n=2,846)
a. Push the person back	**	10.5% (±1.3)	10.9% (±1.3)	11.0% (±1.4)
b. Say nothing and keep on walking	**	47.7% (±1.9)	48.5% (±2.2)	46.9% (±3.3)
c. Say, "Watch where you're going," and keep on walking	**	30.7% (±1.5)	28.2% (±2.0)	30.2% (±2.4)
d. Swear at the person and walk away	**	11.1% (±1.4)	12.4% (±1.2)	12.0% (±1.4)

247. You are at a party at someone's house and one of your friends offers you a drink containing alcohol. What would you say or do?

	(n=0)	(n=4,200)	(n=3,665)	(n=2,842)
a. Drink it	**	11.6% (±1.7)	28.0% (±2.2)	37.5% (±2.7)
b. Tell your friend, "No thanks. I don't drink," and suggest that you and your friend go and do something else	**	48.5% (±2.1)	35.9% (±2.0)	28.2% (±2.2)
c. Just say, "No, thanks," and walk away	**	23.5% (±1.5)	25.9% (±1.6)	27.5% (±1.8)
d. Make up a good excuse, tell your friend you had something else to do, and leave	**	16.3% (±1.6)	10.2% (±1.3)	6.8% (±1.1)

Questions by Topics

Questions about a single topic are not always grouped together in this report. For example, some questions about alcohol are presented in the Alcohol, Tobacco, and Other Drug (ATOD) Use section and others are presented in the Risk and Protective Factor section. The following list, groups the questions by topic so you can easily locate them in this report.

Abuse 122-126

Activities and Clubs 87, 175-177, 233, 242

Alcohol

Access 46, 164
 Age at first use 223-224
 Consequences 54
 Current use 32, 43-44
 Drinking and driving 111-113
 Family influence 47-48, 185-186, 196
 Lifetime use 17
 Perceived norms 50-51, 53, 169, 171, 225, 230, 247
 Perceived risks 52, 220
 Prevention/intervention 62, 150
 Use at school 49, 61
 Usual type 45

Asthma 88-93

Bullying/Harassment 143-149

Demographics 1-9

Family

Food insecurity 68-69
 Living situations 10-12
 Military service 13-14
 Parent's education 7-8

Fighting 119, 240

Fitness

Physical activity 79-82, 85-86
 Sedentary behavior 83-84

Gambling 142

Gangs 120-121

HIV/STD/Pregnancy Prevention 154-156

Health or Health Care

Diabetes 98
 Doctor visit 94
 Oral Health 95-97
 Sleep 99

Honesty 15, 238-239, 241

Marijuana

Access 56, 166
 Age at first use 221
 Current use 33
 Family influence 57-58, 198
 Lifetime use 18
 Perceived norms 60, 168, 173, 227, 231
 Perceived risks 218-219
 Use at school 59, 61
 Using and driving 114-115
 Usual type 55

Mental Health

Anxiety and worry 133-134
 Depression 127
 Help with mental health 135-136
 Suicide 128-132

Nutrition

Breakfast 71
 Food insecurity 68-69
 Fruit/Vegetable servings 70
 Junk food 77-78
 Soda and Sugar Drinks 72-76

Other Drugs

Access 167
 Consequences 54
 Current use 34-37
 Family influence 63, 185
 Lifetime use 19-24
 Perceived norms 64, 228, 232, 234
 Perceived risks 65, 218-219
 Prevention/intervention 62, 150
 Use at school 61

Quality of Life 157-163

Safety-related Behaviors

Swimming 106-107
 Wearing Safety Gear 105

School Climate

Absence 188, 207
 Achievement 199-200
 After-school activities 87, 175-177, 233, 242
 Bullying/harassment 143-149
 Counseling and support 151-153
 Engagement 201-206, 208-214, 216, 235, 237
 Nutrition 73-74, 76-78
 Physical activity 81-82, 85-86
 Prevention/intervention 62, 136, 150
 Safety at 121, 148, 215
 Substance use at 49, 59, 61
 Weapons at 116-117

Sexual Behaviors 100-104

Sexual Orientation 9

Social and Emotional Learning 137-141

Substance Use Prevention 62, 150

Texting and Driving 108-110

Tobacco

Access 165
 Age at first use 222
 Current use 25-31
 Family 40-41, 197
 Lifetime use 16
 Perceived norms 42, 170, 226, 229
 Perceived risks 217
 Prevention/intervention 150
 Secondhand Smoke 38-39

Weapons 116-118, 172, 187

Weight and Weight Loss 66-67

Core Questions

"Core" questions appeared on both Forms A and B (the secondary versions of the survey) are listed below. All other questions appeared on either Form A or Form B, but not both. Core questions may or may not have appeared on Form C (the elementary version).

Activities

Participation in after school activities 87

Alcohol

Age when first drank alcohol 223
Binge drinking 43
Current alcohol drinking (past 30 days) 32
Levels of alcohol use 44
Lifetime alcohol use 17

Bullying

Been bullied (past 30 days) 143

Demographics

Age 1
Education status of father 8
Education status of mother 7
Food insecurity 69
Gender 3
Language spoken in home 5
Loss of home due to family finances 12
Race/Ethnicity 4
Where you live 11
Who you live with 10

Fighting

Physical fighting (past 12 months) 119

Gangs

Gang involvement 120
Gangs at school 121

Honesty

Honesty in completing survey 15

Marijuana

Age when first used marijuana 221
Current marijuana use (past 30 days) 33
Lifetime marijuana use 18

Mental Health

Bothered by anxiety 133
Bothered by worrying 134
Depression (past 12 months) 127
Seriously considered attempting suicide 128

Other Drugs

Current drug use including marijuana (past 30 days) 35
Current drug use not including marijuana (past 30 days) 34
Current painkiller use (past 30 days) 36

School Climate

Been bullied (past 30 days) 143
Enjoyed being in school (past 12 months) 204
Feel safe at school 215
Grades in school (past 12 months) 199
Participation in after school activities 87
Weapon carrying on school property (past 12 months) 116

Texting and Driving

Texting and driving (past 30 days) 110

Tobacco

Current cigarette smoking (past 30 days) 25

Weapons

Weapon carrying on school property (past 12 months) 116

For more information on which survey forms each question was on,
see the Survey Crosswalk at www.AskHYS.net, under HYS Results - QxQ